

**International Nonproprietary Names (INN)
for biological and biotechnological substances**
(a review)



**World Health
Organization**

**International Nonproprietary Names (INN)
for biological and biotechnological substances**
(a review)



International Nonproprietary Names (INN) Programme

**Technologies Standards and Norms (TSN)
Regulation of Medicines and other Health Technologies (RHT)
Essential Medicines and Health Products (EMP)**

International Nonproprietary Names (INN) for biological and biotechnological substances

(a review)

FORMER DOCUMENT NUMBER: INN Working Document 05.179

© World Health Organization 2014

All rights reserved. Publications of the World Health Organization are available on the WHO website (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications –whether for sale or for non-commercial distribution– should be addressed to WHO Press through the WHO website (www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed by the WHO Document Production Services, Geneva, Switzerland

CONTENTS

0.	INTRODUCTION.....	iv
1.	PHARMACOLOGICAL CLASSIFICATION OF BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES	1
2.	CURRENT STATUS OF EXISTING STEMS OR SYSTEMS FOR BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES	4
2.1.	Groups with respective stems	4
2.2.	Groups with respective pre-stems.....	5
2.3.	Groups with INN schemes.....	5
2.4.	Groups without respective stems / pre-stems and without INN schemes.....	5
3.	GENERAL POLICIES FOR BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES	6
3.1.	General policies for blood products	6
3.2.	General policies for fusion proteins	6
3.3.	General policies for gene therapy products	6
3.4.	General policies for glycosylated compounds	7
3.5.	General policies for immunoglobulins fractionated from plasma	8
3.6.	General policies for monoclonal antibodies	8
3.7.	General policies for non-glycosylated compounds	11
3.8.	General policies for skin substitutes	11
3.9.	General policies for transgenic products	11

3.10.	General policies for vaccines	12
3.11.	General policies for cell therapy products	12
4.	SUMMARY OF INN ASSIGNED TO BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES	13
4.1.	Antimicrobial, bactericidal permeability increasing polypeptides	13
4.2.	Antisense oligonucleotides	13
4.3.	Antithrombins	13
4.4.	Blood coagulation cascade inhibitors	14
4.5.	Blood coagulation factors	14
4.6.	Colony stimulating factors	15
4.7.	Enzymes	16
4.8.	Erythropoietin type blood factors	20
4.9.	Gene therapy products	21
4.10.	Gonadotropin-releasing-hormone (GnRH) inhibitors, peptides	21
4.11.	Growth factors	21
4.12.	Growth hormone (GH) derivatives	23
4.13.	Growth hormone antagonists	23
4.14.	Heparin derivatives including low molecular mass heparins	23
4.15.	Hirudin derivatives.....	24
4.16.	Insulins	24
4.17.	Interferons	25
4.18.	Interleukin receptor antagonists	26
4.19.	Interleukin type substances	26

4.20.	Monoclonal antibodies.....	27
4.21.	Oxytocin derivatives	30
4.22.	Peptides and glycopeptides.....	30
4.23.	Peptide vaccines / recombinant vaccines.....	33
4.24.	Pituitary / placental glycoprotein hormones	34
4.25.	Pituitary hormone-release stimulating peptides.....	35
4.26.	Receptor molecules, native or modified	35
4.27.	Synthetic polypeptides with a corticotropin-like action	36
4.28.	Thrombomodulins.....	36
4.29.	Toxins	37
4.30.	Vasoconstrictors, vasopressin derivatives	37
4.31.	Various.....	37
5.	CURRENT CHALLENGES	46
	REFERENCES	47
	ANNEX 1.....	49
	The list of INN for composite proteins published	49
	ANNEX 2.....	69
	Transliteration of Greek letters in English, French and Spanish	69
	ANNEX 3.....	70
	The previous naming scheme for monoclonal antibodies.....	70

0. INTRODUCTION

More than 50 years ago, WHO established the International Nonproprietary Name (INN) Expert Group / WHO Expert Committee on Specifications for Pharmaceutical Preparations, to assign nonproprietary names to medicinal substances, so that each substance would be recognized globally by a unique name. These INNs do not give proprietary rights, unlike a trade mark, and can be used freely as they are public property.

INNs have been assigned to biological products since the early days of the INN Programme. As well as many names for individual substances, animal insulin preparations were given an INN in Recommended list 3 in 1959. In the period up to 1980, names were assigned to antibiotics, synthetic peptides, hormones and other proteins. In names of compounds related by structure and / or function, specific letter groups, called stems, are included to aid recognition by health professionals. The *-actide* for synthetic polypeptides with a corticotrophin-like action is an example.

In 1982, the name *insulin human* was proposed for the recombinant protein identical to natural human insulin, and since then names have been assigned to a growing number of recombinant products. Within the INN Programme, names have not been assigned to natural human blood products or vaccines. For those groups of biological products, the WHO Expert Committee on Biological Standardization (ECBS) has been adopting the scientific names of the biological products within the definitions of respective requirements.

Since the time when *insulin human* became the first recommended INN (rINN) for a recombinant product, the range of biological / biotechnological products has increased in size and complexity. For example, new stems have been introduced for tissue plasminogen activators (*-plase*) among other groups. Analogues of recombinant glycosylated proteins produced in different cell systems have been classified using Greek letters as indicators in the sequence of product introduction: erythropoietin (*epoetin alfa, beta* and so on) and glycoprotein hormones (*follitropin*) are examples. In the 1990s, a systematic scheme for naming monoclonal antibodies was implemented, based on the stem *-mab*, which indicates the origin (mouse, human, etc) of the antibody and its intended use: tumour, immunomodulator and so on.

As a result of the scientific and technical developments currently taking place, new products of biotechnology and other biological products are being introduced and more products can be expected for the treatment or prevention of disease. Examples of such new products include recombinant blood products, transgenic products (human proteins expressed in animals or plants), products for gene therapy and novel vaccines.

As this area is becoming more and more complex and challenging, the INN Expert Group has requested the WHO-INN Secretariat to prepare a working document intended to summarize and review the past and present INN situation in this field.

This document, first published on the website of the INN Programme in 2006, therefore presents an inventory of the policy decisions taken by the INN Expert Group during all these years of change, and of the names assigned to biological and biotechnological substances. Considering the potential for further developments in the field of biologicals, this review is intended to be a *living document* which will be regularly updated to include new policies, and future INNs assigned.

Comments and suggestions from all interested parties are most welcome and will be presented to the INN Expert Group for their consideration and for possible incorporation in future updates of this review.

You are reading the current updated version, also available as pdf-copy at:

<http://www.who.int/medicines/services/inn/publication/en/index.html>.

1. PHARMACOLOGICAL CLASSIFICATION OF BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES ⁽¹⁾

Alimentary tract and metabolism

insulins (see item 4.16).

Anti-infectives

antimicrobial, bactericidal permeability increasing polypeptides (see item 4.1).

Antineoplastics

peptide vaccines / recombinant vaccines (see item 4.23)

toxins (see item 4.29).

Blood and agents acting on the haemopoietic system

antithrombins (see item 4.3)

blood coagulation cascade inhibitors (see item 4.4)

blood coagulation factors (see item 4.5)

erythropoietin type blood factors (see item 4.8)

heparin derivatives including low molecular mass heparins (see item 4.14)

hirudin derivatives (see item 4.15)

thrombomodulins (see item 4.28).

Immunomodulators and immunostimulants

- colony stimulating factors (see item 4.6)
- interferons (see item 4.17)
- interleukin receptor antagonists (see item 4.18)
- interleukin type substances (see item 4.19)
- monoclonal antibodies (see item 4.20)
- receptor molecules, native or modified (see item 4.26).

Hormones, hormone antagonists, hormone-release stimulating peptides or hormone-release inhibiting peptides (excluding insulins)

- gonadotropin-releasing-hormone (GnRH) inhibitors, peptides (see item 4.10)
- growth hormone (GH) derivatives (see item 4.12)
- growth hormone antagonists (see item 4.13)
- oxytocin derivatives (see item 4.21)
- pituitary / placental glycoprotein hormones (see item 4.24)
- pituitary hormone-release stimulating peptides (see item 4.25)
- synthetic polypeptides with a corticotropin-like action (see item 4.27)
- vasoconstrictors, vasopressin derivatives (see item 4.30).

Various

- antisense oligonucleotides (see item 4.2)
- enzymes (see item 4.7)
- gene therapy products (see item 4.9)

growth factors (see item 4.11)

peptides and glycopeptides (for special groups of peptides see *-actide* (see item 4.27), *-pressin* (see item 4.30), *-relin* (see item 4.25), *-tocin* (see item 4.21)) (see item 4.22).

2. CURRENT STATUS OF EXISTING STEMS OR SYSTEMS FOR BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES^{(1) (2) (3) (4) (5) (6) (7)}

2.1. Groups with respective stems

Name of the group	Stem
antisense oligonucleotides	<i>-rsen</i>
blood coagulation cascade inhibitors	<i>-cogin</i>
blood coagulation factors	<i>-cog</i>
colony stimulating factors	<i>-stim</i>
enzymes	<i>-ase</i>
erythropoietin type blood factors	<i>-poetin</i>
growth factors	<i>-ermin</i>
growth hormone derivatives	<i>som-</i>
heparin derivatives including low molecular mass heparins	<i>-parin</i>
hirudin derivatives	<i>-irudin</i>
gonadotropin-releasing-hormone (GnRH) inhibitors, peptides	<i>-relix</i>
interleukin receptor antagonists	<i>-kinra</i>
interleukin type substances	<i>-kin</i>
monoclonal antibodies	<i>-mab</i>
oxytocin derivatives	<i>-tocin</i>
peptides and glycopeptides (for special groups of peptides see <i>-actide</i> , <i>-pressin</i> , <i>-relin</i> , <i>-tocin</i>)	<i>-tide</i>
pituitary hormone-release stimulating peptides	<i>-relin</i>
receptor molecules, native or modified (a preceding infix should designate the target)	<i>-cept</i>
synthetic polypeptides with a corticotropin-like action	<i>-actide</i>
vasoconstrictors, vasopressin derivatives	<i>-pressin</i>

2.2. Groups with respective pre-stems

Name of the group	Pre-stem
aptamers, classical and mirror ones	<i>-apt-</i>
antimicrobial, bactericidal permeability increasing polypeptides	<i>-ganan</i>
neurotrophins	<i>-neurin</i>
small interfering RNA	<i>-siran-</i>

2.3. Groups with INN schemes

Name of the group
antithrombins
gene therapy products
insulins
interferons
pituitary / placental glycoprotein hormones

2.4. Groups without respective stems / pre-stems and without INN schemes

Name of the group
growth hormone antagonists
thrombomodulins
toxins

3. GENERAL POLICIES FOR BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES

3.1. General policies for blood products⁽⁴⁾

- INNs have not been assigned to natural human blood products.
- Many natural blood products have well-established names, so the recombinant version should have a distinctive name reflecting as much as possible the established name used in the field.
- It is essential to add "activated" to the name of the blood product when this is presented for therapeutic use in its activated form.

3.2. General policies for fusion proteins¹⁽⁴⁾

- INNs have been assigned to some fusion proteins. If a stem exists for one or the other part of the fusion protein, this stem should be brought into the name. This allows the constant part of a fusion protein to be recognized in the name.
- At present it is considered unnecessary to indicate that the product is a fusion product within the name, but this position may need to be reviewed in the future.

3.3. General policies for gene therapy products⁽²⁾

In 2005, the two-word nomenclature scheme for gene therapy products was formally adopted by the members of the INN Expert Group designated to deal with the selection of nonproprietary names. The 2012 updated scheme is shown in Table 1.

¹ The list of INN for composite proteins published is given in Annex 1, including some fusion proteins.

Table 1 Two-word scheme for gene therapy products (updated in 2012)

	prefix	infix	suffix
word 1 (gene component)	random to contribute to euphonious and distinctive name e.g. <i>al-</i> ; <i>bet-</i> ; <i>val-</i>	to identify the gene using, when available, existing infixes for biological products or using similar infix as for the protein for which the gene codes. e.g. <i>-cima-</i> : cytosine deaminase <i>-ermin-</i> : growth factor <i>-kin-</i> : interleukin <i>-lim-</i> : immunomodulator <i>-lip-</i> : human lipoprotein lipase <i>-mul-</i> : multiple gene <i>-stim-</i> : colony stimulating factor <i>-tima-</i> : thymidine kinase <i>-tusu-</i> : tumour suppression	<i>-(a vowel)gene</i> e.g. <i>-(o)gene</i>
word 2 (vector component)	random to contribute to euphonious and distinctive name	e.g. <i>-adeno-</i> : adenovirus <i>-cana-</i> : canarypox virus <i>-foli-</i> : fowlpox virus <i>-herpa-</i> : herpes virus <i>-lenti-</i> : lentivirus <i>-morbilli-</i> : paramyxoviridae morbillivirus <i>-parvo-</i> : adeno-associated virus (parvoviridae dependovirus) <i>-retro-</i> : other retrovirus <i>-vaci-</i> : vaccinia virus	<i>-vec</i> (non-replicating viral vector) <i>-repvec</i> (replicating viral vector)
			<i>-plasmid</i> (plasmid vector)

In the case of non-plasmid naked DNA products, there is no need for a second word in the name.

3.4. General policies for glycosylated compounds ⁽⁸⁾

For glycoproteins / glycopeptides

- For groups identified with a stem, e.g. for erythropoetins: *-poetin*, differences in the amino acid chain are indicated by using a

random prefix and differences in the glycosylation pattern are indicated by another designator expressed by a Greek letter² spelt in full and added as a second word to the name (e.g. *epoetin alfa* (66), *darbepoetin alfa* (85); see item 4.8).

- For blood coagulation factors obtained by recombinant biotechnology, the differences in the glycosylation pattern are indicated by a Greek letter spelt in full and added as a second word to the name (e.g. *eptacog alfa (activated)* (77), *octocog alfa* (73)); when the amino acid sequence differs from the natural product this is indicated by using a random prefix (e.g. *beroctocog alfa* (98); see item 4.5).
- Similarly, for enzymes identified with a stem *-ase* obtained by recombinant biotechnology and differing in the amino acid chain, these differences are indicated by using a random prefix and differences in the glycosylation pattern are indicated by a Greek letter spelt in full and added as a second word to the name (e.g. *alglucosidase alfa* (91), *bucelipase alfa* (95); see item 4.7).
- The Greek letters should be used in the Greek alphabetical order (see Annex 2).

3.5. **General policies for immunoglobulins fractionated from plasma**^{(9) (10)}

Not to select an INN for immunoglobulins fractionated from plasma is the current policy.

The "systematic" or descriptive name is essential since the prescriber must know all the information conveyed by it and there is no benefit in assigning an INN from which it will not be readily apparent.

3.6. **General policies for monoclonal antibodies**^{(1) (3) (11)³}

- INN for monoclonal antibodies (mAbs) are composed of a prefix, a substem A, a substem B and a suffix.
- The common stem for mAbs is *-mab*, placed as a suffix.

² The transliteration of Greek letters in English, French and Spanish is given in Annex 2.

³ It contains the revised naming scheme for monoclonal antibodies; the previous naming scheme for monoclonal antibodies is given in Annex 3.

- The stem *-mab* is to be used for all products containing an immunoglobulin variable domain which binds to a defined target.
- Substem B indicates the species on which the immunoglobulin sequence of the mAb is based (shown in Table 2).

Table 2 Substem B for the species

<i>a</i>	rat
<i>axo (pre-sub-stem)</i>	rat-mouse
<i>e</i>	hamster
<i>i</i>	primate
<i>o</i>	mouse
<i>u</i>	human
<i>xi</i>	chimeric
<i>-xizu-</i>	chimeric-humanized
<i>zu</i>	humanized

The distinction between chimeric and humanized antibodies is as follows:

Chimeric: A chimeric antibody is one for which both chain types are chimeric as a result of antibody engineering. A chimeric chain is a chain that contains a foreign variable domain (originating from one species other than human, or synthetic or engineered from any species including human) linked to a constant region of human origin. The variable domain of a chimeric chain has a V region amino acid sequence which, analysed as a whole, is closer to non-human species than to human.

Humanized: A humanized antibody is one for which both chain types are humanized as a result of antibody engineering. A humanized chain is typically a chain in which the complementarity determining regions (CDR) of the variable domains are foreign (originating from one species other than human, or synthetic) whereas the remainder of the chain is of human origin. Humanization assessment is based on the resulting amino acid sequence, and not on the methodology per se, which allows protocols other than grafting to be used. The variable domain of a humanized chain has a V region amino acid sequence which, analysed as a whole, is closer to human than to other species.

The *-xizu-* infix is used for an antibody having both chimeric and humanized chains.

The *-axo-* infix is used for an antibody having both rat and mouse chains.

- Substem A indicates the target (molecule, cell, organ) class (shown in Table 3).

Table 3 Substem A for target class

<i>-b(a)-</i>	bacterial
<i>-c(i)-</i>	cardiovascular
<i>-f(u)-</i>	fungal
<i>-gr(o)-</i>	skeletal muscle mass related growth factors and receptors
<i>-k(i)-</i>	interleukin
<i>-l(i)-</i>	immunomodulating
<i>-n(e)-</i>	neural
<i>-s(o)-</i>	bone
<i>-tox(a)-</i>	toxin
<i>-t(u)-</i>	tumour
<i>-v(i)-</i>	viral

In principle, a single letter, e.g. *-b-* for bacterial is used as substem A. Whenever substem B starts with a consonant (e.g. *x* or *z*), to avoid problems in pronunciation, an additional vowel indicated in the table, e.g. *-ba-* is inserted.

Prefix

The prefix should be random, i.e. the only requirement is to contribute to a euphonious and distinctive name.

Second word

If the monoclonal antibody is conjugated to another protein or to a chemical (e.g. chelator), identification of this conjugate is accomplished by use of a separate, second word or acceptable chemical designation. For instance, for mAbs conjugated to a toxin, the suffix *-tox* is used in the second word.

If the monoclonal antibody is radiolabelled, the radioisotope is listed first in the INN, e.g. *technetium (^{99m}Tc) nofetumomab merpentan (81)*.

The prefix *peg-* can be used for pegylated mAbs, but this should be avoided if it leads to over-long INN. In most cases, it is best to adopt two-word INN for pegylated mAbs, with the first word describing the mAb and the second being *pegol* or a related designation.

3.7. General policies for non-glycosylated compounds⁽⁸⁾

For proteins / peptides:

- Identification of the group with a stem, e.g. for hirudin analogues: *-irudin*, and indication of differences in the amino acid chain by using a random prefix (e.g. *bivalirudin (72)*).
- Identification of the group with a word, e.g. insulin, and indication of differences in the composition of the amino acid chain as a second element of the name (e.g. *insulin argine (58)*).

3.8. General policies for skin substitutes⁽⁴⁾

The products within this system are made of cells within a matrix, and skin substitutes can be considered to be engineered tissue and thus fall outside the scope of the INN system.

3.9. General policies for transgenic products⁽⁴⁾

- If an INN already exists, the same name should be used for the transgenic product, qualified in some way to identify that this product is transgenic.
- A similar system to that used for glycosylated recombinant products is suggested to differentiate new or additional sources of the same substance, and the source of the substance should be included in the definition of the INN.

3.10. General policies for vaccines^{(4) (5) (6) (7)}

- At present, vaccines are not included within the INN system, but names are assigned through recommendations of the Expert Committee on Biological Standardization and through the pharmacopoeial monograph.
- During the INN Consultation in 1993, it was agreed that the prerequisite for an INN application for a recombinant vaccine⁴ would be fulfilled if the manufacturer was able to provide all information outlined in the guidelines entitled Definition of INNs for Substances Prepared by Biotechnology (WHO / Pharm S / Nom 1348⁽¹²⁾).
- During the INN Consultation in 1998, following discussion on recombinant viruses, the experts agreed not to attempt to name live viruses.
- Another approach in vaccine technology is the development of peptide vaccines⁵ (epitopes involved in immune response formation): since these peptides are chemically well-defined, they fall within the INN naming system.

3.11. General policies for cell therapy products

During the 55th INN Consultation in 2012, the INN Expert Group decided in principle to name some cell therapy products. A possible INN naming scheme for cell therapy products is currently under development.

⁴ The definition of recombinant vaccines is given in item 4.23.

⁵ The definition of peptide vaccines is given in item 4.23.

4. SUMMARY OF INN ASSIGNED TO BIOLOGICAL AND BIOTECHNOLOGICAL SUBSTANCES^{(1) (3) (7) (8) (13) (14) (15) (16) (17) (18) (19) (20) (21)}

4.1. Antimicrobial, bactericidal permeability increasing polypeptides

The pre-stem for antimicrobial, bactericidal permeability increasing polypeptides is *-ganan*.

iseganan (85)⁶, *omiganan* (89), *pexiganan* (78).

4.2. Antisense oligonucleotides

The common stem for antisense oligonucleotides is *-rsen*.

aganirsen (103), *alicaforsen* (97), *anivamersen* (105), *apatorsen* (110), *aprinocarsen* (97), *beclanorsen* (101), *cenersen* (97), *custirsen* (99), *drisapersen* (106), *etepirsen* (103), *gataparsen* (103), *mipomersen* (100), *mongersen* (111), *oblimersen* (97), *trabedersen* (98).

-virsen (antivirals): *afovirsen* (97), *fomivirsen* (97), *miravirsen* (101), *radavirsen* (106), *trecovirsen* (97).

4.3. Antithrombins

antithrombin III (60), *antithrombin alfa* (93) (Rec. Glycoprotein (432aa) from transgenic goats), *antithrombin gamma* (107).

⁶ The numbers in parentheses indicate the Proposed list number.

4.4. Blood coagulation cascade inhibitors

The common stem for blood coagulation cascade inhibitors is *-cogin*.

drotrecogin alfa (activated) (86), pegnivacogin (106), taneptacogin alfa (90), tifacogin (78).

4.5. Blood coagulation factors

The common stem for blood coagulation factors is *-cog*.

The sub-stems *-eptacog*, *-octocog*, *-nonacog/-trenonacog* and *-tridecacog* have been selected up to date for recombinant blood coagulation factors.

A prefix will be necessary if the amino acid sequence does not match that of the naturally occurring material.

In accordance with the general policy, *alfa*, *beta*, etc, will be added for the glycoproteins (see item 3.4 - general policies for glycosylated compounds).

When the additional statement "activated" is needed, e.g. for the blood coagulation factor VIIa, it should be spelt out in full and added in parentheses after the name.

blood coagulation factor VII: *-eptacog*

eptacog alfa (activated) (77), eptacog alfa pegol (activated) (101), oreptacog alfa (activated) (109), vatreptacog alfa (activated) (98)

blood factor VIII: *-octocog*

beroctocog alfa (98), damoctocog alfa pegol (109), efmoroctocog alfa (111), lonoctocog alfa (111), moroectocog alfa (72), octocog alfa (73), ruriococog alfa pegol (111), simoectocog alfa (104), turoctocog alfa (108), turoctocog alfa pegol (108)

blood factor IX: *-nonacog* (with Ala at the position 148 (Ala-alloform))

albutrepenonacog alfa (109), nonacog alfa (77), nonacog beta pegol (104), nonacog gamma (108)

-*trenonacog* (with Thr at the position 148 (Thr-alloform))

eftrenonacog alfa (109), *trenonacog alfa* (107)

blood coagulation factor XIII: -*tridecacog*

catridecacog (99)

recombinant von Willebrand factor (vWF): -*vonico*

vonico alfa (102).

4.6. Colony stimulating factors

The common stem for colony stimulating factors is -*stim*.

ancestim (79) (cell growth factor), *garnocestim* (86) (immunomodulator), *pegacaristim* (80) (megakaryocyte growth factor), *romiplostim* (97) (thrombopoietin receptor (MPL) agonist)

combination of two different types of colony stimulating factors: -*distim*

leridistim (80), *milodistim* (75)

granulocyte macrophage colony stimulating factor (GM-CSF) type substances: -*gramostim*

ecogramostim (62), *molgramostim* (64), *regramostim* (65), *sargramostim* (66)

granulocyte colony stimulating factor (G-CSF) type substances: -*grastim*

balugrastim (107), *eplapegrastim* (111), *empegfilgrastim* (107), *filgrastim* (64), *lenograstim* (64), *lipegfilgrastim* (107), *nartograstim* (66), *pegbovigrastim* (109), *pegfilgrastim* (86), *pegnartograstim* (80), *pegteograstim* (109)

macrophage stimulating factors (M-CSF) type substances: -*mostim*

cilmostim (71), *lanimostim* (91), *mirimostim* (65)

interleukin-3 analogues and derivatives: *-plestim*

daniplestim (76), muplestim (74).

4.7. Enzymes

The common stem for enzymes, in general, is *-ase*.

Sub-stems are referring to the activity of the substances.

proteinase:

with *-ase* suffix:

crisantaspase (111), brinase (22), calaspargase pegol (105), kallidinogenase (22), ocrase (28), pegaspargase (64), promelase (47), serrapeptase (31), sfericase (40), streptokinase (6), urokinase (48), urokinase alfa (77)

without *-ase* suffix:

batroxobin (29), bromelains (18), chymopapain (26), chymotrypsin (10), defibrotide (44), fibrinolysin (human) (10), sutilains (18)

lipase: *-lipase*

bucelipase alfa (95), burlulipase (107), rizolipase (22), sebelipase alfa (107)

enzymes with superoxide dismutase activity: *-dismase*

- *ledismase (70), sudismase (58)*
- isomerase (belongs to this group but in which the preferred stem has not been used)

orgotein (31), pegorgotein (72)

plasminogen activator combined with another enzyme: *-diplase*

amediplase (79)

tissue-type-plasminogen activators: *-teplase*

*alteplase (73), desmoteplase (80), dutiplase (62), lanoteplase (76),
monteplase (72), nateplase (73), pamiteplase (78), reteplase (69), silteplase
(65), tenecteplase (79)*

anistreplase (59) (belongs to this group but in which the preferred stem has not been used)

urokinase-type-plasminogen activators: *-uplase*

nasaruplase (76), nasaruplase beta (86), saruplase (76)

others:

agalsidase alfa (84): human alpha-galactosidase isoenzyme A, isolated from human cell line, clone RAG 001, glycoform α

agalsidase beta (84): α -galactosidase (human clone λ AG¹⁸ isoenzyme A subunit protein moiety reduced), glycoform β

alfimeprase (85): [3-L-serine]fibrolase-(3-203)-peptide (fibrolase : fibrinolytic enzyme isolated from *Agkistrodon contrix contrix* venom)

alglucerase (68): glucosylceramidase (human placenta isoenzyme protein moiety reduced)

alglucosidase alfa (91): human lysosomal prepro- α -glucosidase-(57-952)-peptide 199-arginine-223-histidine variant

asfotase alfa (104): tissue-nonspecific alkaline phosphatase- IgG₁ fusion protein; human tissue-nonspecific isozyme alkaline phosphatase (AP-TNAP, EC=3.1.3.1) fusion protein with leucyl-lysyl-human immunoglobulin G1 Fc region {(6-15)-H-CH₂-CH₃ of IGHG1*03} fusion protein with aspartyl-isoleucyl-deca(aspartic acid), dimer (493-493':496-496')-bisdisulfide

cerliponase alfa (111): immature human tripeptidyl-peptidase 1 (cell growth-inhibiting gene 1 protein, lysosomal pepstatin-insensitive protease, TPP-1, EC 3.4.14.9), 544 residues protein, produced in Chinese hamster ovary (CHO) cells, glycoform alfa

condoliase (106): endolyase, chondroitin ABC (C-ABC).
glycosaminoglycan lyase chondroitin ABC endolyase 1 (chondroitinase ABC) *Proteus vulgaris*

dornase alfa (70): deoxyribonuclease (human clone 18-1 protein moiety)

elosulfase alfa (108): human *N*-acetylgalactosamine-6-sulfatase (chondroitinsulfatase, galactose-6-sulfate sulfatase, EC=3.1.6.4) dimer (139-139')-disulfide glycosylated (produced by CHO cells)

epafipase (85): 2-acetyl-1-alkyl-*sn*-glycero-3-phosphocholine deacetylase-(6-400)-peptide(human)

eufausease (84): broad spectrum serine-protease enzyme, extracted from the Antarctic krill (*Euphausia superba*)

galsulfase (92): *N*-acetylgalactosamine 4-sulfatase (human CSL4S-342 cell)

glucarpidase (92): recombinant glutamate carboxypeptidase (carboxypeptidase G2)

hyalosidase (50): hyaluronoglucosaminidase *or* E.C. 3.2.1.35

hyaluronidase (1): enzymes of various origins which depolymerize hyaluronic acid

idursulfase (90): α -L-iduronate sulfate sulfatase

idursulfase beta (106): iduronate 2-sulfatase (α -L-iduronate sulfate sulfatase), human proenzyme produced in CHO cells (glycoform beta)

imiglucerase (72): 495-L-histidineglucosylceramidase (human placenta isoenzyme protein moiety)

laronidase (86): 8-L-histidine- α -L-iduronidase (human)

olipudase alfa (111): recombinant DNA derived des-(1-13)-human sphingomyelin phosphodiesterase (acid sphingomyelinase, EC-3.1.4.12), produced in Chinese hamster ovary (CHO) cells, glycoform alfa

pegademase (63): adenosine deaminase, reaction product with succinic anhydride, esters with polyethylene glycol monomethyl ether
The source of the product should be indicated

pegadricase (105): pegylated Urate Oxidase from *Candida utilis*, [198-threonine(S>T)]uricase (EC 1.7.3.3, urate oxidase) *Pichia jadinii* (Yeast) (*Candida utilis*) tetramer, 6-amino group of an average of 3 lysine residues, mostly in position 16, 19, and 85 of each monomer, are amidified with α -(3-carboxypropanoyl)- ω -methoxypoly(oxyethylene)

pegargininase (111): [111-glutamic acid,209-serine]arginine deiminase (ADI, arginine dihydrolase, AD) from *Mycoplasma hominis*, an average of five amino groups are amidified with 4-[ω -methoxypoly(oxyethylene)]-4-oxobutanoyl, produced in *Escherichia coli*

pegcrisantaspase (111): recombinant L-asparaginase derived from *Erwinia chrysanthemi* pegylated with 5 kDa methoxy polyethylene glycol (m-PEG-NHS), produced in *Escherichia coli*:

L-asparaginase (EC 3.5.1.1, L-asparagine amidohydrolase) *Erwinia chrysanthemi* tetramer α_4 , an average of 10 (*a*) out of 18 amino groups of each monomer are amidified with 5- $\{[\alpha$ -methylpoly(oxyethylene)]amino $\}$ -5-oxopentanoyl

pegloticase (98): tetramer α_4 of des-(1-5)-[6-threonine,45-threonine, 290-lysine, 300-serine]uricase (EC 1.7.3.3, urate oxidase) from *Sus scrofa* (porcine), non acetylated, of which some of the lysine 6-amine residues are engaged in a carbamate linkage with a monomethyl ether of polyoxyethylene (macrogol)

pegvaliase (111): pegylated, recombinant DNA derived *Anabaena variabilis* phenylalanine ammonia lyase mutein (S 503, S 565), produced in *Escherichia coli*:

[503,565-diserine (C>S)]phenylalanine ammonia-lyase (EC 4.3.1.24) *Anabaena variabilis* in which an average of 5 lysyl residues are N6- $\{6$ -[ω -methoxypoly(oxyethylene)]hexanoyl $\}$ substituted

penicillinase (111): an enzyme obtained by fermentation from cultures of *Bacillus Cereus*

ranpirnase (81): ribonuclease (*Rana pipiens*)

rasburicase (82): urate oxydase (tetramer of the *N*-acetylpolypeptide of 301 amino acids

reveglucosidase alfa (111): des-(2-7)-human insulin-like growth factor II fusion protein with glycyl-L-alanyl- L-prolyl-human lysosomal alpha-glucosidase (acid maltase, aglucosidase alfa) produced in Chinese hamster ovary (CHO) cells, glycoform alfa

senrebotase (107): L-methionylglycyl-L-seryl-des-(445-glycine,446-L-tyrosine)-[2-L-glutamic acid,432,442,444,447-tetra-L-aspartic acid]botulinum neurotoxin A precursor 27-L-alanine variant light chain (433-41')-disulfide with [14-L-arginine,15-L-lysine]human nociceptin fusion protein with L-alanyl-L-leucyl-L-alanyltris(tetraglycyl-L-seryl)-[3-L-valine,4-L-leucine,5-L-glutamine-418-L-leucine,419-L-aspartic acid]botulinum neurotoxin A heavy chain-(1-419)-peptide

streptodornase (6): enzyme obtained from cultures of various strains of *Streptococcus hemolyticus* and capable of hydrolysing desoxyribonucleoproteins

taliglucerase alfa (101): L-glutamyl-L-phenylalanyl-[495(497)-L-histidine(R>H)]human glucosylceramidase (beta-glucocerebrosidase) peptide with L-aspartyl-L-leucyl-L-leucyl-L-valyl-L-aspartyl-L-threonyl-L-methionine, glycosylated peptide 1-506

tilactase (50): β -D-galactosidase or EC 3.2.1.23

velaglucerase alfa (98): human glucosylceramidase (EC 3.2.1.45 or beta-glucocerebrosidase), glycoform α .

vonapanitase (111): recombinant DNA derived type I pancreatic elastase, produced in *Pichia pastoris*: [26-tryptophan(Arg>Trp),202-leucine(Val>Leu),225-arginine(Gln>Arg)]mature human CELA1 (chymotrypsin-like elastase family member 1, pancreatic elastase 1, elastase 1, EC 3.4.21.36)non-glycosylated

vorhyaluronidase alfa (111): human hyaluronidase PH-20 (hyaluronoglucosaminidase PH-20, sperm adhesion molecule 1, EC 3.2.1.35) precursor-(36-482)-peptide (mature-(1-447)-peptide), produced in Chinese hamster ovary (CHO) DG44dhfr- cells, glycoform α .

4.8. Erythropoietin type blood factors

The common stem for erythropoietin type blood factors is *-poetin*.

In the case of erythropoietins, it was decided to select *epoetin* together with a Greek letter to differentiate between compounds of the same amino acid sequence as human erythropoietin which vary in the glycosylation pattern (see item 3.4 - general policies for glycosylated compounds).

Substances with different amino acid sequences will be named using the *-poetin* stem and a random prefix.

darbepoetin alfa (85), epoetin alfa (66), epoetin beta (62), epoetin gamma (67), epoetin delta (85), epoetin epsilon (72), epoetin zeta (95), epoetin theta (95), epoetin kappa (97), epoetin omega (73).

4.9. Gene therapy products

alferminogene tadenovec (95), alipogene tiparvovec (99), amolimogene bepiplasmid (98), beperminogene perplasmid (95), contusugene ladenovec (97), golnerminogene pradenovec (101), pexastimogene devacirepvec (108), riferminogene pecaplastmid (100), rilimogene galvacirepvec (107), rilimogene glafolivec (107), sitimagene ceradenovec (97), taberminogene vadenovec (100), talimogene laherparepvec (104), tipapkinogene sovacivec (102), velimogene aliplasmid (97), vocimagene amiretrorepvec (107).

4.10. Gonadotropin-releasing-hormone (GnRH) inhibitors, peptides

The common stem for gonadotropin-releasing-hormone (GnRH) inhibitors, peptides is *-relix*.

abarelix (78), cetrotorelix (66), degarelix (86), detirelix (56), ganirelix (65), iturelix (79), ozarelix (94), prazarelix (81), ramorelix (69), teverelix (78).

4.11. Growth factors

The common stem for growth factors is *-ermin*.

Sub-stems allow distinction between the various types of growth factors.

INNs for tumour necrosis factors (TNF) are also classified under the stem *-ermin*.

vascular endothelial growth factors: *-bermin*

telbermin (85)

epidermal growth factors: *-dermin*

murodermin (63), nepidermin (97)

fibroblast growth factors: *-fermin*

ersofermin (66), palifermin (88), repifermin (82), sprifermin (105), trafermin (74), velafermin (94)

leukaemia-inhibiting factors: *-filermin*

emfilermin (82)

tumour necrosis factors: *-nermin*

ardenermin (88), dulanermin (99), plusonermin (73), sonermin (68), tasonermin (78)

platelet-derived growth factors: *-plermin*

becaplermin (74)

insulin-like growth factors: *-sermin*

mecasermin (66), mecasermin rinfabate (92)

transforming growth factors: *-termin*

cetermin (74), liatermin (81)

bone morphogenetic proteins: *-otermin*

avotermin (77), dibotermin alfa (89), eptotermin alfa (92), nebotermin (109), radotermin (92)

others:

cimaglermin alfa (110) (recombinant DNA derived glial growth factor 2 (GGF2))

dapiclermin (93) (modified ciliary neurotrophic factor (CNTF)).

4.12. Growth hormone (GH) derivatives

The common stem for growth hormone derivatives is *som-*.

human growth hormone derivatives:

somatrem (54), *somatropin* (74), *somatropin pegol* (103)

For substances other than human, suffixes are added to indicate the species specificity of the structure.

bovine-type substances: *-bove*

somagrebove (63), *somavubove* (63), *sometribove* (74), *somidobove* (58)

porcine-type substances: *-por*

somalapor (62), *somenopor* (62), *somfasepor* (66), *sometripor* (75)

salmon-type substances: *-salm*

somatosalm (69)

others (growth hormone related peptides):

somatorelin (57) (pituitary hormone-release stimulating peptides, see item 4.25), *somatostatin* (46) (growth hormone release inhibitor).

4.13. Growth hormone antagonists

pegvisomant (82).

4.14. Heparin derivatives including low molecular mass heparins

The common stem for heparin derivatives including low molecular mass heparins is *-parin*.

ardeparin sodium (68), adomiparin sodium (104), bemiparin sodium (75), certoparin sodium (70), dalteparin sodium (77), deligoparin sodium (89), enoxaparin sodium (77), heparin sodium (54), livaraparin calcium (86), minolteparin sodium (74), nadroparin calcium (78), parnaparin sodium (77), reviparin sodium (78), semuloparin sodium (99), sevuparin sodium (106), tafoxiparin sodium (102), tinzaparin sodium (77).

4.15. Hirudin derivatives

The common stem for hirudin derivatives is *-irudin*.

bivalirudin (72), desirudin (76), lepirudin (76), pegmusirudin (77).

4.16. Insulins

Up to now, the insulin derivatives have been named using the two-word approach. The compounds named represent a structure with an additional amino acid, such as *insulin argine (58)*, or represent modifications of the amino acid sequence, i.e. *insulin aspart (76)*.

biphasic insulin injection (16), compound insulin zinc suspension (06), dalanated insulin (104), globin zinc insulin injection (06), insulin argine (58), insulin aspart (76), insulin defalan (37), insulin degludec (101), insulin detemir (80), insulin glargine (76), insulin glulisine (84), insulin human (48), insulin lispro (72), insulin peglispro (107), insulin tregopil (103), insulin zinc suspension (amorphous) (04), insulin zinc suspension (crystalline) (04), isophane insulin (04), neutral insulin injection (15), protamine zinc insulin injection (06)

argine: B30-yl-L-arginyl-L-arginine

aspart: [B28-L-aspartic acid]

dalanated: des-B30-alanine

defalan: des-B1-phenylalanine

degludec: N⁶, B²⁹-[N-(15-carboxypentadecanoyl)-L-γ-glutamyl]-des-30B-L-threonine

detemir: N^{6,B29}-tetradecanoyl-des-B30-L-threonine

glargine: [A21-glycine], B30-yl-L-arginyl-L-arginine

glulisine: [B3-lysine, B29-glutamic acid]

lispro: [B28-L-lysine, B29-L-proline]

tregopil: N^{6,B29}-(4,7,10,13-tetraoxatetradecanoyl).

4.17. Interferons

Interferon was published as an INN in 1962 with a general definition based on the origin and activity, e.g. "a protein formed by the interaction of animal cells with viruses capable of conferring on animal cells resistance to virus infection".

The name was revised in the 1980s when human interferon and its variations *alfa*, *beta* and *gamma* were produced by recombinant biotechnology. The INN Expert Group would have preferred to replace the old INN interferon by alfaferon, betaferon and gammaferon; however, this approach could not be adopted as these names had already been registered as trade marks. The system adopted was thus to take interferon alfa, interferon beta and interferon gamma, and to provide, when necessary, for further distinction by additional numbers, or in the case of mixtures, by additional codes. Additional Arabic numbers can be used to distinguish subspecies which differ significantly in primary amino acid sequence, but are still considered to belong to one of the primary groups e.g. Interferon alfa-1, interferon alfa- 2. Small (lower case) letters are used to subdivide such groups further on the basis of less significant differences like one, two or three amino acid differences or post translational modifications, including glycosylation e.g. Interferon alfa-2a, Interferon alfa-2b, Interferon beta-1a, Interferon beta-1b.

albinterferon alfa-2b (99), *cepeginterferon alfa-2b* (105), *interferon alfa* (73), *interferon alfacon-1* (77), *interferon beta* (73), *interferon gamma* (73), *peginterferon alfa-2a* (84), *peginterferon alfa-2b* (84), *peginterferon beta-1a* (108), *peginterferon lambda-1a* (105), *ropeginterferon alfa-2b* (109).

4.18. Interleukin receptor antagonists

The common stem for interleukin receptor antagonists is *-kinra*.

interleukin-1 (IL-1) receptor antagonists: *-nakinra*

anakinra (72)

interleukin-4 (IL-4) receptor antagonists: *-trakinra*

pitrakinra (87).

4.19. Interleukin type substances

The common stem for interleukin type substances is *-kin*.

In accordance with general policy for naming glycosylated proteins (see item 3.4), it was agreed to publish the INNs for glycosylated interleukins with alfa, beta.

interleukin-1 (IL-1) analogues and derivatives: *-nakin*

interleukin-1 α analogues and derivatives: *-onakin*

pifonakin (77)

interleukin-1 β analogues and derivatives: *-benakin*

mobenakin (72)

interleukin-2 (IL-2) analogues and derivatives: *-leukin*

adargileukin alfa (89), *aldesleukin* (63), *celmoleukin* (65), *denileukin diftitox* (78), *pegaldesleukin* (74), *teceleukin* (67), *tucotuzumab celmoleukin* (95)

interleukin-3 (IL-3) analogues and derivatives: *-plestim* (belongs to this group but in which the preferred stem has not been used)

daniplestim (76), *muplestim* (74)

interleukin-4 (IL-4) analogues and derivatives: *-trakin*

binetrakin (82)

interleukin-6 (IL-6) analogues and derivatives: *-exakin*

atexakin alfa (72)

interleukin-8 (IL-8) analogues and derivatives: *-octakin*

canoctakin (110), *emoctakin* (74)

interleukin-10 (IL-10) analogues and derivatives: *-decakin*

ilodecakin (81)

interleukin-11 (IL-11) analogues and derivatives: *-elvekin*

oprelvekin (76)

interleukin-12 (IL-12) analogues and derivatives: *-dodekin*

edodekin alfa (79)

interleukin-13 (IL-13) analogues and derivatives: *-tredekin*

cintredekin besudotox (92)

interleukin-18 (IL-18) analogues and derivatives: *-octadekin*

iboctadekin (92)

interleukin-21 (IL-21) analogues and derivatives: *-enicokin*

denenicokin (99)

neurotrophins (interleukin-78, brain derived neurotropic factor): *-neurin*
(pre-stem, belongs to this group but in which the preferred stem has not
been used)

abrineurin (84).

4.20. Monoclonal antibodies

The common stem for monoclonal antibodies is *-mab*.

INNs for monoclonal antibodies alphabetically by origin:

-axomab (pre-sub-stem, rat-murine hybrid)

catumaxomab (93), ertumaxomab (93)

-omab (mouse origin)

abagovomab (95), afelimomab (80), altumomab (80), anatumomab mafenatox (86), arcitumomab (74), bectumomab (81), begelomab (111), besilesomab (92), biciromab (66), blinatumomab (100), capromab (80), detumomab (80), dorlimomab aritox (66), edobacomab (80), edrecolomab (74), elsilimomab (89), enlimomab (80), enlimomab pegol (77), epitumomab (97), epitumomab cituxetan (89), faralimomab (81), gavilimomab (84), ibritumomab tiuxetan (86), igovomab (86), imciromab (66), inolimomab (80), lemalesomab (86), maslimomab (66), minretumomab (80), mitumomab (82), moxetumomab pasudotox (102), nacolomab tafenatox (80), naptumomab estafenatox (96), nerelimomab (81), odulimomab (81), oregovomab (86), racotumomab (100), satumomab (81), solitomab (106), sulesomab (86), taplitumomab paptox (84), technetium (^{99m}Tc) fanolesomab (86), technetium (^{99m}Tc) nofetumomab merpentan (81), technetium (^{99m}Tc) pintumomab (86), telimomab aritox (66), tenatumomab (99), tositumomab (80), vepalimomab (80), zolimomab aritox (80)

-umab (human origin)

abrilumab (111), actoxumab (111), adalimumab (85), adecatumumab (90), aducanumab (110), alirocumab (107), anatumab ravtansine (109), anifrolumab (109), atinumab (104), atorolimumab (80), belimumab (89), bertilimumab (88), bezlotoxumab (107), bimagrumab (111), briakinumab (101), brodalumab (105), canakinumab (97), carlumab (104), cixutumumab (100), conatumumab (99), daratumumab (101), denosumab (94), diridavumab (111), drozitumab (103), dupilumab (108), dusigitumab (108), efungumab (95), eldelumab (109), enfortumab vedotin (109), enoticumab (107), evolocumab (108), exbivirumab (91), fasinumab (107), fezakinumab (101), figitumumab (100), firivumab (111), flavotumab (106), fletikumab (110), foralumab (103), foravirumab (100), fresolimumab (101), fulranumab (104), ganitumab (103), gantenerumab (108), glembatumumab (102), golimumab (91), guselkumab (109), icrucumab (104), imalumab (111), inclacumab (106), intetumumab (101), ipilimumab (94), iratumumab (94), lenzilumab (111), lerdelimumab (86), lexatumumab (95), libivirumab (91), lirilumab (107), lucatumumab (98), mapatumumab (93), mavrilimumab (102), metelimumab (88), morolimumab (79), namilumab (104), narnatumab (105), nebacumab (66), necitumumab (100), nesvacumab (108), nivolumab (111), ofatumumab (93), olaratumab (103), orticumab (107), oxelumab (105), panitumumab (96), panobacumab (100), patritumab (106),

placulumab (107), primumab (89), radretumab (104), rafivirumab (100), ramucirumab (110), raxibacumab (92), regavirumab (80), rilatumumab (101), robatumumab (100), roledumab (103), sarilumab (106), secukinumab (102), seribantumab (108), sevirumab (66), sifalimumab (104), sirukumab (105), stamulumab (95), tabalumab (105), tarextumab (109), teprotumumab (108), tosatoxumab (109), tovetumab (109), tralokinumab (102), tremelimumab (97), tuvirumab (66), ulocuplumab (110), urelumab (104), ustekinumab (99), vantictumab (109), varlilumab (111), vesencumab (104), votumumab (80), zalutumumab (93), zanolimumab (92), ziralimumab (84)

-ximab (chimeric origin)

abciximab (80), amatuximab (104), basiliximab (81), bavituximab (95), brentuximab vedotin (103), cetuximab (82), clenoliximab (77), coltuximab ravtansine (109), dinutuximab (109), ecomeximab (87), ensituximab (103), futuximab (107), galiximab (89), girentuximab (101), indatuximab ravtansine (105), infliximab (77), iodine (¹²⁴I) girentuximab (101), keliximab (81), lumiliximab (90), margetuximab (109), modotuximab (110), pagibaximab (93), priliximab (80), pritoxaximab (108), rituximab (77), setoxaximab (108), siltuximab (100), teneliximab (87), ublituximab (104), vapaliximab (87), volociximab (93)

-xizumab (chimeric-humanized origin)

ontuxizumab (109), otelixizumab (99), pasotuxizumab (111)

-zumab (humanized origin)

abitudzumab (109), alacizumab pegol (98), alemtuzumab (83), anrukinzumab (98), apolizumab (87), aselizumab (88), bapineuzumab (93), benralizumab (102), bevacizumab (86), bimekizumab (110), bivatumab (86), blosozumab (105), bococizumab (110), brontictuzumab (111), cantuzumab mertansine (105), cantuzumab ravtansine (105), caplacizumab (106), cedelizumab (81), certolizumab pegol (97), citatuzumab bogatox (99), clazakizumab (107), codrituzumab (109), concizumab (108), crenezumab (105), dacetuzumab (98), daclizumab (78), dalotuzumab (107), dapirolizumab pegol (110), demcizumab (107), denintuzumab mafodotin (111), duligotuzumab (110), eculizumab (87), efalizumab (85), elotuzumab (100), emactuzumab (111), emibetuzumab (111), enavatuzumab (104), enokizumab (104), epratuzumab (82), erlizumab (84), etaracizumab (99), etrolizumab (104), farletuzumab (100), felvizumab (77), ficlatuzumab (105), fontolizumab (87), gemtuzumab (83), gevokizumab (104), ibalizumab (97), idarucizumab (109), imgatuzumab (107), inotuzumab ozogamicin (92), itolizumab (103), ixekizumab (105), labetuzumab (85), lampaizumab (107), lebrikizumab (101), lifastuzumab vedotin (110), ligelizumab (107), lintuzumab (86), lodelcizumab (108), lorvotuzumab mertansine (103),

lulizumab pegol (111), lumretuzumab (111), matuzumab (88), mepolizumab (81), milatuzumab (98), mogamulizumab (104), motavizumab (95), natalizumab (79), nimotuzumab (94), obinutuzumab (109), ocaratuzumab (107), ocrelizumab (95), olokizumab (103), omalizumab (84), onartuzumab (104), oportuzumab monatox (100), otlertuzumab (110), ozanezumab (108), ozoralizumab (105), palivizumab (79), parsatuzumab (107), pascolizumab (87), pateclizumab (105), pembrolizumab (110), perakizumab (108), pertuzumab (89), pexelizumab (86), pidilizumab (108), pinatuzumab vedotin (108), polatuzumab vedotin (110), ponezumab (104), quilizumab (106), ralpancizumab (110), ranibizumab (90), reslizumab (85), romosozumab (106), rontalizumab (101), rovelizumab (81), ruplizumab (83), samalizumab (105), sibrotuzumab (86), simtuzumab (107), sipilizumab (87), sofituzumab vedotin (110), solanezumab (107), sontuzumab (94), suvizumab (102), tadocizumab (94), talizumab (89), tanezumab (99), tefibazumab (92), teplizumab (97), tigatuzumab (98), tiltrakizumab (108), tocilizumab (90), toralizumab (87), trastuzumab (78), trastuzumab emtansine (103), tregalizumab (104), tucotuzumab celmoleukin (95), urtoxazumab (90), vanucizumab (111), vatelizumab (105), vedolizumab (100), veltuzumab (98), visilizumab (84), vorsetuzumab (107), vorsetuzumab mafodotin (107), yttrium (⁹⁰Y) clivatuzumab tetraxetan (102), yttrium ⁹⁰Y tacatuzumab tetraxetan (93)

Others: *muromonab-CD3 (59)* (the first monoclonal antibody to which an INN was assigned belongs to this group but it was named before the stem was established).

4.21. Oxytocin derivatives

The common stem for oxytocin derivatives is *-tocin*.

argiprestocin (13), aspartocin (11), carbetocin (45), carginocin (35), demoxytocin (22), merotocin (111), nacartocin (51), oxytocin (13).

4.22. Peptides and glycopeptides

for special groups of peptides see *-actide* (see item 4.27), *-pressin* (see item 4.30), *-relin* (see item 4.25), *-tocin* (see item 4.21)

The common stem for peptides and glycopeptides is *-tide*.

analgesic: *leconotide (86), ziconotide (78)*

angiogenesis inhibitor: *cilengitide* (81)

angiotensin converting-enzyme inhibitor: *teprotide* (36)

antianaemic: *peginesatide* (108)

antiarrhythmic: *danegaptide* (101), *rotigaptide* (94)

anti-inflammatory: *icrocaptide* (89)

antidepressant: *nemifitide* (87)

antidiabetic: *albenatide* (111), *amlintide* (76), *davalintide* (101), *exenatide* (89), *langlenatide* (111), *lixisenatide* (99), *pramlintide* (74), *seglitide* (57)

antidiarrhoeal: *lagatide* (75)

antineoplastic: *paclitaxel trevatide* (109)

antithrombotic: *eptifibatide* (78) (-fiba- is a pre-substem for platelet aggregation inhibitor (GPIIb/IIIa receptor antagonist))

antiviral: *enfuvirtide* (85), *tifuvirtide* (91)

natriuretic peptides: *anaritide* (57), *carperitide* (65), *cenderitide* (105), *neseritide* (80), *ularitide* (69)

autoimmune disorders: *dalazatide* (111), *dirucotide* (100)

β 1-adrenergic receptor analogue: *mibenratide* (111)

calcium sensing receptor agonist: *velcalcetide* (109)

cicatrisation promoter: *aclerastide* (110), *ensereptide* (107)

diagnostic: *betiatide* (58), *bibapcitide* (78), *ceruletide* (34), *depreotide* (80), *flotegatide* (^{18}F) (108), *fluciclatide* (^{18}F) (103), *maraciclaitide* (103), *mertiatile* (60), *pendetide* (70), *technetium* ($^{99\text{m}}\text{Tc}$) *apcitide* (86), *technetium* ($^{99\text{m}}\text{Tc}$) *etarfolatide* (107), *teriparatide* (50)

expectorant (in cystic fibrosis): *lancovutide* (99)

gastro-intestinal bleeding / antineoplastic: *edotreotide* (84), *ilatretotide* (68), *lanretotide* (64), *octretotide* (52),

pentetreotide (66), vapreotide (62)

gastro-intestinal functions normalizing agent: *linaclotide (97), plecanatide (104)*

glucagon-like peptide (GLP) analogues: *-glutide*

albiglutide (97), dulaglutide (103), elsiglutide (104), liraglutide (87), semaglutide (101), taspoglutide (99), teduglutide (90)

growth stimulant-veterinary: *nosiheptide (35)*

gut motility increasing: *ociltide (52)*

hormone analogue: *abaloparatide (109), semparatide (80), teriparatide (50)*
(see diagnostic)

immunological agents - antineoplastics: *almurtide (74), delmitide (92), edratide (89), goralatide (72), mifamurtide (95), murabutide (49), pentigetide (60), pimelautide (53), prezatide copper acetate (67), rolipoltide (94), romurtide (61), tabilautide (60), temurtide (60), tigapotide (95)*

immunological agents for active immunization: *-motide (see item 4.23)*

abecomotide (109), alicdamotide (109), amilomotide (105), asudemotide (107), disomotide (94), elpamotide (103), latromotide (107), ovemotide (94), pradimotide (107), tanurmotide (109), tecemotide (108), tertomotide (98), tiplimotide (82), trempamotide (107), zastumotide (110)

inhibition of growth hormone release: *pasireotide (90)*

kallicrein inhibitor: *ecallantide (93)*

melanocortin receptor agonist: *afamelanotide (99), bremelanotide (95), modimelanotide*

neuromodulator / neuroprotective agent: *davunetide (100), ebiratide (56), obinepitide (96), vanutide cridificar (100)*

peptic ulcer: *sulglicotide (29), triletide (50)*

pulmonary surfactant: *lusupultide* (80), *sinapultide* (78)

sedative: *emideltide* (70)

thrombin receptor as an agonist, promoter of bone and skin wound healing:
rusalptide (96)

transforming growth factor beta-1 inhibitor: *disitertide* (99)

treatment of Parkinson's disease: *doreptide* (59), *pareptide* (38)

zonulin antagonist (in celiac disease): *larazotide* (99)

other: *defibrotide* (44) (nucleotide, the preferred stem has been used but not in accordance with its definition see item 4.7).

4.23. Peptide vaccines / recombinant vaccines

Definition of peptide vaccines: vaccine in which antigens are produced from synthetic peptides and transported through the bloodstream by an adjuvant, in order to stimulate an immune response.

Definition of recombinant vaccines: vaccine produced from a cloned gene.

Description of recombinant vaccines: there are certain antigens on viruses and bacteria which are better at stimulating an antibody response by the animal than others. The genes for these antigens can be isolated, and made to produce large quantities of the antigens they code for. A recombinant vaccine contains these antigens, not the whole organism. Compare with "modified live vaccine" and "killed vaccine".

peptide vaccines (see item 4.22): *-motide*

abecomotide (109), *alicdamotide* (109), *amilomotide* (105), *asudemotide* (107), *disomotide* (94), *elpamotide* (103), *latromotide* (107), *ovemotide* (94), *pradimotide* (107), *tanurmotide* (109), *tecemotide* (108), *tertomotide* (98), *tiplimotide* (82), *trepamotide* (107), *zastumotide* (110).

recombinant vaccine:

verpasep caltespen (95) (heat-shock protein HSP 65 (*Mycobacterium bovis* strain BCG) fusion protein with transcription factor E7 (human papilloma virus 16)).

The suffix *-tespen* is the indicator of heat shock protein.

4.24. Pituitary / placental glycoprotein hormones

The names selected by the International Union of Pure and Applied Chemistry–International Union of Biochemistry (IUPAC-IUB) have, to date, been chosen for compounds with an amino acid sequence identical to that of the naturally occurring human hormones. Addition of a Greek letter as the second part of the name will allow differentiation of different glycosylation patterns for compounds produced by biotechnology (see item 3.4 - general policies for glycosylated compounds).

follicle stimulating hormones: ending in (-)*follitropin*

corifollitropin alfa (80), *follitropin alfa (71)*, *follitropin beta (75)*, *follitropin gamma (106)*, *urofollitropin (57)*, *varfollitropin alfa (101)*

gonadotropin: ending in *-gonadotropin*

choriogonadotropin alfa (76), *chorionic gonadotrophin (01)*: chorionic gonadotropins, obtained from human serum and urine during pregnancy and has both lutropin and follitropin activity

serum gonadotrophin (01): used for the follicle stimulating hormone (FSH, follitropin) from serum of pregnant mares

luteinizing hormones: ending in (-)*lutropin*

lutropin alfa (71).

4.25. Pituitary hormone-release stimulating peptides

The common stem for pituitary hormone-release stimulating peptides is *-relin*.

LHRH-release-stimulating peptides:

avorelin (74), buserelin (36), deslorelin (61), fertirelin (42), gonadorelin (32), goserelin (55), histrelin (53), leuprorelin (47), lutrelin (51), nafarelin (50), peforelin (93), triptorelin (58), zoptarelin doxorubicin (107)

growth hormone release-stimulating peptides: *-morelin*

anamorelin (97), capromorelin (83), dmorelin (59), examorelin (72), ipamorelin (78), lenomorelin (106), macimorelin (100), pralmorelin (77), rismorelin (74), sermorelin (56), somatorelin (57), tabimorelin (86), tesamorelin (96), ulimorelin (103)

thyrotropin releasing hormone analogues: *-tirelin*

azetirelin (60), montirelin (58), orotirelin (58), posatirelin (60), protirelin (31), taltirelin (75)

thyrotropin alfa (78) (thyrotropin releasing hormone (TRH) analog, belongs to this group but in which the preferred stem has not been used)

other: *corticorelin (66)* (diagnostic agent).

4.26. Receptor molecules, native or modified

The stem for receptor molecules, native or modified is *-cept*.

A preceding infix should designate the target.

B-cell activating factor receptors: *-babiobcept (98)*

vascular endothelial growth factor receptors: *-beraflibercept (96), conbercept (105)*

complement receptors: *-comirococept (91)*

subgroup of interferon receptors: *-far-*
bifarcept (86)

frizzled family receptors: *-fri-*
ipafricept (109)

lymphocyte function-associated antigen 3 receptors: *-lefa-*
alefacept (84)

interleukin-1 receptors: *-na-*
rilonacept (95)

tumour necrosis factor (TNF) receptors: *-ner-*
baminercept (99), *etanercept (81)*, *lenercept (72)*, *onercept (86)*,
pegsunercept (95)

cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) receptors: *-ta-*
abatacept (91), *belatacept (93)*

transmembrane activator and calcium modulator and cyclophilin ligand
interactor: *-taci-*
atacicept (95)

transforming growth factor receptors: *-ter-*
dalantercept (105), *luspatercept (110)*, *ramatercept (108)*, *sotatercept (104)*

antiviral receptors: *-vir-*
alvircept sudotox (69).

4.27. Synthetic polypeptides with a corticotropin-like action

The common stem for synthetic polypeptides with a corticotropin-like action is *-actide*.

alsactide (45), *codactide (24)*, *giractide (29)*, *norleusactide (18)*, *seractide (31)*, *tetracosactide (18)*, *tosactide (24)*, *tricosactide (44)*, *tridecactide (97)*.

4.28. Thrombomodulins

sothrombomodulin alfa (101), *thrombomodulin alfa (94)*.

4.29. Toxins

toxin ML-1 (mistletoe lectin I) (*Viscum album*): *aviscumine* (86).

4.30. Vasoconstrictors, vasopressin derivatives

The common stem for vasoconstrictors, vasopressin derivatives is *-pressin*.

argipressin (13), *desmopressin* (33), *felypressin* (13), *lypressin* (13), *ornipressin* (22), *selepressin* (105), *terlipressin* (46), *vasopressin injection* (16).

4.31. Various⁷

- *abicipar pegol* (108): pegylated composite protein for clinical applications (CPCA), with alternative scaffold domain to antigen receptors based on ankyrin repeats, anti- [*Homo sapiens* VEGFA (vascular endothelial growth factor A, VEGF-A, VEGF)]; glycyL-seryl-ankyrin repeats (3-35, 36-68, 69-101, 102-123)-lysyl-dialanyl-bis(triglycyl-seryl) linker (127-134)-cysteinyl (1-135), conjugated via a maleimide group linker (thioether bond to C135) to a single linear methoxy polyethylene glycol 20 (mPEG20)
- *alisporivir* (100): [8-(*N*-methyl-D-alanine),9-(*N*-ethyl-L-valine)]cyclosporine
- *agatolimod* (98): DNA, d(*P*-thio)(T-C-G-T-C-G-T-T-T-T-G-T-C-G-T-T-T-T-G-T-C-G-T-T)
- *andexanet alfa* (110): des-(6-39)-human blood-coagulation factor X light chain (98-108')-disulfide with [185'-alanine (S>A)]human activated factor Xa heavy chain, produced in Chinese hamster ovary (CHO) cells (glycoform alfa)

⁷ The descriptions following the INN names may not be the complete definitions as shown in the publications of INN Lists.

- *angiotensin II (65)*: 5-L-isoleucineangiotensin II (the source of the material should be indicated)
- *angiotensinamide (12)*: N-{1-{N-{N-{N-[N-(N²-asparaginyllarginyl)valyl]tyrosyl}valyl}histidyl}prolyl}-3-phenylalanine
- *asvasiran (111)*: small interfering RNA (siRNA) inhibitor of human Respiratory Syncytial Virus replication;
duplex of guanylyl-(3'→5')-guanylyl-(3'→5')-cytidylyl-(3'→5')-uridylyl-(3'→5')-cytidylyl-(3'→5')-uridylyl-(3'→5')-uridylyl-(3'→5')-adenylyl-(3'→5')-guanylyl-(3'→5')-cytidylyl-(3'→5')-adenylyl-(3'→5')-adenylyl-(3'→5')-adenylyl-(3'→5')-guanylyl-(3'→5')-uridylyl-(3'→5')-cytidylyl-(3'→5')-adenylyl-(3'→5')-adenylyl-(3'→5')-guanylyl-(3'→5')-thymidylyl-(3'→5')-thymidine and thymidylyl-(5'→3')-thymidylyl-(5'→3')-cytidylyl-(5'→3')-cytidylyl-(5'→3')-guanylyl-(5'→3')-adenylyl-(5'→3')-guanylyl-(5'→3')-adenylyl-(5'→3')-adenylyl-(5'→3')-uridylyl-(5'→3')-cytidylyl-(5'→3')-guanylyl-(5'→3')-uridylyl-(5'→3')-uridylyl-(5'→3')-uridylyl-(5'→3')-cytidylyl-(5'→3')-adenylyl-(5'→3')-guanylyl-(5'→3')-uridylyl-(5'→3')-uridylyl-(5'→3')-uridylyl-(5'→3')-cytidine
- *bamosiran (106)*: siRNA inhibitor of β2-adrenergic receptor production
- *bevasiranib (108)*: siRNA inhibitor of Vascular Endothelial Growth Factor (VEGF) production
- *blisibimod (107)*: B-cell activating factor (BAFF)-binding peptide fragment/human IgG1 Fc fusion protein
- *calcitonin (80)*: a polypeptide hormone that lowers the calcium concentration in blood (the species specificity should be indicated in brackets behind the name)
- *conestat alfa (107)*: human plasma protease C1 inhibitor (C1 esterase inhibitor) (N,O-glycosylated recombinant protein expressed in the mammary gland of transgenic rabbits), glycoform α
- *dianexin (109)*: recombinant DNA derived annexin A5 dimer covalently linked by a 14 residues peptide linker, produced in *Escherichia coli* (nonglycosylated): L-methionyl-human annexin A5 fusion protein with glycyl-L-seryl-L-leucyl-L-α-glutamyl-L-valyl-L-leucyl-L-phenylalanyl-L-glutaminylglycyl-L-prolyl-L-serylglycyl-L-lysyl-L-leucyl-human annexin A5

- *delcasertib (105)*: human immunodeficiency virus 1 protein Tat-(46-57)-peptide (1→1')-disulfide with L-cysteinyl-[mouse protein kinase C delta type-(8-17)-peptide]
- *edifoligide (89)*: oligonucleotide
- *egaptivon pegol (111)*: a pegylated aptamer which binds von Willebrand factor; 5'-O-[[6-(carboxyamino)hexyl]hydroxyphosphoryl]-2'-O-methylguanylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-deoxyguanylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-deoxyadenylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-O-methyl-P-thioguanlylyl-(3'→5')-thymidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-deoxyguanylyl-(3'→5')-2'-deoxyguanylyl-(3'→5')-thymidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-deoxycytidylyl-(3'→5')-2'-O-methyladenylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-O-methylcytidylyl-(3'→3')-thymidine, carbamate ester with monomethyl ether of polyethylene glycol (20 kDa)
- *emapticap pegol (108)*: β-L-guanylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-adenylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-cytidylyl-(3'→5')-β-L-uridylyl-(3'→5')-β-L-guanylyl-(3'→5')-β-L-guanosine 6-{2-(N-[ω-methylpoly(oxyethan-1,2-diyl)]-2-[[ω-methylpoly(oxyethan-1,2-diyl)]oxy}acetamido)acetamido}hexyl hydrogen 5'-phosphate
- *epelestat (92)*: human recombinant neutrophil elastase inhibitor, bovine pancreatic trypsin inhibitor (BPTI) homologue

- *hemoglobin glutamer (80)*: the species specificity should be indicated in brackets behind the name, "(bovine)"; the average mass of the polymer is given as e.g. haemoglobin glutamer-250 for 250kD
- *hemoglobin crosfumaril (76)*: hemoglobin A₀ (human α₂β₂ tetrameric subunit), α-chain 99,99'-diamide with fumaric acid
- *hemoglobin crosfumaril (bovine) (108)*: S^{3.β92}, S^{3.β92}-bis(2-amino-2-oxoethyl)-N^{6.α99}, N^{6.α99}-(but-2-enedioyl)bovine hemoglobulin (α₂β₂ tetramer)
- *hemoglobin raffimer (89)*
- *imetelstat (101)*: oligonucleotide telomerase inhibitor;
3'-amino-3'-deoxy-*P*-thiothymidylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioadenylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioguanilyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioguanilyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioguanilyl-(3'→5')-3'-amino-3'-deoxy-*P*-thiothymidylyl-(3'→5')-3'-amino-3'-deoxy-*P*-thiothymidylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioadenylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioguanilyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioadenylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thiocytidylyl-(3'→5')-3'-amino-2',3'-dideoxy-*P*-thioadenylyl-(3'→5')-3'-amino-2',3'-dideoxyadenosine 5'-{ *O*-[2-hydroxy-3-(hexadecanoylamino)propyl] hydrogen phosphorothioate }
- *enadenotucirev (111)*: chimeric oncolytic adenovirus Ad3/Ad11p containing two deletions in the viral genome in the E3 region (2444 bp) and in the E4 region (24bp) and 197 non-homologous nucleotides in the E2B region
- *entolimod (108)*: L -methionyl- L -arginylglycyl- L -seryl-hexa(L-histidyl)glycyl- (*Enterobacteria phage T7* major capsid protein 10A-(1-11)-peptidyl)- L -arginyl- L -aspartyl- L -leucyl- L -tyrosyl-tetra(L-aspartyl)- L -lysyl- L -aspartyl- L -prolyl- (*Salmonella dublin* flagellin-(1-176)-peptidyl)- L -seryl- L -prolylglycyl- L -isoleucyl- L -seryl-pentaglycyl- L -isoleucyl- L -leucyl- L -aspartyl- L -seryl- L -methionylglycyl- (*Salmonella dublin* flagellin-(402-505)-peptide)
- *iodinated (¹²⁵I) human serum albumin (24)*: human serum albumin iodinated with radioactive iodine (¹²⁵I)
- *iodinated (¹³¹I) human serum albumin (24)*: human serum albumin iodinated with radioactive iodine (¹³¹I)

- *iropilact (74)*: N-L-methionyl blood platelet factor 4 (human subunit)
- *ismomultin alfa (91)*: 47-261-Glycoprotein gp 39 (human clone CDM8-gp39 reduced)
- *lexaptepid pegol (108)*: β -L-guanylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-cytidine 6-{2-(N-[ω -methylpoly(oxyethan-1,2-diyl)]-2-{[ω -methylpoly(oxyethan-1,2-diyl)]oxy}acetamido)acetamido}hexyl hydrogen 5'-phosphate
- *litenimod (96)*: (3'-5')d(P-thio)(T-A-A-A-C-G-T-T-A-T-A-A-C-G-T-T-A-T-G-A-C-G-T-C-A-T)
- *macrosalb (¹³¹I) (33)*: macroaggregated iodinated (¹³¹I) human albumin
- *macrosalb (^{99m}Tc)(33)*: technetium (^{99m}Tc) labelled macroaggregated human serum albumin
- *metenkefalin (97)*: L-tyrosylglycylglycyl-L-phenylalanyl-L-methionine β -endorphin human-(1-5)-peptide
- *metreleptin (82)*: N-methionylleptin (human)
- *mirostipen (85)*: [23-methionine] human myeloid progenitor inhibitory factor 1-(23-99)-peptide
- *nagrestipen (76)*: 26-L-alaninelymphokine MIP 1 α (human clone pAT464 macrophage inflammatory)

- *ocriplasmin (101)*: truncated human plasmin:
human plasmin heavy chain A-(543-561)-peptide (548-666;558-566)-bisdisulfide with human plasmin light chain B
- *olaptosed pegol (109)*: β -L-guanylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-uridylyl-(3'→5')- β -L-adenylyl-(3'→5')- β -L-cytidylyl-(3'→5')- β -L-guanylyl-(3'→5')- β -L-cytidine 6-{2-(N-[ω -methylpoly(oxyethan-1,2-diyl)]-2-[[ω -methylpoly(oxyethan-1,2-diyl)]oxy}acetamido)acetamido}hexyl hydrogen 5'-phosphate
- *opebacan (83)*: 132-L-alanine-1-193-bactericidal / permeability-increasing protein (human)
- *orgotein (31)*: a group of soluble metalloproteins isolated from liver, red blood cells, and other mammalian tissues
- *ovandrotone albumin (52)*: 3-[(3,17-dioxoandrost-4-en-7 α -yl)thio]propionic acid, serum albumin conjugate
- *parathyroid hormone (90)*: non glycosylated human parathyroid hormone, the origin should be indicated between brackets after the INN, for example (r. *E. coli*) for recombinant produced by *Escherichia coli*
- *patisiran (109)*: small interfering RNA (siRNA);
RNA duplex of guanylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-adenylyl-(3'→5')-adenylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-adenylyl-(3'→5')-adenylyl-(3'→5')-guanylyl-(3'→5')-adenylyl-(3'→5')-guanylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-adenylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-O-methylcytidylyl-(3'→5')-adenylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-thymidylyl-(3'→5')-thymidine with thymidylyl-(5'→3')-thymidylyl-(5'→3')

cytidyl-(5'→3')-adenyl-(5'→3')-2'-O-methyluridylyl-(5'→3')-uridylyl-(5'→3')-
 guanylyl-(5'→3')-guanylyl-(5'→3')-uridylyl-(5'→3')-uridylyl-(5'→3')-cytidyl-(5'
 →3')-uridylyl-(5'→3')-cytidyl-(5'→3')-adenyl-(5'→3')-2'-O-methyluridylyl-(5'
 →3')-adenyl-(5'→3')-adenyl-(5'→3')-guanylyl-(5'→3')-guanylyl-(5'→3')-
 uridylyl-(5'→3')-adenosine

- *pegaptanib (88)*: 5'-ester of (2'-deoxy-2'-fluoro)C-Gm-Gm-A-A-(2'-deoxy-2'-
 fluoro)U-(2'-deoxy-2'-fluoro)C-Am-Gm-(2'-deoxy-2'-fluoro)U-Gm-Am-Am-(2'-
 deoxy-2'-fluoro)U-Gm-(2'-deoxy-2'-fluoro)C-(2'-deoxy-2'-fluoro)U-(2'-deoxy-2'-
 fluoro)U-Am-(2'-deoxy-2'-fluoro)U-Am-(2'-deoxy-2'-fluoro)C-Am-(2'-deoxy-2'-
 fluoro)U-(2'-deoxy-2'-fluoro)C-(2'-deoxy-2'-fluoro)C-Gm-(3'→3')-dT with α,α' -
 [(1S)-1-[[5-(phosphonoxy)pentyl]carbamoyl]pentane-1,5-
 diyl]bis(iminocarbonyl)]bis[ω -methoxypoly(oxyethane-1,2-diyl)]
- *pegdinetanib (103)*: 94 residues protein derived from human fibronectin 10th type
 III domain, pegylated:
 glycyll[1438-L-arginine(D>R),1439-L-histidine(A>H),1441-L-
 histidine(A>H),1442-L-phenylalanine(V>F),1443-L-proline(T>P),1444-L-
 threonine(V>T),1467-L-leucine(G>L),1468-L-glutamine(S>Q),1469-L-
 proline(K>P),1470-L-proline(S>P),1492-L-aspartic acid(G>D),1493-
 glycine(R>G),1494-L-arginine(G>R),1495-L-asparagine(D>N),1496-
 glycine(S>G),1497-L-arginine(P>R),1498-L-leucine(A>L),1499-L-
 leucine(S>L),1501-L-isoleucine(K>I),1515-S-[(3RS)-1-(1-[[α -
 methylpoly(oxyethylene)]carbamoyl]-3-[[α -
 methylpoly(oxyethylene)]carbamoyl]oxy)methyl]-8,13-dioxo-
 1,4-dioxo-9,12-diazapentadecan-15-yl]-2,5-dioxopyrrolidin-3-yl]-
 L-cysteine(S>C)]human fibronectin-(1424-1516)-peptide
- *revusiran (111)*: [(2S,4R)-1-{30-(2-acetamido-2-deoxy- β -D-galactopyranosyl)-
 14,14-bis[16-(2-acetamido-2-deoxy- β -D-galactopyranosyl)-5,11-dioxo-2,16-
 dioxo-6,10-diazahexadecyl]-12,19,25-trioxo-16,30-dioxo-13,20,24-
 triazatriacontanoyl}-4-hydroxypyrrolidin-2-yl]methyl hydrogen 2'-
 deoxy-2'-fluorouridylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-deoxy-
 2'-fluoroguanlylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-deoxy-2'-
 fluoroadenylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-deoxy-2'-
 fluorouridylyl-(3'→5')-2'-O-methyluridylyl-(3'→5')-2'-deoxy-2'-
 fluorocytidylyl-(3'→5')-2'-deoxy-2'-fluoroadenylyl-(3'→5')-2'-deoxy-2'-
 fluorouridylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-deoxy-2'-
 fluorouridylyl-(3'→5')-2'-O-methyladenylyl-(3'→5')-2'-O-methyladenylyl-(
 3'→5')-2'-O-methylcytidylyl-(3'→5')-2'-deoxy-2'-
 fluorocytidylyl-(3'→5')-2'-O-methyladenylyl-(3'→5')-2'-deoxy-2'-
 fluoroadenylyl-(3'→5')-2'-O-methylguanylyl-(3'→5')-2'-deoxy-2'-
 fluoroadenylate duplex with 2'-O-methyl-P-thiocytidylyl-(5'→3')-2'-
 deoxy-2'-fluoro-P-thiouridylyl-(5'→3')-2'-O-methyladenylyl-(5'→3')-2'-
 deoxy-2'-fluorocytidylyl-(5'→3')-2'-O-methylcytidylyl-(5'→3')-2'-deoxy-

2'-fluorocytidylyl-(5'→3')-2'-*O*-methyluridylyl-(5'→3')-2'-deoxy-2'-
 fluoroadenylyl-(5'→3')-2'-*O*-methyladenylyl-(5'→3')-2'-deoxy-2'-
 fluoroadenylyl-(5'→3')-2'-*O*-methylguanylyl-(5'→3')-2'-*O*-methyluridylyl-(
 5'→3')-2'-*O*-methyladenylyl-(5'→3')-2'-deoxy-2'-
 fluorocytidylyl-(5'→3')-2'-*O*-methyladenylyl-(5'→3')-2'-deoxy-2'-
 fluorouridylyl-(5'→3')-2'-deoxy-2'-fluorouridylyl-(5'→3')-2'-deoxy-2'-
 fluoroguanlylyl-(5'→3')-2'-*O*-methylguanylyl-(5'→3')-2'-deoxy-2'-
 fluorouridylyl-(5'→3')-2'-*O*-methyluridylyl-(5'→3')-2'-deoxy-2'-
 fluorocytidylyl-(5'→3')-2'-*O*-methyluridine

- *rintatolimod (102)*: poly[5']-inosinylyl-(3'→) duplex with poly[dodecakis[3']-
 cytidylyl-(5'→)3')-uridylyl-(5'→)
- *secretin (01)*: hormone of the duodenal mucosa which activates the pancreatic
 secretion and lowers the blood-sugar level
- *secretin human (106)*: human peptide hormone secretin
- *serelaxin (105)*: human relaxin 2 (relaxin H2)
- *talactoferrin alfa (93)*: recombinant human lactoferrin
- *tadekinig alfa (90)*: interleukin-18 binding protein (human gene IL 18BP isoform
 a precursor)
- *thrombin alfa (97)*: human thrombin (recombinant, glycoform α)
- *tiprelestat (103)*: human elafin (elastase-specific inhibitor, skin-derived
 antileukoproteinase, peptidase inhibitor 3)
- *topsalysin (111)*: recombinant DNA derived proaerolysin, pore-forming protein,
 from *Aeromonas hydrophila*, with the furin site substituted with a prostate
 specific antigen (PSA) cleavage site, fusion protein with 6 histidines, produced in
Escherichia coli (nonglycosylated):
 [427-L-histidine(K>H),428-L-serine(V>S),429-L-serine(R>S),430-L-
 lysine(R>K),431-L-leucine(A>L),432-L-glutamine(R>Q)]proaerolysin *Aeromonas*
hydrophila fusion protein with hexa-L-histidine

- *torapsel (91)*: 42-89-glycoprotein (human clone PMT21:PL85 P-selectin glycoprotein ligand 1) fusion protein with immunoglobulin (human constant region)
- *trebananib (106)*: immunoglobulin G1 Fc fragment fused with two synthetic polypeptides that bind the *Homo sapiens* ANGPT2 (angiopoietin 2); methionyl (1) -gamma1 heavy chain fragment (2-228) [*Homo sapiens* IGHG1*01 hinge (EPKSC 1-5>del) (2-11), CH2 (12-121), CH3 (122-228)] fused, at the C-terminal end, with a synthetic polypeptide that comprises two 14-mer amino acid repeats that bind angiopoietin 2 (229-287) [linker (229-235) -14-mer (236-249) – linker (250-271) -14-mer (272-285) -leucyl-glutamate]; (7-7':10-10')-bisdisulfide dimer
- *tremacamra (78)*: 1-453-glycoprotein ICAM-I (human reduced)
- *tropasminogen alfa (99)*: thrombin-activable plasminogen; endo-[(558a(559)-558h(365))-human coagulation factor XI-(363-370)-peptide]-des-(559-562)-[606(610)-lysine,623(627)-lysine]human plasminogen, glycoform α
- *votucalis (96)*: methionyl[145-leucine]FS-HBP2 (*Rhipicephalus appendiculatus* (Brown ear tick) Female-Specific Histamine-Binding Protein 2).

5. CURRENT CHALLENGES

The challenges currently faced include the following:

- The INN Expert Group, when selecting names for recombinant proteins, has to deal not with substances with well-defined structures but with products of highly complex composition or even with mixtures of such products.
- It is not only modified proteins that might differ from their naturally occurring counterparts, products derived by expression of the natural gene in foreign host cells may also differ structurally, biologically or immunologically from the natural protein.
- Glycoproteins particularly may occur in forms that differ in the structure of one or more of their carbohydrate units, a phenomenon known as microheterogeneity and resulting in a heterogeneous population of molecules. Such differences may affect both the size and the charge of individual glycoproteins.
- A variety of novel biotechnology-derived products are under development, all of which will require specific policies on how to deal with such products.
- Clearly, the INN nomenclature of biological medicinal products is an area of increasing complexity.

REFERENCES

1. The use of stems in the selection of International Nonproprietary Names (INN) for pharmaceutical substances, 2013, *WHO/EMP/RHT/TSN/2013.1* *
2. International nomenclature and gene therapy products (*WHO Drug Information*, Vol.19, N°2, 2005, p.103) *
3. Pre-stems: Suffixes used in the selection of INN (document regularly updated on the INN website) *
4. Consultation on International Nonproprietary Names (INN) and biological products, *INN Working Document 00.118 (2002)*
5. INNs for biotechnological products: collaboration with other parties, *WHO/Pharm S/Nom 1763 (1999)*
6. INNs for recombinant vaccines and viruses, *WHO/Pharm S/Nom 1719 (1998)*
7. INNs for biosynthetic vaccines, *WHO/Pharm S/Nom 1419 (1994)*
8. INN nomenclature for peptides, glycopeptides, proteins and glycoproteins, *WHO/Pharm S/Nom 1428 (1994)*
9. INNs for immunoglobulins, *WHO/Pharm S/Nom 1517 (1995)*
10. INNs for immunoglobulins, *WHO/Pharm S/Nom 101 (1967)*
11. General policies for monoclonal antibodies, *INN Working Document 09.251, 24/06/2009* *
12. Definition of INNs for substances prepared by biotechnology, *WHO/Pharm S/Nom 1348 (1992)*
13. International Nonproprietary Names (INN) for pharmaceutical substances, Lists 1-111 of proposed INN and Lists 1-72 of Recommended INN, Cumulative List N°15, 2013
14. Guidelines on the use of International Nonproprietary Names (INNs) for pharmaceutical substances, *WHO/Pharm S/Nom 1570 (1997)*

15. INNs of the *-tropin (trophin)* series: pituitary hormones, *WHO/Pharm S/Nom 1406 (1993-1995)*
16. INNs for blood factors, *WHO/Pharm S/Nom 1362 (1994)*
17. INNs for growth factor, *WHO/Pharm S/Nom 1318 (1991)*
18. INNs for heparin derivatives, *WHO/Pharm S/Nom 1031 (1985-1991)*
19. Nomenclature of insulin injections, *WHO/Pharm S/Nom 1127 (1986)*
20. Generic names for genetically-engineered insulins, *WHO/Pharm S/Nom 737 (1980-1983)*
21. International nonproprietary names (INN) for pharmaceutical substances: names for radicals, groups & others (Comprehensive list), *WHO/EMP/RHT/TSN/2014.2**

** These documents are available on the INN Programme Website at:
<http://www.who.int/medicines/services/inn/en/>.*

The list of INN for composite proteins published⁸

classified by groups

alb- (human serum albumin)**alb- & -cog***albutrepenonacog alfa (109)*

human coagulation factor IX (EC 3.4.21.22, Christmas factor, plasma thromboplastin component) 148-threonine variant fusion protein with prolyl(human coagulation factor IX 148-threonine variant-(137-153)-peptide) fusion protein with human serum albumin, produced in CHO cells (alfa glycoform)

alb- & -interferon*albinterferon alfa-2b (99)*

human serum albumin (585 residues) fusion protein with human interferon α -2b (165 residues)

alb- & -tide*albenatide (111)*

$S^{3,34}$ -{1-[(23S)-23-{[exendin-4 *Heloderma suspectum* precursor-(48-86)-peptidyl (exenatidyl)]amino}-3,12,24-trioxo-7,10-dioxa-4,13,18,25-tetraazapentacosyl]-2,5-dioxopyrrolidin-3-yl}human serum albumin.

Peptide is synthetic, and human serum albumin is produced in *Saccharomyces cerevisiae*.

⁸ It should be noted that this list may not be comprehensive (eg. pegylated substances are not included) and the descriptions under the names are the ones published.

albiglutide (97)

([8-glycine]human glucagon-like peptide 1-(7-36)-peptidyl)([8-glycine]human glucagon-like peptide 1-(7-36)-peptidyl)(human serum albumin (585 residues))

Others:

-al- & -grastim

balugrastim (107)

human serum albumin (585 residues) fusion protein with des-(1-alanine,37-valine,38-serine,39-glutamic acid)-human granulocyte colony-stimulating factor (pluripoietin)

-ase

revelucosidase alfa (111)

des-(2-7)-human insulin-like growth factor II fusion protein with glycy-L-alanyl-L-prolyl-human lysosomal alpha-glucosidase (acid maltase, aglucosidase alfa) produced in Chinese hamster ovary (CHO) cells, glycoform alfa

senrebotase (107)

L-methionylglycyl-L-seryl-des-(445-glycine,446-L-tyrosine)-[2-L-glutamic acid,432,442,444,447-tetra-L-aspartic acid]botulinum neurotoxin A precursor 27-L-alanine variant light chain (433-41')-disulfide with [14-L-arginine,15-L-lysine]human nociceptin fusion protein with L-alanyl-L-leucyl-L-alanyltris(tetraglycyl-L-seryl)-[3-L-valine,4-L-leucine,5-L-glutamine-418-L-leucine,419-L-aspartic acid]botulinum neurotoxin A heavy chain-(1-419)-peptide

-cept

abatacept (91)

1-25-oncostatin M (human precursor) fusion protein with CTLA-4 (antigen) (human) fusion protein with immunoglobulin G1 (human heavy chain fragment), bimolecular (146→146')-disulfide

aflibercept (96)

des-432-lysine-[human vascular endothelial growth factor receptor 1-(103-204)-peptide (containing Ig like C2 type 2 domain) fusion protein with human vascular endothelial growth factor receptor 2-(206-308)-peptide (containing Ig like C2 type 3 domain fragment) fusion protein with human immunoglobulin G1-(227 C-terminal residues)-peptide (Fc fragment)], (211-211':214-214')-bisdisulfide dimer

alefacept (84)

1-92-antigen LFA-3 (human) fusion protein with human immunoglobulin G1 (hinge-C_H2-C_H3 γ 1-chain), dimer

atacicept (95)

[86-serine,101-glutamic acid,196-serine,197-serine,222-aspartic acid,224-leucine][human tumor necrosis factor receptor superfamily member 13B-(30-110)-peptide (TACI fragment containing TNFR-Cys 1 and TNFR-Cys 2) fusion protein with human immunoglobulin G1-(232 C-terminal residues)-peptide (γ 1-chain Fc fragment), (92-92':95-95')-bisdisulfide dimer

baminercept (99)

human tumor necrosis factor receptor superfamily member 3 (lymphotoxin- β receptor, TNF C receptor)-(2-195)-peptide (fragment of extracellular domain) fusion protein with human immunoglobulin heavy constant γ 1 chain Fc fragment [227 residues, hinge (195-205) des-(1-4),C5>V, CH2 (206-315), CH3 (316-421) des-K¹⁰⁷]

belatacept (93)

[Tyr²⁹,Glu¹⁰⁴,Gln¹²⁵,Ser¹³⁰,Ser¹³⁶,Ser¹³⁹,Ser¹⁴⁸](antigen CTLA-4 human-3-126)-peptide (fragment containing the human extracellular domain) fusion protein with immunoglobulin G1-[233 amino acids from the C-terminal of the heavy chain]-peptide (fragment containing the human monoclonal Fc domain), bimolecular (120 \rightarrow 120')-disulfide

briobcept (98)

aspartyl[1-valine,20-asparagine,27-proline](human tumor necrosis factor receptor superfamily member 13C (BAFF receptor, BlyS receptor 3 or CD268 antigen)-(1-71)-peptidyl (part of the extracellular domain))valyl(human immunoglobulin G1 Fc fragment, *Homo sapiens* IGHG1-(104-329)-peptide) (79-79':82-82')-bisdisulfide dimer

conbercept (105)

fusion protein for immune applications (FPIA) comprising *Homo sapiens* FLT1 (fms-related tyrosine kinase 1, vascular endothelial growth factor receptor 1, VEGFR1, vascular permeability factor receptor, tyrosine-protein kinase FRT) fragment, fused with *Homo sapiens* KDR (kinase insert domain receptor, vascular endothelial growth factor receptor 2, VEGFR2, protein-tyrosine kinase receptor FLK1, CD309) fragment, fused with *Homo sapiens* immunoglobulin G1 Fc fragment;

FLT1, 132-232 precursor fragment (1-101)-KDR, 227-421 precursor fragment (102-296) -glycyl-prolyl-glycyl (297-299) -gamma1 chain H-CH2-CH3 fragment (300-526) [*Homo sapiens* IGHG1*03 hinge 6-15 P13>L (307) (300-309), CH2 (310-419), CH3-CH-S (420-526)]; (305-305':308-308')-bisdisulfide dimer

dalantercept (105)

fusion protein for immune applications (FPIA) comprising *Homo sapiens* ACVRL1 (activin A receptor type II-like 1, activin receptor-like kinase 1, ALK1, ALK-1, serine/threonine-protein kinase receptor R3, SKR3, transforming growth factor-beta superfamily receptor type I, TGF-B superfamily receptor type I, TSR-I, HHT2, ORW2) fragment, fused with *Homo sapiens* immunoglobulin G1 Fc fragment;

ACVR2L1, 22-120 precursor fragment (1-99) -threonyl-triglycyl (100-103) - gamma1 chain H-CH2-CH3 fragment (104-328) [*Homo sapiens* IGHG1*03 hinge 8-15 (104-111), CH2 L1.3>A (115), G1>A (118), A115>V (211) (112-221), CH3 S85.3>P (284) (222-328)]; (107-107':110-110')-bisdisulfide dimer

etanercept (81)

1-235-tumor necrosis factor receptor (human) fusion protein with 236-467-immunoglobulin G1 (human γ 1-chain Fc fragment), dimer

ipafricept (109)

fusion protein for immune applications (FPIA) comprising *Homo sapiens* FZD8 (frizzled family receptor 8, Frizzled-8) extracellular domain, fused with *Homo sapiens* immunoglobulin G1 Fc fragment;
Homo sapiens FZD8 precursor fragment 28-158 (1-131) -*Homo sapiens* IGHG1*01 H-CH2-CH3 fragment (hinge 1-15 C5>S (136) (132-146), CH2 (147-256), CH3 (257-361), CHS (362-363)) (132-363); dimer (142-142':145-145')-bisdisulfide

lenercept (72)

1-182-tumor necrosis factor receptor (human reduced), (182→104')-protein with 104-330-immunoglobulin G1 (human clone pTJ5 C γ 1 reduced)

luspatercept (110)

fusion protein for immune applications (FPIA) comprising the *Homo sapiens* ACVR2B (activin receptor type 2B, activin A receptor type IIB, activin receptor type IIB, ACTR-IIB, ActR-IIB) extracellular domain, fused with *Homo sapiens* immunoglobulin G1 Fc fragment;
Homo sapiens ACVR2B precursor fragment 25-131 L79>D (55) (1-107) -linker triglycyl (108-110) - γ 1 chain H-CH2-CH3 fragment [*Homo sapiens* IGHG1*03 (hinge 8-15 (111-118), CH2 (119-228), CH3 (229-333), CHS (334-335))] (111-335); dimer (114-114':117-117')-bisdisulfide

ramatercept (108)

fusion protein for immune applications (FPIA) comprising *Homo sapiens* ACVR2B (activin A receptor type IIB, ActR-IIB) fragment, fused with *Homo sapiens* immunoglobulin G1 Fc fragment;
Homo sapiens ACVR2B precursor fragment 20-134 (1-115) –triglycyl (116-118) -*Homo sapiens* IGHG1*03 H-CH2-CH3 fragment (hinge 8-15 (119-126), CH2 A115>V (226) (127-236), CH3 (237-341), CHS (342-343)) (119-343); dimer (122-122':125-125')-bisdisulfide

rilonacept (95)

[653-glycine][human interleukin-1 receptor accessory protein-(1-339)-peptide (extracellular domain fragment) fusion protein with human type 1 interleukin-1 receptor-(5-316)-peptide (extracellular domain fragment) fusion protein with

human immunoglobulin G1-(229 C-terminal residues)-peptide (Fc fragment)],
(659-659':662-662')-bisdisulfide dimer

sotatercept (104)

fusion protein for immune applications (FPIA) comprising *Homo sapiens* ACVR2A (activin receptor type 2A, activin receptor type IIA) fragment fused with *Homo sapiens* immunoglobulin G1 Fc fragment;
Homo sapiens ACVR2A, 21-135 precursor fragment (1-115) -threonyl-triglycyl linker (116-119) -gamma1 chain H-CH2-CH3 fragment (120-344) [*Homo sapiens* IGHG1*03 hinge (120-127), CH2, A115>V (227) (128-237), CH3 (238-344)]; (123-123':126-126')-bisdisulfide dimer

-cept & -tox⁹ (-tox is for toxins (active or inactivated proteins))

alvircept sudotox (69)

N²-L-methionyl-1-178-antigen CD4 (human clone pT4B protein moiety reduced)(178→248')-protein with 248-L-histidine-249- L-methionine-250- L-alanine-251- L-glutamic acid-248-613-exotoxin A(*Pseudomonas aeruginosa* reduced)

-kin & -tox

cintridekin besudotox (92)

toxin hIL13-PE38QQR (plasmid phuIL13-Tx)

denileukin diftotox (78)

N-L-methionyl-387-L-histidine-388-L-alanine-1-388-toxin (*Corynebacterium diphtheriae* strain C7) (388→2')-protein with 2-133-interleukin 2 (human clone pTIL2-21a)

⁹ The names and the descriptions of toxins are published in Annex 4-1 of "International Nonproprietary Names (INN) for pharmaceutical substances. Names for radicals, groups & others: comprehensive list (WHO/EMP/RHT/TSN/2014.2)".

-mab & -dotin¹⁰

brentuximab vedotin (103)

immunoglobulin G1-kappa auristatin E conjugate, anti-[*Homo sapiens* TNFRSF8 (tumor necrosis factor receptor superfamily member 8, KI-1, CD30)], chimeric monoclonal antibody conjugated to auristatin E;

gamma1 heavy chain (1-446) [Mus musculus VH (IGHV1-84*02 -(IGHD)-IGHJ3*01) [8.8.10] (1-117) -*Homo sapiens* IGHG1*01 CH3 K130>del (118-446)], (220-218')-disulfide (if not conjugated) with kappa light chain (1'-218') [Mus musculus V-KAPPA (IGKV3-4*01 -IGKJ1*01) [10.3.9] (1'-111') -*Homo sapiens* IGKC*01 (112'-218''); (226-226'')-disulfide dimer; conjugated, on an average of 3 to 5 cysteinyl, to monomethylauristatin E (MMAE), via a maleimidecaproyl-valyl-citrullinyl-*p*-aminobenzylcarbamate (mc-val-cit-PABC) linker

vedotin

1-(6-{[(2S)-1-({(2S)-5-carbamoylamino-1-[(4-{[(2S)-{[(2S)-1-{(3R,4S,5S)-1-{(2S)-2-[(1R,2R)-3-{[(1S,2R)-1-hydroxy-1-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl}-3-methoxy-5-methyl-1-oxoheptan-4-yl](methyl)amino}-3-methyl-1-oxobutan-2-yl]amino}-3-methyl-1-oxobutan-2-yl]methylcarbamoyloxy}phenyl)amino]-1-oxopentan-2-yl}amino)-3-methyl-1-oxobutan-2-yl]amino}-6-oxohexyl)-2,5-dioxopyrrolidin-3-yl

enfortumab vedotin (109)

immunoglobulin G1-kappa, anti-[*Homo sapiens* PVRL4 (poliovirus receptor-related 4, nectin-4, nectin 4, PPR4, LNIR), *Homo sapiens* monoclonal antibody conjugated to auristatin E;

gamma1 heavy chain (1-447) [*Homo sapiens* VH (IGHV3-48*02 (98.00%) - (IGHD)-IGHJ6*01) [8.8.10] (1-117) -IGHG1*03 (CH1 (118-215), hinge (216-230), CH2 (231-340), CH3 (341-445), CHS (446-447)) (118-447)], (220-214')-disulfide with kappa light chain (1'-214') [*Homo sapiens* V-KAPPA (IGKV1-12*01 (96.80%) -IGKJ4*01) [6.3.9] (1'-107') -IGKC*01 (108'-214')]; dimer (226-226'':229-229'')-bisdisulfide; conjugated, on an average of 3 to 4 cysteinyl, to monomethylauristatin E (MMAE), via a cleavable maleimidecaproyl-valyl-citrullinyl-*p*-aminobenzylcarbamate (mc-val-cit-PABC) linker

vedotin (for *vedotin*, please refer to *brentuximab vedotin (103)*)

¹⁰ The names ending in *-dotin* and the descriptions are published in Annex 4-2 of "International Nonproprietary Names (INN) for pharmaceutical substances. Names for radicals, groups & others: comprehensive list (WHO/EMP/RHT/TSN/2014.2)".

lifastuzumab vedotin (110)

immunoglobulin G1-kappa auristatin E conjugate, anti-[*Homo sapiens* SLC34A2 (solute carrier family 34 sodium phosphate member 2, sodium/phosphate cotransporter 2B, NaPi2b, NaPi3b)], humanized monoclonal antibody conjugated to auristatin E;

gamma1 heavy chain (1-450) [humanized VH (*Homo sapiens* IGHV3-23*04 (85.70%) -(IGHD)-IGHJ5*01) [8.8.13] (1-120) -*Homo sapiens* IGHG1*03 (CH1 R120>K (217) (121-218), hinge (219-233), CH2 (234-343), CH3 (344-448), CHS (449-450)) (121-450)], (223-219')-disulfide with kappa light chain (1'-219') [humanized V-KAPPA (*Homo sapiens* IGKV1-39*01 (78.00%) -IGKJ1*01) [11.3.9] (1'-112')-*Homo sapiens* IGKC*01 (113'-219')]; dimer (229-229":232-232")-bisdisulfide; conjugated, on an average of 3 to 4 cysteinyl, to monomethylauristatin E (MMAE), via a cleavable maleimidocaproylvalyl-citrullinyl-*p*-aminobenzyloxycarbonyl (mc-val-cit-PABC) type linker

vedotin (for *vedotin*, please refer to *brentuximab vedotin (103)*)

pinatuzumab vedotin (108)

immunoglobulin G1-kappa auristatin E conjugate, anti-[*Homo sapiens* CD22 (sialic acid binding Ig-like lectin 2, SIGLEC2, SIGLEC-2, Blymphocyte cell adhesion molecule, BL-CAM, Leu-14)], humanized monoclonal antibody conjugated to auristatin E;

gamma1 heavy chain (1-450) [humanized VH (*Homo sapiens* IGHV3-66*01 (79.60%) -(IGHD)-IGHJ4*01) [8.8.13] (1-120) -*Homo sapiens* IGHG1*03 (CH1 R120>K (217) (121-218), hinge (219-233), CH2 (234-343), CH3 (344-448), CHS (449-450)) (121-450)], (223-219')-disulfide (if not conjugated) with kappa light chain (1'-219') [humanized V-KAPPA (*Homo sapiens* IGKV1-39*01 (80.00%) -IGKJ1*01) [11.3.9] (1'-112') -*Homo sapiens* IGKC*01 (113'-219')]; dimer (229-229":232-232")-bisdisulfide; conjugated, on an average of 3 to 4 cysteinyl, to monomethylauristatin E (MMAE), via a cleavable maleimidecaproyl-valylcitrullinyl-*p*-aminobenzylcarbamate (mc-val-cit-PABC) linker

vedotin (for *vedotin*, please refer to *brentuximab vedotin (103)*)

polatuzumab vedotin (110)

immunoglobulin G1-kappa auristatin E conjugate, anti-[*Homo sapiens* CD79B (immunoglobulin-associated CD79 beta)], humanized monoclonal antibody conjugated to auristatin E;

gamma1 heavy chain (1-447) [humanized VH (*Homo sapiens* IGHV3-23*04 (76.50%)-(IGHD)-IGHJ4*01) [8.8.10] (1-117) -*Homo sapiens* IGHG1*03 (CH1

R120>K (214)(118-215), hinge (216-230), CH2 (231-340), CH3 (341-445), CHS (446-447)) (118-447)], (220-218')-disulfide with kappa light chain (1'-218') [humanized V-KAPPA (*Homo sapiens* IGKV1-39*01 (85.90%) -IGKJ1*01 [10.3.9] (1'-111') -*Homo sapiens* IGKC*01 (112'-218'))]; dimer (226-226":229-229")-bisdisulfide; conjugated, on an average of 3 to 4 cysteinyl, to monomethylauristatin E (MMAE), via a cleavable maleimidocaproyl-valyl-citrullinyl-*p*-aminobenzyloxycarbonyl (mc-val-cit-PABC) type linker

vedotin (for *vedotin*, please refer to *brentuximab vedotin* (103))

***sofituzumab vedotin* (110)**

immunoglobulin G1-kappa auristatin E conjugate, anti-[*Homo sapiens* MUC16 (mucin 16, MUC-16, cancer antigen 125, CA125)], humanized monoclonal antibody conjugated to auristatin E;
gamma1 heavy chain (1-446) [humanized VH (*Homo sapiens* IGHV3-48*03 (79.80%) - (IGHD)-IGHJ4*01) [9.8.9] (1-116) -*Homo sapiens* IGHG1*03 (CH1 R120>K (213) (117-214), hinge (215-229), CH2 (230-339), CH3 (340-444), CHS (445-446)) (117-446)], (219-214')-disulfide with kappa light chain (1'-214') [humanized V-KAPPA (*Homo sapiens* IGKV1-5*01 (87.90%) -IGKJ1*01) [6.3.9] (1'-107') -*Homo sapiens* IGKC*01 (108'-214'))]; dimer (225-225":228-228")-bisdisulfide; conjugated, on an average of 3 to 4 cysteinyl, to monomethylauristatin E (MMAE), via a cleavable maleimidocaproylvalyl-citrullinyl-*p*-aminobenzyloxycarbonyl (mc-val-cit-PABC) type linker

vedotin (for *vedotin*, please refer to *brentuximab vedotin* (103))

***denintuzumab mafodotin* (111)**

immunoglobulin G1-kappa auristatin F conjugate, anti-[*Homo sapiens* CD19 (B lymphocyte surface antigen B4, Leu-12)], humanized monoclonal antibody;
gamma1 heavy chain (1-450) [humanized VH (*Homo sapiens* IGHV4-31*02 (84.80%) - (IGHD)-IGHJ4*01) [10.7.12] (1-120) -*Homo sapiens* IGHG1*01 (CH1 (121-218), hinge (219-233), CH2 (234-343), CH3 (344-448), CHS (449-450)) (121-450)], (223-213')-disulfide with kappa light chain (1'-213') [humanized V-KAPPA (*Homo sapiens* IGKV3-11*01 (85.30%) -IGKJ2*02) [5.3.9] (1'-106') -*Homo sapiens* IGKC*01 (107'-213'))]; dimer (229-229":232-232")-bisdisulfide; conjugated, on an average of 4 cysteinyl, to monomethylauristatin F (MMAF), via a noncleavable maleimidocaproyl (mc) linker

mafodotin

N-{(2*R*,3*R*)-3-[(2*S*)-1-[(3*R*,4*S*,5*S*)-4-({*N*-[6-(2,5-dioxo-2,5-dihydro-1*H*-pyrrol-1-yl)hexanoyl]-*N*-methyl-L-valyl-L-valyl}methylamino)-

3-methoxy-5-methylheptanoyl]pyrrolidin-2-yl]-3-methoxy-2-methylpropanoyl}-L-phenylalanine

vorsetuzumab mafodotin (107)

immunoglobulin G1-kappa auristatin F conjugate, anti- [*Homo sapiens* CD70 (tumor necrosis factor superfamily member 7, TNFSF7, CD27LG, CD27L)], humanized monoclonal antibody conjugated to auristatin F; gamma1 heavy chain (1-448) [humanized VH (*Homo sapiens* IGHV1-2*02 (86.70%) -(IGHD)-IGHJ6*01) [8.8.11] (1-118) -*Homo sapiens* IGHG1*01 (119-448)], (221-218')-disulfide (if not conjugated) with kappa light chain (1'-218') [humanized V-KAPPA (*Homo sapiens* IGKV4-1*01 (79.20%) -IGKJ1*01) [10.3.9] (1'-111') -*Homo sapiens* IGKC*01 (112'-218')]; (227-227":230-230")-bisdisulfide dimer; conjugated, on an average of 3 to 5 cysteinyl, to monomethylauristatin F (MMAF), via a non-cleavable maleimidocaproyl (mc) linker

mafodotin (for *mafodotin*, please refer to *denintuzumab mafodotin (111)*)

-mab & -kin

tucotuzumab celmoleukin (95)

immunoglobulin G1, anti-(tumor associated calcium signal transducer 1 (KS 1/4 antigen)) (human-mouse monoclonal huKS-IL2 heavy chain) fusion protein with interleukin 2 (human), disulfide with human-mouse monoclonal huKS-IL2 light chain, dimer

-mab & -tansine¹¹

anetumab ravtansine (109)

immunoglobulin G1-lambda2, anti- [*Homo sapiens* MSLN (mesothelin, pre-pro-megakaryocyte-potentiating factor, megakaryocyte potentiating factor, MPF, CAK1)], *Homo sapiens* monoclonal antibody conjugated to maytansinoid DM4; gamma1 heavy chain (1-450) [*Homo sapiens* VH (IGHV5-51*01 (94.90%) -(IGHD)-IGHJ4*01) [8.8.13] (1-120) -IGHG1*01 (CH1 (121-218), hinge (219-233), CH2 (234-343), CH3 (344-448), CHS (449-450)) (121-450)], (223-216')-

¹¹ The names ending in *-tansine* and the descriptions are published in Annex 4-2 of "International Nonproprietary Names (INN) for pharmaceutical substances. Names for radicals, groups & others: comprehensive list (WHO/EMP/RHT/TSN/2014.2)".

disulfide with lambda light chain (1'-217') [*Homo sapiens* V-LAMBDA (IGLV2-14*01 (95.60%) -IGLJ2*01) [9.3.11] (1'-111') -IGLC2*01 A43>G (155) (112'-217')]; dimer (229-229":232-232")-bisdisulfide; conjugated, on an average of 3 lysyl, to maytansinoid DM4 [*N*^{2'}-deacetyl-*N*^{2'}-(4-mercapto-4-methyl-1-oxopentyl)-maytansine] via the reducible SPDB linker [*N*-succinimidyl 4-(2-pyridyldithio)butanoate]

ravtansine

4-[(5-{[(1*S*)-1-{[(1*S*,2*R*,3*S*,5*S*,6*S*,16*E*,18*E*,20*R*,21*S*)-11-chloro-21-hydroxy-12,20-dimethoxy-2,5,9,16-tetramethyl-8,23-dioxo-4,24-dioxo-9,22-diazatetracyclo[19.3.1.1^{10,14}.0^{3,5}]hexacos-10,12,14(26),16,18-pentaen-6-yl]oxy}-1-oxopropan-2-yl](methylamino)-2-methyl-5-oxopentan-2-yl)disulfanyl]butanoyl

cantuzumab ravtansine (105)(66)

immunoglobulin G1-kappa, anti- [*Homo sapiens* MUC1 sialylated carbohydrate, tumour-associated (CA242, cancer antigen 242)], humanized monoclonal antibody conjugated to maytansinoid DM4;
gamma1 heavy chain (1-449) [humanized VH (*Homo sapiens* IGHV7-4-1*02 (76.50%) - (IGHD)-IGHJ2*01 R120>Q (111), L123>T (114)) [8.8.12] (1-119) - *Homo sapiens* IGHG1*01 (120-449)], (222-219')-disulfide with kappa light chain (1'-219') [humanized V-KAPPA (*Homo sapiens* IGKV2-28*01 (82.00%) - IGKJ3*01 V124>L (109), D125>E (110), I126>L (111)) [11.3.9] (1'-112') - *Homo sapiens* IGKC*01 (113'-219')]; (228-228":231-231")-bisdisulfide dimer; conjugated, on an average of 3 to 4 lysyl, to maytansinoid DM4 [*N*^{2'}-deacetyl-*N*^{2'}-(4-mercapto-4-methyl-1-oxopentyl)-maytansine] via the reducible SPDB linker [*N*-succinimidyl 4-(2-pyridyldithio)butanoate]

ravtansine (for *ravtansine*, please refer to *anetumab ravtansine (109)*)

coltuximab ravtansine (109)

immunoglobulin G1-kappa, anti- [*Homo sapiens* CD19 (B lymphocyte surface antigen B4, Leu-12)], chimeric monoclonal antibody conjugated to maytansinoid DM4;
gamma1 heavy chain (1-450) [Mus musculus VH (IGHV1-69*02 - (IGHD)-IGHJ4*01) [8.8.13] (1-120) - *Homo sapiens* IGHG1*01 (CH1 (121-218), hinge (219-233), CH2 (234-343), CH3 (344-448), CHS (449-450)) (121-450)], (223-211')-disulfide with kappa light chain (1'-211') [Mus musculus V-KAPPA (IGKV4-70*01 -IGKJ1*01) [5.3.7] (1'-104') - *Homo sapiens* IGKC*01 (105'-211')]; dimer (229-229":232-232")-bisdisulfide; conjugated, on an average of 3 to 4 lysyl, to maytansinoid DM4 [*N*^{2'}-deacetyl-*N*^{2'}-(4-mercapto-4-methyl-1-

oxopentyl)-maytansine] via the reducible SPDB linker [*N*-succinimidyl 4-(2-pyridyldithio)butanoate]

ravtansine (for *ravtansine*, please refer to *anetumab ravtansine* (109))

***indatuximab ravtansine* (105)(67)**

immunoglobulin G4-kappa, anti-*[Homo sapiens* SDC1 (syndecan-1, CD138)], chimeric monoclonal antibody conjugated to maytansinoid DM4; gamma4 heavy chain (1-449) [*Mus musculus* VH (IGHV1-9*01 - (IGHD)-IGHJ4*01) [8.8.15] (1-122) -*Homo sapiens* IGHG4*01 (123-449)], (136-214')-disulfide with kappa light chain (1'-214') [*Mus musculus* V-KAPPA (IGKV10-94*01 -IGKJ1*01) [6.3.9] (1'-107') -*Homo sapiens* IGKC*01 (108'-214')]; (228-228":231-231")-bisdisulfide dimer; conjugated, on an average of 3 to 4 lysyl, to maytansinoid DM4 [*N*²-deacetyl-*N*²-(4-mercapto-4-methyl-1-oxopentyl)-maytansine] via the reducible SPDB linker [*N*-succinimidyl 4-(2-pyridyldithio)butanoate]

ravtansine (for *ravtansine*, please refer to *anetumab ravtansine* (109))

***cantuzumab mertansine* (105)(66)**

immunoglobulin G1-kappa, anti-*[Homo sapiens* MUC1 sialylated carbohydrate, tumour-associated (CA242, cancer antigen 242)], humanized monoclonal antibody conjugated to maytansinoid DM1; gamma1 heavy chain (1-449) [humanized VH (*Homo sapiens* IGHV7-4-1*02 (76.50%) - (IGHD)-IGHJ2*01 R120>Q (111), L123>T (114)) [8.8.12] (1-119) - *Homo sapiens* IGHG1*01 (120-449)], (222-219')-disulfide with kappa light chain (1'-219') [humanized V-KAPPA (*Homo sapiens* IGKV2-28*01 (82.00%) - IGKJ3*01 V124>L (109), D125>E (110), I126>L (111)) [11.3.9] (1'-112') -*Homo sapiens* IGKC*01 (113'-219')]; (228-228":231-231")-bisdisulfide dimer; conjugated, on an average of 4 lysyl, to maytansinoid DM1 [*N*²-deacetyl-*N*²-(3-mercapto-1-oxopropyl)-maytansine] via the reductible SPP linker [*N*-succinimidyl 4-(2-pyridyldithio)pentanoate]

mertansine

x(4*RS*)-4[(3-{{(1*S*)-2-{{(1*S*,2*R*,3*S*,5*S*,6*S*,16*E*,18*E*,20*R*,21*S*)-11-chloro-21-hydroxy-12,20-dimethoxy-2,5,9,16-tetramethyl-8,23-dioxo-4,24-dioxa-9,22-diazatetracyclo[19.3.1.1^{10,14}.0^{3,5}]hexacos-10,12,14(26),16,18-pentaen-6-yl]oxy}-1-methyl-2-oxoethyl]methylamino}-3-oxopropyl]disulfanyl]pentanoyl}

lorvotuzumab mertansine (103)(65)

immunoglobulin G1-kappa, anti-[*Homo sapiens* NCAM1 (neural cell adhesion molecule 1, CD56, NCAM-1)], humanized monoclonal antibody conjugated to maytansinoid DM1;

gamma1 heavy chain (1-448) [humanized VH (*Homo sapiens* IGHV3-30*03 (91.80%) -(IGHD)-IGHJ4*01) [8.8.11] (1-118) -*Homo sapiens* IGHG1*01 (119-448)], (221-219')-disulfide with kappa light chain (1'-219') [humanized V-KAPPA (*Homo sapiens* IGKV2-30*02 (92.00%) -IGKJ1*01) [11.3.9] (1'-112') -*Homo sapiens* IGKC*01 (113'-219')]; (227-227":230-230")-bisdisulfide dimer; conjugated, on an average of 3 to 4 lysyl, to maytansinoid DM1 via a thiopentanoate linker

mertansine (for *mertansine*, please refer to *cantuzumab mertansine (105)(66)*)

trastuzumab emtansine (103)(65)

immunoglobulin G1-kappa, anti-[*Homo sapiens* ERBB2 (epidermal growth factor receptor 2, HER-2, p185c-erbB2, NEU, EGFR2)], humanized monoclonal antibody conjugated to maytansinoid DM1;

gamma1 heavy chain (1-449) [humanized VH (*Homo sapiens* IGHV3-66*01 (81.60%) -(IGHD)-IGHJ6*01 T123>L) [8.8.13] (1-120) -*Homo sapiens* IGHG1*03 (121-449) CH1 R120>K], (223-214')-disulfide with kappa light chain (1'-214') [humanized V-KAPPA (*Homo sapiens* IGKV1-39*01 (86.30%) -IGKJ1*01) [6.3.9] (1'-107') -*Homo sapiens* IGKC*01 (108'-214')]; (229-229":232-232")-bisdisulfide dimer; conjugated, on an average of 3 to 4 lysyl, to maytansinoid DM1 via a succinimidyl-4-(*N*-maleimidomethyl) cyclohexane-1-carboxylate (SMCC) linker

emtansine

4-({3-[(3-[(1*S*)-2-[(1*S*,2*R*,3*S*,5*S*,6*S*,16*E*,18*E*,20*R*,21*S*)-11-chloro-21-hydroxy-12,20-dimethoxy-2,5,9,16-tetramethyl-8,23-dioxo-4,24-dioxa-9,22-diazatetracyclo[19.3.1.1^{10,14}.0^{3,5}]hexacos-10,12,14(26),16,18-pentaen-6-yl]oxy}-1-methyl-2-oxoethyl]methylamino)-3-oxopropyl)sulfanyl]-2,5-dioxopyrrolidin-1-yl}methyl)cyclohexylcarbonyl

-mab & -tox

anatumomab mafenatox (86)

immunoglobulin G 1, anti-(human tumor-associated glycoprotein 72) (human-mouse clone pMB125 Fab fragment γ 1-chain) fusion protein with enterotoxin A

(227-alanine) (*Staphylococcus aureus*) complex with mouse clone pMB125 κ -chain)

citatumumab bogatox (99)

immunoglobulin Fab fusion protein, anti-[*Homo sapiens* tumor-associated calcium signal transducer 1 (TACSTD1, gastrointestinal tumor-associated protein 2, GA733-2, epithelial glycoprotein 2, EGP-2, epithelial cell adhesion molecule Ep-CAM, KSA, KS1/4 antigen, M4S, tumor antigen 17-1A, CD326)], humanized Fab fused with *Bougainvillea spectabilis* Willd rRNA N-glycosidase [type I ribosome inactivating protein (RIP), bouganin], VB6-845; gamma1 heavy chain fragment (1-225) [hexahistidyl (1-6) -humanized VH from 4D5MOC-B (*Homo sapiens* FR/*Mus musculus* CDR, *Homo sapiens* IGHJ4*01, V124>L) [8.8.9] (7-122) -*Homo sapiens* IGHG1*01 CH1-hinge fragment EPKSC (123-225)], (225-219')-disulfide with kappa fusion chain (1'-481') [humanized V-KAPPA from clone 4D5MOC-B (*Homo sapiens* FR/*Mus musculus* CDR, *Homo sapiens* IGKJ1*01, I126>L) [11.3.9] (1'-112') -*Homo sapiens* IGKC*01 (113'-219') -12-mer furin linker (proteolytic cleavage spacer from *Pseudomonas* exotoxin A) (220'-231') -*Bougainvillea spectabilis* Willd bouganin fragment (27-276 from precursor, V354'>A, D358'>A, Y364'>N, I383'>A) (232'-481')]

dorlimomab aritox (66)

ricin A chain-antibody ST 1 F(ab')₂ fragment immunotoxin

moxetumomab pasudotox (102)

immunoglobulin Fv fragment fused to *Pseudomonas* toxin, anti-[*Homo sapiens* CD22 (sialic acid-binding Ig-like lectin 2, Siglec-2, SIGLEC2, Leu-14, B-lymphocyte cell adhesion molecule, BL-CAM)], *Mus musculus* monoclonal antibody disulfide stabilized Fv fragment with the variable heavy VH domain fused with the truncated form PE38 of *Pseudomonas aeruginosa* exotoxin A (VH-PE38), disulfide linked with the variable kappa domain (V-KAPPA)]; VH-PE38 (1-476) comprising the VH domain (1-123) [methionyl -*Mus musculus* VH [(IGHV5-12-1*01 -(IGHD)-IGHJ3*01) [8.8.16] (2-123)] fused with a 7-mer linker (124-130) and with the *Pseudomonas aeruginosa* exotoxin A (ETA) PE38 fragment (131-476) [277-638 precursor fragment with del 389-405>N (131-476), containing domain II (131-243) with furin proteolytic cleavage site (152-164), domain Ib (244-267), domain III (268-476)], (45-101')-disulfide with V-KAPPA (1'-108') [methionyl -*Mus musculus* V-KAPPA [(IGHKV10-96*01 -IGKJ1*01) [6.3.9] (2'-108')]

nacolomab tafenatox (80)

immunoglobulin G1, anti-(human colorectal tumor antigen C242) Fab fragment (mouse monoclonal r-C242Fab-SEA clone pkP941 γ 1-chain) fusion protein with enterotoxin A (*Staphylococcus aureus*), disulfide with mouse monoclonal r-C242Fab-SEA clone pkP941 κ -chain

naptumomab estafenatox (96)

immunoglobulin fragment, anti-[trophoblast glycoprotein (TPBG, 5T4)] monoclonal 5T4 gamma1 heavy chain fragment fusion protein [*Mus musculus* VH (5T4V14: H41>P, S44>G, I69>T, V113>G)-IGHG1_CH1] - [Glycyl-Glycyl-Prolyl] - superantigen SEA/E-120 (synthetic), non-disulfide linked with monoclonal 5T4 kappa light chain [*Mus musculus* V-KAPPA (5T4V18: F10>S, T45>K, I63>S, F73>L, T77>S, L78>V, L83>A)-IGKC]

oportuzumab monatox (100)

immunoglobulin scFv fusion protein, anti- [*Homo sapiens* tumor-associated calcium signal transducer 1 (TACSTD1, gastrointestinal tumor-associated protein 2, GA733-2, epithelial glycoprotein 2, EGP- 2, epithelial cell adhesion molecule Ep-CAM, KSA, KS1/4 antigen, M4S1, tumor antigen 17-1A, CD326)] humanized monoclonal antibody scFv fused with *Pseudomonas aeruginosa* exotoxin A; hexahistidyl -humanized scFv [V-KAPPA (*Homo sapiens* IGKV1- 39*01 (78%)-IGKJ1*01, I126>L) [11.3.9] (7-118) -26-mer linker -VH (*Homo sapiens* IGHV7-4-1*02 -(IGHD)-IGHJ4*01, V124>L) [8.8.9] (145-260)] -20-mer linker - *Pseudomonas aeruginosa* exotoxin A (ETA) [277-633 precursor fragment, containing domain II (281-393) with furin proteolytic cleavage site (302-313), domain Ib (394-433), domain III (434-637)] (281-637) -hexahistidyl-lysyl-aspartyl-glutamylleucyl

taplitumomab paptox (84)

immunoglobulin G1, anti-(human antigen CD19) (mouse monoclonal B43 γ 1-chain), disulfide with mouse monoclonal B43 κ -chain, dimer, disulfide with protein PAP (pokeweed antiviral)

telimomab aritox (66)

ricin A chain-antibody T 101 Fab fragment immunotoxin

zolimomab aritox (80)

immunoglobulin G1, anti-(human CD5 (antigen) heavy chain) (mouse monoclonal H65-RTA γ 1-chain), disulfide with mouse monoclonal H65-RTA light chain, dimer, disulfide with ricin (castor bean A-chain)

-tide

cenderitide (105)

natriuretic peptide receptor type B (NPR-B) agonist;
human C-type natriuretic peptide-(32-53)-peptide (CNP-22) fusion protein with eastern green mamba (*Dendroaspis angusticeps*) natriuretic peptide-(24-38)-peptide

elsiglutide (104)

[2-glycine(A>G),3-glutamic acid(D>E),8-serine(D>S),10-leucine(M>L),11-serine(N>S),16-alanine(N>A),24-alanine(N>A),28-alanine(Q>A)]human glucagon-like peptide 2 (GLP-2) fusion protein with hexalysinamide

langlenatide (109)

exenatide derivative and human IgG4 Fc dimer linked together with polyethylene glycol derivative:
 $N^{6,27},N^1$ -[ω -(oxypropane-1,3-diyl)- α -(propane-1,3-diyl)poly(oxyethylene)] [1-(imidazol-4-ylacetic acid)]exendin-4 *Heloderma suspectum* (Gila monster), human immunoglobulin G4 Fc fragment-(9'-229')-peptide dimer (3'-3'')-disulfide

vanutide cridificar (100)

inactivated diphtheria toxin (carrier) covalently linked to human beta-amyloid protein 42 short fragments: pentadecakis[N^{6-Lys} -(sulfanylacetyl)]-[52-glutamic acid(G>E)]diphtheria toxin *Corynebacterium diphtheriae* thioether with human beta-amyloid protein 42-(1-7)-peptidylcysteine

-motide

amilomotide (105)

virus like particle of bacteriophage Q-beta coat protein that is coupled to multiple copies of human beta-amyloid1-6 peptide fragment;
reaction products of bacteriophage Q-beta coat protein with human beta-amyloid protein-(1-6)-peptidylglycylglycyl-L-cysteine and 3-(2,5-dioxo-2,5-dihydro-1H-pyrrole-1-yl)-N-{6-[(2,5-dioxopyrrolidin-1-yl)oxy]-6-oxohexyl}propanamide

tecemotide (108)

human mucin-1 (carcinoma-associated mucin, episialin, CD227)-(107-131)-peptide (sequence 40 times repeated) fusion protein with 6-N-hexadecanoyl-L-lysylglycine

zastumotide (110)

19,137,308,342,395-penta[S-(2-amino-2-oxoethyl)]-[2-aspartic acid(K²>D),3-proline(L³>P)]glycerophosphoryl diester phosphodiesterase (*Haemophilus influenzae* strain 86-028NP EC 3.1.4.46)-(1-127)-peptide fusion protein with [2-aspartic acid(P²>D)]human melanoma-associated antigen 3 (MAGE-3 antigen, antigen MZ2-D, cancer/testis antigen 1.3 or CT1.3) fusion protein with diglycylheptahistidine }

CPCA with Fc

This item includes INN assigned to composite proteins for clinical applications (CPCA) (or peptides) fused with immunoglobulin Fc with the aim of increasing their half-life.

Before the *ef*-suffix was implemented:

asfotase alfa (104)

tissue-nonspecific alkaline phosphatase- IgG₁ fusion protein;human tissue-nonspecific isozyme alkaline phosphatase (AP-TNAP, EC=3.1.3.1) fusion protein with leucyl-lysyl-human immunoglobulin G1 Fc region {(6-15)-H-CH2-CH3 of IGHG1*03} fusion protein with aspartyl-isoleucyl-deca(aspartic acid), dimer (493-493':496-496')-bisdisulfide

blisibimod (107)

B-cell activating factor (BAFF)-binding peptide fragment/human IgG1 Fc fusion protein

dulaglutide (103)

glucagon-like peptide-1-immunoglobulin G4 fusion protein, [2-glycyl,16-L-glutamyl,30-glycyl][human glucagon-like peptide 1-(7-37)-peptide] {(8-A>G,22-G>E,36-R>G)-GLP-1(7-37)} fusion protein with tris(tetraglycyl-L-seryl)-L-alanine (linker) fusion protein with des-276-lysine-[57-L-proline,63-L-alanine,64-L-alanine]human immunoglobulin G4 Fc region {(10-S>P)-H-(4-F>A,5-L>A)-CH2-(107-K>-)-CH3 of IGHG4*01}, dimer (55-55':58-58')-bisdisulfide

romiplostim (97)

L-methionyl[human immunoglobulin heavy constant gamma 1-(227 C-terminal residues)-peptide (Fc fragment)] fusion protein with 41 amino acids peptide, (7-7':10,10')-bisdisulfide dimer

torapsel (91)

42-89-glycoprotein (human clone PMT21:PL85 P-selectin glycoprotein ligand fusion protein with immunoglobulin (human constant region)

trebananib (106)

immunoglobulin G1 Fc fragment fused with two synthetic polypeptides that bind the *Homo sapiens* ANGPT2 (angiopoietin 2); methionyl (1) -gamma1 heavy chain fragment (2-228) [*Homo sapiens* IGHG1*01 hinge (EPKSC 1-5>del) (2-11), CH2 (12-121), CH3 (122-228)] fused, at the C-terminal end, with a synthetic polypeptide that comprises two 14-mer amino acid repeats that bind angiopoietin 2 (229-287) [linker (229-235) -14-mer (236-249) – linker (250-271) -14-mer (272-285) -leucyl-glutamate]; (7-7':10-10')-bisdisulfide dimer

ef-

eflapgrastim (111)

human granulocyte colony-stimulating factor and human IgG4 Fc dimer linked together with polyethylene glycol derivative, produced in *Escherichia coli*:
Na.1,N1.9'-[ω-(oxypropane-1,3-diyl)-α-(propane-1,3-diyl)poly(oxyethylene)]
des-(1-L-alanine,37-39)-[18-L-serine(C>S)]human granulocyte colony-
stimulating factor (G-CSF, pluripointin) (1-174)-peptide and des-(1-8)-human
immunoglobulin G4 Fc fragment (IGHG4*01 H-CH2-CH3) (1'-221')-peptide
dimer (11'-11'')-disulfide

efmoroctocog alfa (111)

recombinant DNA derived (1-742)-(1637-2332)-human blood coagulation factor
VIII fusion protein with immunoglobulin G1 Fc domain fragment, produced in
HEK293H cells, glycoform alfa:
des-(743-1636)-human blood coagulation factor VIII (antihemophilic factor,
procoagulant component) fusion protein with human immunoglobulin G1 Fc
fragment (IGHG1*01 H-CH2-CH3)-(6-231)-peptide (1444-6':1447-9')-
bisdisulfide with human immunoglobulin G1 Fc fragment (IGHG1*01 H-CH2-
CH3)-(6-231)-peptide

eftrenonacog alfa (109)

recombinant DNA derived human blood coagulation factor IX fusion protein with
one Fc fragment of the human immunoglobulin G1 Fc fragment dimer, produced
in HEK293H cells (glycoform alfa):
human blood coagulation factor IX (EC 3.4.21.22, Christmas factor, plasma
thromboplastin component) variant 148-T, fusion protein with human
immunoglobulin G1 Fc fragment (IGHG1*01 H-CH2-CH3)-(6-231)-peptide
(421-6':424-9')-bisdisulfide with human immunoglobulin G1 Fc fragment
(IGHG1*01 H-CH2-CH3)-(6-231)-peptide

Others

dianexin (109)

recombinant DNA derived annexin A5 dimer covalently linked by a 14 residues
peptide linker, produced in *Escherichia coli* (nonglycosylated):
L-methionyl-human annexin A5 fusion protein with glycyl-L-seryl-L-leucyl-L-α-
glutamyl-L-valyl-L-leucyl-L-phenylalanyl-L-glutaminyglycyl-L-prolyl-L-
serylglycyl-L-lysyl-L-leucyl-human annexin A5

mipsagargin (110)

sarcoplasmic/endoplasmic reticulum Ca²⁺ dependent ATPase (SERCA) inhibitor conjugated to a peptide targeting prostate-specific membrane antigen (PSMA):
N⁴-(12-[[[(3*S*,3*aR*,4*S*,6*S*,6*aR*,7*S*,8*S*,9*bS*)-6-(acetyloxy)-3,3*a*-dihydroxy-3,6,9-trimethyl-8-[[[(2*Z*)-2-methylbut-2-enoyl]oxy]-7-(octanoyloxy)-2-oxo-2,3,3*a*,4,5,6,6*a*,7,8,9*b*-decahydroazuleno[4,5-*b*]furan-4-yl]oxy]-12-oxododecyl)-L-asparaginyl-L-γ-glutamyl-L-γ-glutamyl-L-γ-glutamyl-L-glutamic acid

topsalysin (111)

recombinant DNA derived proaerolysin, pore-forming protein, from *Aeromonas hydrophila*, with the furin site substituted with a prostate specific antigen (PSA) cleavage site, fusion protein with 6 histidines, produced in *Escherichia coli* (nonglycosylated):
[427-L-histidine(K>H),428-L-serine(V>S),429-L-serine(R>S),430-L-lysine(R>K),431-L-leucine(A>L),432-L-glutamine(R>Q)]proaerolysin *Aeromonas hydrophila* fusion protein with hexa-L-histidine

transferrin aldifitox (95)

a conjugate of the precursor of human serotransferrin (siderophillin) with a primary amine group used to form an amidine with (4-iminobutane-1,4-diyl)sulfanediyl[(3*RS*)-2,5-dioxopyrrolidine-1,3-diyl]-1,3-phenylenecarbonyl and forming an *N*-benzoyl derivative of a primary amine group of diphtheria [550-L-phenylalanine]toxin from *Corynebacterium diphtheriae*-(26-560)-peptide

verpasep caltespen (95)

60 kDa chaperonin 2 (heat shock protein 65 from *Mycobacterium bovis* strain BCG) fusion protein with L-histidylprotein E7 from human papillomavirus type 16.

zoptarelin doxorubicin (107)

[6-D-lysine]human gonadoliberin-1 (LHRH) and doxorubicin covalently linked together with glutaric acid:
5-oxo-L-prolyl-L-histidyl-L-tryptophyl-L-seryl-L-tyrosyl-N⁶-[5-(2-[(2*S*,4*S*)-4-[(3-amino-2,3,6-trideoxy-α-L-*lyxo*-hexopyranosyl)oxy]-2,5,12-trihydroxy-7-methoxy-6,11-dioxo-1,2,3,4,6,11-hexahydrotetracen-2-yl]-2-oxoethoxy)-5-oxopentanoyl]-D-lysine-L-leucyl-L-arginyl-L-prolylglycinamide

ANNEX 2

Transliteration of Greek letters in English, French and Spanish

Upper case	Lower case	English	French	Spanish
A	α	alfa (and not alpha)	alfa (and not alpha)	alfa
B	β	beta	bêta	beta
Γ	γ	gamma	gamma	gamma
Δ	δ	delta	delta	delta
E	ε	epsilon	epsilon	épsilon
Z	ζ	zeta	zêta	<u>d</u>seta
H	η	eta	êta	eta
Θ	θ	theta	thêta	<u>z</u>eta
I	ι	iota	iota	iota
K	κ	kappa	kappa	kappa
Λ	λ	lambda	lambda	lambda
M	μ	mu	mu	mi
N	ν	nu	nu	ni
Ξ	ξ	xi	xi	xi
O	ο	omicron	omicron	ómicron
Π	π	pi	pi	pi
P	ρ	rho	rhô	ro
Σ	σ	sigma	sigma	sigma
T	τ	tau	tau	tau
Υ	υ	upsilon	upsilon	ípsilon
Φ	φ	phi	phi	fi
X	χ	chi	khi	ji
Ψ	ψ	psi	psi	psi
Ω	ω	omega	oméga	omega

* letters to be avoided

The previous naming scheme for monoclonal antibodies

General policies for monoclonal antibodies

- The common stem for monoclonal antibodies is *-mab*.
- Sub-stems for source of product:

<i>a</i>	rat
<i>axo (pre-sub-stem)</i>	rat-murine hybrid
<i>e</i>	hamster
<i>i</i>	primate
<i>o</i>	mouse
<i>u</i>	human
<i>xi</i>	chimeric
<i>zu</i>	humanized

The distinction between chimeric and humanized antibodies is as follows:

A chimeric antibody is one that contains contiguous foreign-derived amino acids comprising the entire variable region of both heavy and light chains linked to heavy and light constant regions of human origin.

A humanized antibody has segments of foreign-derived amino acids interspersed among variable region segments of human-derived amino acid residues and the humanized heavy-variable and light-variable regions are linked to heavy and light constant regions of human origin.

- Sub-stems for disease or target class:

<i>-ba(c)-</i>	bacterial
<i>-ci(r)-</i>	cardiovascular
<i>-fung-</i>	fungal
<i>-ki(n)- (pre-sub-stem)</i>	interleukin
<i>-le(s)-</i>	inflammatory lesions
<i>-li(m)-</i>	immunomodulator
<i>-os-</i>	bone
<i>-vi(r)-</i>	viral

tumours:

<i>-co(l)-</i>	colon
<i>-go(t)-</i>	testis
<i>-go(v)-</i>	ovary
<i>-ma(r)-</i>	mammary
<i>-me(l)-</i>	melanoma
<i>-pr(o)-</i>	prostate
<i>-tu(m)-</i>	miscellaneous

Whenever there is a problem in pronunciation, the final letter of the sub-stems for diseases or targets may be deleted, e.g. *-vi(r)-*, *-ba(c)-*, *-li(m)-*, *-co(l)-*, etc.

Prefix

Should be random e.g. the only requirement is to contribute to a euphonious and distinctive name.

Second word

If the product is radiolabelled or conjugated to another chemical, such as toxin, identification of this conjugate is accomplished by use of a separate, second word or acceptable chemical designation.

If the monoclonal antibody is used as a carrier for a radioisotope, the latter will be listed first in the INN, e.g. *technetium (^{99m}Tc) pintumomab (86)*.

-toxa- infix

For monoclonals conjugated to a toxin, the infix *-toxa-* can be inserted either into the first (main) name or included in the second word.