

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



WIPO | PCT



(10) International Publication Number  
**WO 2014/012007 A2**

(43) International Publication Date  
16 January 2014 (16.01.2014)

- (51) International Patent Classification:  
*C07K 16/18* (2006.01) *A61K 45/06* (2006.01)  
*A61K 39/395* (2006.01) *C12Q 1/68* (2006.01)
- (21) International Application Number:  
PCT/US2013/050300
- (22) International Filing Date:  
12 July 2013 (12.07.2013)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
61/671,421 13 July 2012 (13.07.2012) US  
61/753,184 16 January 2013 (16.01.2013) US  
61/789,156 15 March 2013 (15.03.2013) US  
61/826,747 23 May 2013 (23.05.2013) US
- (71) Applicant: **ONCOMED PHARMACEUTICALS, INC.**  
[US/US]; 800 Chesapeake Drive, Redwood City, California 94063 (US).
- (72) Inventors: **GURNEY, Austin L.**; 946 Diamond Street, San Francisco, California 94114 (US). **BOND, Christopher J.**; 511 30th Avenue, San Mateo, California 94403 (US).
- (74) Agents: **CALVO, Paul A.** et al.; Sterne, Kessler, Goldstein & Fox, PLLC, 1100 New York Ave., NW, Washington, DC 20005 (US).

- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*
- *with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))*
- *with sequence listing part of description (Rule 5.2(a))*

(54) Title: RSPO3 BINDING AGENTS AND USES THEREOF

(57) Abstract: The present invention relates to RSPO-binding agents, particularly RSPO3-binding agents and methods of using the agents for treating diseases such as cancer. The present invention provides antibodies that specifically bind human RSPO3 proteins and modulate  $\beta$ -catenin activity. The present invention further provides methods of using agents that modulate the activity of RSPO3 proteins and inhibit tumor growth. Also described are methods of treating cancer comprising administering a therapeutically effect amount of an agent or antibody of the present invention to a patient having a tumor or cancer.



WO 2014/012007 A2

## RSPO3 BINDING AGENTS AND USES THEREOF

## CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims priority benefit of U.S. Provisional Application No. 61/671,421, filed July 13, 2012, U.S. Provisional Application No. 61/753,184, filed January 16, 2013, U.S. Provisional Application No. 61/789,156, filed March 15, 2013, and U.S. Provisional Application No. 61/826,747, filed May 23, 2013, each of which is hereby incorporated by reference herein in its entirety.

## FIELD OF THE INVENTION

[002] The field of this invention generally relates to antibodies and other agents that bind R-Spondin proteins (RSPO), particularly human R-Spondin protein RSPO3, as well as to methods of using the antibodies or other agents for the treatment of diseases such as cancer.

## BACKGROUND OF THE INVENTION

[003] The R-Spondin (RSPO) family of proteins is conserved among vertebrates and comprises four members, RSPO1, RSPO2, RSPO3, and RSPO4. These proteins have been referred to by a variety of names, including roof plate-specific spondins, hPWTSR (hRSPO3), THS2D (RSPO3), Cristin 1-4, and Futrin 1-4. The RSPOs are small secreted proteins that overall share approximately 40-60% sequence homology and domain organization. All RSPO proteins contain two furin-like cysteine-rich domains at the N-terminus followed by a thrombospondin domain and a basic charged C-terminal tail (Kim et al., 2006, *Cell Cycle*, 5:23-26).

[004] Studies have shown that RSPO proteins have a role during vertebrate development (Kamata et al., 2004, *Biochim. Biophys. Acta*, 1676:51-62) and in *Xenopus* myogenesis (Kazanskaya et al., 2004, *Dev. Cell*, 7:525-534). RSPO1 has also been shown to function as a potent mitogen for gastrointestinal epithelial cells (Kim et al., 2005, *Science*, 309:1256-1259). It has been reported that RSPO3 is prominently expressed in or close by endothelial cells and their cellular precursors in *Xenopus* and mouse. Furthermore, it has been suggested that RSPO3 may act as an angiogenic factor in embryogenesis (Kazanskaya et al., 2008, *Development*, 135:3655-3664). RSPO proteins are known to activate  $\beta$ -catenin signaling similar to Wnt signaling, however the relationship between RSPO proteins and Wnt signaling is still being investigated. It has been reported that RSPO proteins possess a positive modulatory activity on Wnt ligands (Nam et al., 2006, *JBC* 281:13247-57). This study also reported that RSPO proteins could function as Frizzled8 and LRP6 receptor ligands and induce  $\beta$ -catenin signaling (Nam et al., 2006, *JBC* 281:13247-57). Recent studies have identified an interaction between RSPO proteins and LGR (leucine-rich repeat containing, G protein-coupled receptor) proteins, such as LGR5 (U.S. Patent Publication Nos.

2009/0074782 and 2009/0191205), and these data present an alternative pathway for the activation of  $\beta$ -catenin signaling.

[005] The Wnt signaling pathway has been identified as a potential target for cancer therapy. The Wnt signaling pathway is one of several critical regulators of embryonic pattern formation, post-embryonic tissue maintenance, and stem cell biology. More specifically, Wnt signaling plays an important role in the generation of cell polarity and cell fate specification including self-renewal by stem cell populations. Unregulated activation of the Wnt pathway is associated with numerous human cancers where it is believed the activation can alter the developmental fate of cells. The activation of the Wnt pathway may maintain tumor cells in an undifferentiated state and/or lead to uncontrolled proliferation. Thus carcinogenesis can proceed by overtaking homeostatic mechanisms which control normal development and tissue repair (reviewed in Reya & Clevers, 2005, *Nature*, 434:843-50; Beachy et al., 2004, *Nature*, 432:324-31).

[006] The Wnt signaling pathway was first elucidated in the *Drosophila* developmental mutant wingless (wg) and from the murine proto-oncogene int-1, now Wnt1 (Nusse & Varmus, 1982, *Cell*, 31:99-109; Van Ooyen & Nusse, 1984, *Cell*, 39:233-40; Cabrera et al., 1987, *Cell*, 50:659-63; Rijsewijk et al., 1987, *Cell*, 50:649-57). Wnt genes encode secreted lipid-modified glycoproteins of which 19 have been identified in mammals. These secreted ligands activate a receptor complex consisting of a Frizzled (FZD) receptor family member and low-density lipoprotein (LDL) receptor-related protein 5 or 6 (LRP5/6). The FZD receptors are seven transmembrane domain proteins of the G-protein coupled receptor (GPCR) superfamily and contain a large extracellular N-terminal ligand binding domain with 10 conserved cysteines, known as a cysteine-rich domain (CRD) or Fri domain. There are ten human FZD receptors, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, and FZD10. Different FZD CRDs have different binding affinities for specific Wnt proteins (Wu & Nusse, 2002, *J. Biol. Chem.*, 277:41762-9), and FZD receptors have been grouped into those that activate the canonical  $\beta$ -catenin pathway and those that activate non-canonical pathways (Miller et al., 1999, *Oncogene*, 18:7860-72).

[007] A role for Wnt signaling in cancer was first uncovered with the identification of Wnt1 (originally int1) as an oncogene in mammary tumors transformed by the nearby insertion of a murine virus (Nusse & Varmus, 1982, *Cell*, 31:99-109). Additional evidence for the role of Wnt signaling in breast cancer has since accumulated. For instance, transgenic over-expression of  $\beta$ -catenin in the mammary glands results in hyperplasias and adenocarcinomas (Imbert et al., 2001, *J. Cell Biol.*, 153:555-68; Michaelson & Leder, 2001, *Oncogene*, 20:5093-9) whereas loss of Wnt signaling disrupts normal mammary gland development (Tepera et al., 2003, *J. Cell Sci.*, 116:1137-49; Hatsell et al., 2003, *J. Mammary Gland Biol. Neoplasia*, 8:145-58). In human breast cancer,  $\beta$ -catenin accumulation implicates activated Wnt signaling in over 50% of carcinomas, and though specific mutations have not been identified, up-regulation of Frizzled

receptor expression has been observed (Brennan & Brown, 2004, *J. Mammary Gland Biol. Neoplasia*, 9:119-31; Malovanovic et al., 2004, *Int. J. Oncol.*, 25:1337-42).

[008] Activation of the Wnt pathway is also associated with colorectal cancer. Approximately 5-10% of all colorectal cancers are hereditary with one of the main forms being familial adenomatous polyposis (FAP), an autosomal dominant disease in which about 80% of affected individuals contain a germline mutation in the adenomatous polyposis coli (APC) gene. Mutations have also been identified in other Wnt pathway components including Axin and  $\beta$ -catenin. Individual adenomas are clonal outgrowths of epithelial cells containing a second inactivated allele, and the large number of FAP adenomas inevitably results in the development of adenocarcinomas through additional mutations in oncogenes and/or tumor suppressor genes. Furthermore, activation of the Wnt signaling pathway, including loss-of-function mutations in APC and stabilizing mutations in  $\beta$ -catenin, can induce hyperplastic development and tumor growth in mouse models (Oshima et al., 1997, *Cancer Res.*, 57:1644-9; Harada et al., 1999, *EMBO J.*, 18:5931-42).

[009] Similar to breast cancer and colon cancer, melanoma often has constitutive activation of the Wnt pathway, as indicated by the nuclear accumulation of  $\beta$ -catenin. Activation of the Wnt/ $\beta$ -catenin pathway in some melanoma tumors and cell lines is due to modifications in pathway components, such as APC, ICAT, LEF1 and  $\beta$ -catenin (see e.g., Larue et al. 2006, *Frontiers Biosci.*, 11:733-742). However, there are conflicting reports in the literature as to the exact role of Wnt/ $\beta$ -catenin signaling in melanoma. For example, one study found that elevated levels of nuclear  $\beta$ -catenin correlated with improved survival from melanoma, and that activated Wnt/ $\beta$ -catenin signaling was associated with decreased cell proliferation (Chien et al., 2009, *PNAS*, 106:1193-1198).

[010] The focus of cancer drug research is shifting toward targeted therapies aimed at genes, proteins, and pathways involved in human cancer. There is a need for new agents targeting signaling pathways and new combinations of agents that target multiple pathways that could provide therapeutic benefit for cancer patients. Thus, biomolecules (e.g., anti-RSPO3 antibodies) that disrupt  $\beta$ -catenin signaling are a potential source of new therapeutic agents for cancer, as well as other  $\beta$ -catenin-associated diseases.

#### BRIEF SUMMARY OF THE INVENTION

[011] The present invention provides binding agents, such as antibodies, that bind RSPO3 proteins, as well as compositions, such as pharmaceutical compositions, comprising the binding agents. Binding agents that bind RSPO3 as well as at least one additional antigen or target, and pharmaceutical compositions of such binding agents, are also provided. In certain embodiments, the RSPO3-binding agents are novel polypeptides, such as antibodies, antibody fragments, and other polypeptides related to such antibodies. The invention further provides methods of inhibiting the growth of a tumor by administering the RSPO3-binding agents to a subject with a tumor. The invention further provides



methods of treating cancer by administering the RSPO3-binding agents to a subject in need thereof. In some embodiments, the methods of treating cancer or inhibiting tumor growth comprise targeting cancer stem cells with the RSPO3-binding agents. In some embodiments, the methods comprise disrupting  $\beta$ -catenin signaling. In some embodiments, the methods comprise modulating (e.g., inhibiting) angiogenesis. In certain embodiments, the methods comprise reducing the frequency of cancer stem cells in a tumor, reducing the number of cancer stem cells in a tumor, reducing the tumorigenicity of a tumor, and/or reducing the tumorigenicity of a tumor by reducing the number or frequency of cancer stem cells in the tumor.

**[012]** In one aspect, the invention provides a binding agent, such as an antibody, that specifically binds human RSPO3. The sequence of human RSPO3 is known in the art and is included herein as SEQ ID NO:3. In certain embodiments, the RSPO3-binding agent binds within amino acids 22-272 of human RSPO3. In certain embodiments, the RSPO3-binding agent binds within amino acids 22-207 of human RSPO3. In certain embodiments, the RSPO3-binding agent binds within amino acids 35-135 of human RSPO3. In certain embodiments, the RSPO3-binding agent binds within amino acids 35-86 of human RSPO3. In certain embodiments, the RSPO3-binding agent binds within amino acids 92-135 of human RSPO3. In some embodiments, the RSPO3-binding agent (e.g., an antibody) specifically binds at least one other human RSPO selected from the group consisting of RSPO1, RSPO2, and RSPO4. In some embodiments, the RSPO3-binding agent or antibody modulates  $\beta$ -catenin activity, is an antagonist of  $\beta$ -catenin signaling, inhibits  $\beta$ -catenin signaling, and/or inhibits activation of  $\beta$ -catenin. In some embodiments, the RSPO3-binding agent inhibits RSPO3 signaling. In some embodiments, the RSPO3-binding agent inhibits, interferes with, and/or disrupts binding of RSPO3 to one or more LGR proteins (e.g., LGR4, LGR5, and/or LGR6). In some embodiments, the RSPO3-binding agent inhibits binding of RSPO3 to LGR5.

**[013]** In certain embodiments, the RSPO3-binding agent is an antibody which binds human RSPO3. In some embodiments, the antibody binds human RSPO3 and mouse RSPO3. In certain embodiments, the antibody comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments, the antibody further comprises a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83). In some embodiments, the antibody comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY

(SEQ ID NO:35), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14). In some embodiments, the antibody comprises a heavy chain CDR1 comprising DYSH (SEQ ID NO:78), a heavy chain CDR2 comprising YIPSNNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLTF (SEQ ID NO:83). In some embodiments, the antibody comprises a heavy chain CDR1 comprising DYSH (SEQ ID NO:78), a heavy chain CDR2 comprising YIPSNNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14). In some embodiments, the antibody comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14).

**[014]** In certain embodiments, the RSPO3-binding agent is an antibody which comprises: (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), DYSH (SEQ ID NO:78), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (b) a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), YIPSNNGDSGYNQKFK (SEQ ID NO:79), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (c) a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), TYFANNFD (SEQ ID NO:80), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (d) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), KASQSVDDYDGDSYMN (SEQ ID NO:81), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (e) a light chain CDR2 comprising AAS (SEQ ID NO:13), AASNLES (SEQ ID NO:82), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; and (f) a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14), QQSNEPLTF (SEQ ID NO:83), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions. In some embodiments, the amino acid substitutions are conservative amino acid substitutions. In some embodiments, the substitutions are made as part of a germline humanization process.

**[015]** In certain embodiments, the RSPO3-binding agent is an antibody which comprises: (a) a heavy chain variable region having at least 80% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62; and/or (b) a light chain variable region having at least 80% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID

NO:86. In certain embodiments, the RSPO3-binding agent is an antibody that comprises: (a) a heavy chain variable region having at least 90% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62; and/or (b) a light chain variable region having at least 90% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.

**[016]** In some embodiments, the RSPO3-binding agent is a monoclonal antibody. In some embodiments, the monoclonal antibody is an IgG1 antibody. In some embodiments, the monoclonal antibody is an IgG2 antibody. In some embodiments, the RSPO3-binding agent is monoclonal antibody 131R002 or monoclonal antibody 131R003. In some embodiments, the RSPO3-binding agent is an affinity-matured variant of monoclonal antibody 131R002 or monoclonal antibody 131R003. In some embodiments, the RSPO3-binding agent is a chimeric antibody comprising the antigen-binding sites from antibody 131R002 or antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized form of antibody 131R002 or antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011.

**[017]** In another aspect, the invention provides a binding agent (e.g., an antibody) that competes for specific binding to human RSPO3 with an antibody of the invention. In some embodiments, the binding agent (e.g., an antibody) competes for specific binding to human RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In some embodiments, the antibody with which the RSPO3-binding agent competes is antibody 131R002 or antibody 131R003. In some embodiments, the antibody with which the RSPO3-binding agent competes is a humanized form of antibody 131R002 or antibody 131R003. In some embodiments, the antibody with which the RSPO3-binding agent competes is antibody h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011. In some embodiments, the binding agent competes for specific binding to RSPO3 with an antibody of the invention in an *in vitro* competitive binding assay.

**[018]** In certain embodiments, the binding agent is an antibody that binds the same epitope, or essentially the same epitope, on RSPO3 as an antibody of the invention (e.g., 131R002, 131R003, or humanized forms/variants thereof). In certain embodiments, the binding agent is an antibody that binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R005/131R007, h131R008, h131R010, or h131R011.

**[019]** In still another aspect, the binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by an antibody of the invention (e.g., 131R002, 131R003, or humanized forms/variants thereof). In some embodiments, the binding agent is an antibody that binds an

epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R005/131R007, h131R008, h131R010, or h131R011.

[020] In certain embodiments of each of the aforementioned aspects or embodiments, as well as other aspects and/or embodiments described elsewhere herein, the binding agent is a bispecific antibody. In some embodiments, the bispecific antibody specifically binds human RSPO3 and a second target. In some embodiments, the bispecific antibody specifically binds human RSPO3 and human RSPO1. In some embodiments, the bispecific antibody specifically binds human RSPO3 and human RSPO2. In some embodiments, the bispecific antibody specifically binds human RSPO3 and human RSPO4. In some embodiments, the bispecific antibody modulates  $\beta$ -catenin activity. In certain embodiments, the bispecific antibody inhibits  $\beta$ -catenin activity. In certain embodiments, the bispecific antibody inhibits  $\beta$ -catenin signaling. In certain embodiments, the bispecific antibody inhibits activation of  $\beta$ -catenin. In some embodiments, the bispecific antibody reduces the number of frequency of cancer stem cells. In certain embodiments, the bispecific antibody comprises two identical light chains. In certain embodiments, the bispecific antibody is an IgG antibody. In certain embodiments, the bispecific antibody is an IgG1 antibody. In certain embodiments, the bispecific antibody is an IgG2 antibody.

[021] In some embodiments, the bispecific antibody comprises: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments, the first antigen-binding site comprises a light chain CDR1 comprising QSVDYDGDSYM (SEQ ID NO:12) or KASQSVDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14) or QQSNEPLTF (SEQ ID NO:83). In some embodiments, the bispecific antibody further comprises a second antigen-binding site that specifically binds human RSPO1. In some embodiments, the bispecific antibody further comprises a second antigen-binding site that specifically binds human RSPO2. Non-limiting examples of antibodies to RSPO1 or antibodies to RSPO2 have been described in, for example, International Patent Application Pub. No. WO 2013/012747. In some embodiments, the first and second binding sites comprise a common (e.g., identical) light chain.

[022] In some embodiments, the bispecific antibody comprises: a) a first antigen-binding site that specifically binds human RSPO3, and b) a second antigen-binding site that specifically binds human RSPO1, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID

NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments, the bispecific antibody comprises: a) a first antigen-binding site that specifically binds human RSPO3, and b) a second antigen-binding site that specifically binds human RSPO2, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments, the first antigen-binding site comprises a light chain CDR1 comprising QSVDYDGDSYM (SEQ ID NO:12) or KASQSVDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83).

**[023]** In some embodiments, the bispecific antibody specifically binds human RSPO3 and comprises: a heavy chain variable region having at least 90% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62. In some embodiments, the bispecific antibody specifically binds human RSPO3 and comprises: a heavy chain variable region having at least 95% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62. In some embodiments, the bispecific antibody comprises a first and second binding site, wherein the first and second binding sites comprise a common (e.g., identical) light chain. In some embodiments, the bispecific antibody comprises a light chain variable region having at least 95% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.

**[024]** In certain embodiments of each of the aforementioned aspects, as well as other aspects and/or embodiments described elsewhere herein, the RSPO3-binding agent or antibody is isolated. In some embodiments, the RSPO3-binding agent or antibody is substantially pure.

**[025]** In another aspect, the invention provides polypeptides. In some embodiments, the polypeptide comprises a sequence selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:86, SEQ ID NO:87, and SEQ ID NO:88. In some embodiments, the polypeptide comprises SEQ ID NO:15 and/or SEQ ID NO:17. In some embodiments, the polypeptide comprises SEQ ID NO:16 and/or SEQ ID NO:17. In some embodiments, the polypeptide comprises SEQ ID NO:36 and/or SEQ ID NO:17. In some embodiments, the polypeptide comprises SEQ



comprises SEQ ID NO:48 and/or SEQ ID NO:88. In some embodiments, the polypeptide comprises SEQ ID NO:49 and/or SEQ ID NO:88. In some embodiments, the polypeptide comprises SEQ ID NO:64 and/or SEQ ID NO:88. In some embodiments, the polypeptide comprises SEQ ID NO:69 and/or SEQ ID NO:88.

[028] In some embodiments, the polypeptide is isolated. In certain embodiments, the polypeptide is substantially pure. In certain embodiments, the polypeptide is an antibody or part of any antibody, such as an antibody fragment.

[029] In another aspect, the invention provides isolated polynucleotide molecules comprising a polynucleotide that encodes the antibodies and/or polypeptides of each of the aforementioned aspects, as well as other aspects and/or embodiments described herein. In some embodiments, the polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, and SEQ ID NO:95. In some embodiments, the polynucleotide comprises a polynucleotide that encodes a polypeptide selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:86, SEQ ID NO:87, and SEQ ID NO:88.

[030] The invention further provides expression vectors that comprise the polynucleotides, as well as cells that comprise the expression vectors and/or the polynucleotides. In some embodiments, the cell is a hybridoma cell line. In some embodiments, the cell is a monoclonal cell line. In some embodiments, the cell is a prokaryotic cell. In some embodiments, the cell is an eukaryotic cell.

[031] In other aspects, the invention provides methods of inhibiting growth of a tumor, comprising contacting the tumor with an effective amount of a RSPO3-binding agent or antibody, including each of those described herein.

[032] In another aspect, the invention provides a method of inhibiting the growth of a tumor in a subject, comprising administering to the subject a therapeutically effective amount of a RSPO3-binding agent or antibody, including each of those described herein.

[033] In another aspect, the invention provides a method of inhibiting  $\beta$ -catenin signaling in a cell, comprising contacting the cell with an effective amount of a RSPO3-binding agent or antibody, including

each of those described herein. In some embodiments, the cell is a tumor cell. In some embodiments, the tumor is a colorectal tumor. In some embodiments, the tumor is an ovarian tumor. In some embodiments, the tumor is a pancreatic tumor. In some embodiments, the tumor is a lung tumor. In some embodiments, the tumor is a breast tumor. In some embodiments, the tumor expresses elevated levels of at least one RSPO protein. In some embodiments, the tumor expresses elevated levels of RSPO1. In some embodiments, the tumor expresses elevated levels of RSPO2. In some embodiments, the tumor expresses elevated levels of RSPO3. In some embodiments, the tumor expresses a high level of at least one RSPO protein. In some embodiments, the tumor expresses a high level of RSPO1. In some embodiments, the tumor expresses a high level of RSPO2. In some embodiments, the tumor expresses a high level of RSPO3. In certain embodiments, the RSPO3-binding agent inhibits growth of the tumor, for example, by reducing the number and/or frequency of cancer stem cells in the tumor. In some embodiments, the tumor contains a RSPO gene fusion. In some embodiments, the tumor contains a RSPO2 gene fusion. In some embodiments, the tumor contains a RSPO3 gene fusion.

**[034]** In another aspect, the invention provides methods of treating cancer in a subject. In some embodiments, the method comprises administering to a subject a therapeutically effective amount of any of the RSPO3-binding agents or antibodies described above, as well as those described elsewhere herein. In some embodiments, the cancer is pancreatic cancer. In some embodiments, the cancer is colorectal cancer. In some embodiments, the colorectal cancer comprises an inactivating mutation in the adenomatous polyposis coli (APC) gene. In some embodiments, the colorectal cancer does not comprise an inactivating mutation in the APC gene. In some embodiments, the colorectal cancer comprises a wild-type APC gene. In some embodiments, the colorectal cancer comprises a RSPO gene fusion. In some embodiments, the colorectal cancer comprises a RSPO2 gene fusion. In some embodiments, the colorectal cancer comprises a RSPO3 gene fusion. In some embodiments, the cancer is ovarian cancer. In some embodiments, the cancer is lung cancer. In some embodiments, the cancer is breast cancer. In some embodiments, the cancer expresses elevated levels of at least one RSPO protein. In some embodiments, the cancer is an ovarian cancer that expresses elevated levels of RSPO3. In some embodiments, the cancer is lung cancer that expresses elevated levels of RSPO3. In some embodiments, the cancer is breast cancer that expresses elevated levels of RSPO3. In some embodiments, the cancer is pancreatic cancer that expresses elevated levels of RSPO3.

**[035]** In another aspect, the invention provides methods of treating a disease in a subject wherein the disease is associated with activation of  $\beta$ -catenin, increased  $\beta$ -catenin signaling, and/or aberrant  $\beta$ -catenin signaling, wherein the method comprises administering to the subject a therapeutically effective amount of a RSPO3-binding agent or antibody, including each of those described herein.



[036] In certain embodiments of each of the aforementioned aspects, as well as other aspects and/or embodiments described elsewhere herein, the treatment methods further comprise a step of determining the expression level of at least one RSPO protein in the tumor or cancer.

[037] In another aspect, the invention provides a method of identifying a human subject or selecting a human subject for treatment with a RSPO3-binding agent or antibody, including but not limited to, each of those described herein. In some embodiments, the method comprises determining if the subject has a tumor that has an elevated expression level of a specific RSPO (e.g., RSPO3) as compared to the expression of the same RSPO protein in normal tissue or to a pre-determined level of the same RSPO protein. In some embodiments, the method comprises identifying a subject for treatment or selecting a subject for treatment if the tumor has an elevated level of RSPO expression. In some embodiments, the method comprises determining if the subject has a tumor that comprises an inactivating mutation in the APC gene. In some embodiments, the method comprises identifying a subject for treatment or selecting a subject for treatment if the tumor comprises an inactivating mutation in the APC gene. In some embodiments, the method comprises determining if the subject has a tumor that comprises a RSPO gene fusion (e.g., a RSPO3 gene fusion). In some embodiments, the method comprises identifying a subject for treatment or selecting a subject for treatment if the tumor comprises a RSPO gene fusion (e.g., a RSPO3 gene fusion).

[038] In certain embodiments of each of the aforementioned aspects, as well as other aspects and/or embodiments described elsewhere herein, the treatment methods comprise administering to the subject the RSPO3-binding agent and at least one additional therapeutic agent.

[039] Pharmaceutical compositions comprising a RSPO3-binding agent or antibody described herein and a pharmaceutically acceptable carrier are further provided, as are cell lines that produce the RSPO3-binding agents. Methods of treating cancer and/or inhibiting tumor growth in a subject (e.g., a human) comprising administering to the subject an effective amount of a pharmaceutical composition comprising the RSPO3-binding agents are also provided.

[040] Where aspects or embodiments of the invention are described in terms of a Markush group or other grouping of alternatives, the present invention encompasses not only the entire group listed as a whole, but also each member of the group individually and all possible subgroups of the main group, and also the main group absent one or more of the group members. The present invention also envisages the explicit exclusion of one or more of any of the group members in the claimed invention.

## BRIEF DESCRIPTION OF THE FIGURES

[041] Figure 1. RSPO expression in tumors and normal tissues. Shown is a summary of microarray data from normal, benign, and malignant tissue human samples. Individual tick marks indicate the expression level of RSPO mRNA. (A) RSPO1 (B) RSPO2 (C) RSPO3

[042] Figure 2. Binding studies of RSPO proteins and LGR5. FACS analysis of HEK-293 cells expressing LGR5. HEK-293 cells were transiently transfected with a cDNA expression vector encoding FLAG-LGR5-CD4TM-GFP and then subsequently mixed with soluble RSPO1-Fc, RSPO2-Fc, RSPO3-Fc, or RSPO4-Fc fusion proteins. An anti-FLAG antibody was used as a positive control, and soluble FZD8-Fc was used as a negative control. Specific binding is indicated by the presence of signal within the dark lined box overlay on each FACS plot.

[043] Figure 3. Anti-RSPO3 antibodies inhibit  $\beta$ -catenin signaling induced by RSPO3 and WNT3A. A TOPflash luciferase reporter assay was used to measure  $\beta$ -catenin signaling in HEK-293 cells after exposure to a combination of WNT3a (5ng/ml) and RSPO3 (10ng/ml) and in the presence of increasing concentrations of anti-RSPO3 antibodies (131R002 or 131R003). Antibodies were used as 4-fold serial dilutions from 20 $\mu$ g/ml to 0.02 $\mu$ g/ml. Controls included exposure to control medium (no WNT3a and no RSPO), WNT3a alone, or a combination of WNT3a and RSPO3 in the absence of antibody.

[044] Figure 4. Affinity-matured 131R003 antibody variants inhibit  $\beta$ -catenin signaling induced by RSPO3 and WNT3A. A TOPflash luciferase reporter assay was used to measure  $\beta$ -catenin signaling in HEK-293 cells after exposure to a combination of WNT3a and RSPO3 and in the presence of increasing concentrations of anti-RSPO3 antibodies (131R003 (-▲-), 131R003 CDR1 variant (-■-), or 131R003 CDR3 variant (-●-)). Antibodies were used as 5-fold serial dilutions from 20 $\mu$ g/ml to 0.006 $\mu$ g/ml. Controls included exposure to control medium (no WNT3a and no RSPO)/cells only (-Δ-), a control antibody (-▼-), WNT3a alone (-◆-), or a combination of WNT3a and RSPO3 in the absence of antibody (-□-).

[045] Figure 5. Inhibition of tumor growth with anti-RSPO antibodies. OV38 ovarian tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with a combination of anti-RSPO1 antibody 89M5 and anti-RSPO3 antibody 131R003 (-●-), taxol (-▲-), a combination of anti-RSPO1 antibody 89M5, anti-RSPO3 antibody 131R003, and taxol (-▼-), or a control antibody (-■-). Data is shown as tumor volume (mm<sup>3</sup>) over days post-treatment.

[046] Figure 6. Inhibition of tumor growth with anti-RSPO antibodies. OV38 ovarian tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with a combination of anti-RSPO1 antibody 89M5 and anti-RSPO3 antibody 131R002 (-▲-), a combination of anti-RSPO1 antibody 89M5 and taxol (-○-), a combination of anti-RSPO3 antibody 131R002 and taxol (-□-), a combination of anti-

RSPO1 antibody 89M5, anti-RSPO3 antibody 131R002, and taxol ( $\Delta$ ), taxol alone ( $\nabla$ ), or a control antibody ( $\blacksquare$ ). Data is shown as tumor volume ( $\text{mm}^3$ ) over days post-treatment.

[047] Figure 7. Inhibition of tumor growth with anti-RSPO3 antibodies. (A) LU45 lung tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R002 ( $\circ$ ) or a control antibody ( $\blacksquare$ ). (B) LU25 lung tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R002 ( $\circ$ ) or a control antibody ( $\blacksquare$ ). Data is shown as tumor volume ( $\text{mm}^3$ ) over days post-treatment.

[048] Figure 8. Affinity-matured antibody variants inhibit  $\beta$ -catenin signaling induced by RSPO3 and WNT3A. A TOPflash luciferase reporter assay was used to measure  $\beta$ -catenin signaling in HEK-293T cells after exposure to a combination of WNT3a and human RSPO3 and in the presence of increasing concentrations of anti-RSPO3 antibody 131R002 ( $\blacktriangle$ ), 131R006 ( $\bullet$ ), or 131R007 ( $\blacksquare$ ). Antibodies were used as 5-fold serial dilutions from  $20\mu\text{g/ml}$  to  $0.0064\mu\text{g/ml}$ . Controls included exposure to control medium (no WNT3a and no RSPO/cells ( $\circ$ )), WNT3a alone ( $\nabla$ ), or a combination of WNT3a and human RSPO3 in the absence of antibody ( $\blacklozenge$ ).

[049] Figure 9. Inhibition of RSPO3 and LGR5 interaction by anti-RSPO3 antibodies. FACS analysis of HEK-293T cells expressing LGR5. HEK-293T cells were transiently transfected with a cDNA expression vector encoding the extracellular domain of human LGR5 (FLAG-LGR5-CD4TM-GFP) and then subsequently mixed with RSPO3-biotin fusion protein in combination with anti-RSPO3 antibodies 131R006 or 131R007. Binding was detected with PE-conjugated streptavidin. Relative RSPO3-biotin binding is shown on the y-axis and expression of the FLAG-LGR5-CD4TM-GFP fusion protein is indicated on the x-axis. Positive binding is indicated by the presence of signal within the dark lined box overlay on each FACS plot.

[050] Figure 10. Inhibition of tumor growth with anti-RSPO antibodies. NCI-H2030 cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R002 ( $\otimes$ ), carboplatin alone ( $\Delta$ ), a combination of anti-RSPO3 antibody 131R002 and carboplatin ( $\circ$ ), or a control antibody ( $\blacksquare$ ). Data is shown as tumor volume ( $\text{mm}^3$ ) over days post-treatment.

[051] Figure 11. Inhibition of tumor growth with anti-RSPO antibodies. LU102 lung tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R002 ( $\bullet$ ), carboplatin alone ( $\Delta$ ), a combination of anti-RSPO3 antibody 131R002 and carboplatin ( $\circ$ ), or a control antibody ( $\blacksquare$ ). (A) Data is shown as tumor volume ( $\text{mm}^3$ ) over days post-treatment. (B) Gene set enrichment analysis results.

[052] Figure 12. Inhibition of tumor growth with anti-RSPO antibodies. PN35 pancreatic tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R002 ( $\bullet$ ), a combination of gemcitabine and nab-paclitaxel (ABRAXANE) ( $\Delta$ ), a combination of anti-RSPO3 antibody 131R002 and gemcitabine and nab-paclitaxel (ABRAXANE) ( $\circ$ ), or a control

antibody (-■-). Data is shown as tumor volume (mm<sup>3</sup>) over days post-treatment. (A) All four treatment groups; (B) the gemcitabine and nab-paclitaxel treatment group and the anti-RSPO3 antibody gemcitabine and nab-paclitaxel treatment on an expanded scale.

[053] Figure 13. Inhibition of  $\beta$ -catenin signaling induced by RSPO3 and WNT3A. A TOPflash luciferase reporter assay was used to measure  $\beta$ -catenin signaling in HEK-293T cells after exposure to a combination of WNT3a and human RSPO3 and in the presence of increasing concentrations of anti-RSPO3 antibody 131R007 (-□-) or 131R010 (-●-). Antibodies were used as 5-fold serial dilutions from 20 $\mu$ g/ml to 0.0064 $\mu$ g/ml. Controls included exposure to control medium (no WNT3a and no RSPO/cells (-▲-)), WNT3a alone (-▼-), or a combination of WNT3a and human RSPO3 in the absence of antibody (-◆-).

[054] Figure 14. Inhibition of tumor growth with anti-RSPO antibodies. LU25 NSCLC lung tumor cells were injected subcutaneously into NOD/SCID mice. Mice were treated with anti-RSPO3 antibody 131R008 (-▲-), paclitaxel alone (-○-), a combination of anti-RSPO3 antibody 131R008 and paclitaxel (-●-), or a control antibody (-■-). Data is shown as tumor volume (mm<sup>3</sup>) over days post-treatment.

#### DETAILED DESCRIPTION OF THE INVENTION

[055] The present invention provides novel agents, including, but not limited to polypeptides such as antibodies, that bind RSPO proteins, particularly human RSPO3. The RSPO3-binding agents include, but are not limited to, antagonists of  $\beta$ -catenin signaling. The RSPO3-binding agents include, but are not limited to, inhibitors of RSPO3 and LGR protein interactions. Related polypeptides and polynucleotides, compositions comprising the RSPO3-binding agents, and methods of making the RSPO3-binding agents are also provided. Methods of using the novel RSPO3-binding agents, such as methods of inhibiting tumor growth, methods of treating cancer, methods of modulating angiogenesis, methods of reducing the frequency of cancer stem cells in a tumor, methods of inhibiting  $\beta$ -catenin signaling, and/or methods of identifying and/or selecting subjects for treatment, are further provided.

[056] Monoclonal antibodies that specifically bind human RSPO3 have been identified - monoclonal antibodies 131R002 and 131R003 (Example 3). Anti-RSPO3 antibodies 131R002 and 131R003 have binding affinities for human RSPO3 of less than 10 nM (Example 3). Anti-RSPO3 antibodies 131R002 and 131R003 inhibit RSPO3-induced  $\beta$ -catenin signaling (Example 4, Fig. 3). Affinity-matured variants of 131R003 inhibit RSPO3-induced  $\beta$ -catenin signaling and have greater activity than parental 131R003 (Example 5, Fig. 4). Anti-RSPO3 antibodies inhibit tumor growth as single agents, in combination with anti-RSPO1 antibodies, and in combination with one or more chemotherapeutic agents (Examples 6, 7, 11, 12 and 14; Figs. 5-7, 10-12 and 14). Humanized anti-RSPO3 antibodies h131R006 and h131R007 are stronger inhibitors of  $\beta$ -catenin activity than antibody 131R002 (Example 8, Fig. 8). Anti-RSPO3

antibodies h131R006 and h131R007 block binding of RSPO3 to LGR5 (Example 9, Fig. 9). Humanized anti-RSPO3 antibody h131R010 isotype IgG1 inhibits  $\beta$ -catenin activity similar to the IgG2 isotype antibody h131R007 (Example 13, Fig. 13).

#### I. Definitions

[057] To facilitate an understanding of the present invention, a number of terms and phrases are defined below.

[058] The terms “antagonist” and “antagonistic” as used herein refer to any molecule that partially or fully blocks, inhibits, reduces, or neutralizes a biological activity of a target and/or signaling pathway (e.g., the  $\beta$ -catenin signaling). The term “antagonist” is used herein to include any molecule that partially or fully blocks, inhibits, reduces, or neutralizes the activity of a protein (e.g., a RSPO protein). Suitable antagonist molecules specifically include, but are not limited to, antagonist antibodies or antibody fragments.

[059] The terms “modulation” and “modulate” as used herein refer to a change or an alteration in a biological activity. Modulation includes, but is not limited to, stimulating or inhibiting an activity. Modulation may be an increase or a decrease in activity (e.g., a decrease in RSPO signaling; a decrease in  $\beta$ -catenin signaling), a change in binding characteristics, or any other change in the biological, functional, or immunological properties associated with the activity of a protein, pathway, or other biological point of interest.

[060] The term “antibody” as used herein refers to an immunoglobulin molecule that recognizes and specifically binds a target, such as a protein, polypeptide, peptide, carbohydrate, polynucleotide, lipid, or combinations of the foregoing, through at least one antigen-binding site within the variable region(s) of the immunoglobulin molecule. As used herein, the term encompasses intact polyclonal antibodies, intact monoclonal antibodies, single chain antibodies, antibody fragments (such as Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments), single chain Fv (scFv) antibodies, multispecific antibodies such as bispecific antibodies, monospecific antibodies, monovalent antibodies, chimeric antibodies, humanized antibodies, human antibodies, fusion proteins comprising an antigen-binding site of an antibody, and any other modified immunoglobulin molecule comprising an antigen recognition site (i.e., antigen-binding site) as long as the antibodies exhibit the desired biological activity. An antibody can be any of the five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, or subclasses (isotypes) thereof (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2), based on the identity of their heavy chain constant domains referred to as alpha, delta, epsilon, gamma, and mu, respectively. The different classes of immunoglobulins have different and well-known subunit structures and three-dimensional configurations. Antibodies can be naked or conjugated to other molecules, including but not limited to, toxins and radioisotopes.

[061] The term “antibody fragment” refers to a portion of an intact antibody and refers to the antigenic determining variable regions of an intact antibody. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments, linear antibodies, single chain antibodies, and multispecific antibodies formed from antibody fragments. “Antibody fragment” as used herein comprises an antigen-binding site or epitope-binding site.

[062] The term “variable region” of an antibody refers to the variable region of an antibody light chain, or the variable region of an antibody heavy chain, either alone or in combination. The variable regions of the heavy and light chains each consist of four framework regions (FR) connected by three complementarity determining regions (CDRs), also known as “hypervariable regions”. The CDRs in each chain are held together in close proximity by the framework regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of the antibody. There are at least two techniques for determining CDRs: (1) an approach based on cross-species sequence variability (i.e., Kabat et al., 1991, *Sequences of Proteins of Immunological Interest, 5th Edition*, National Institutes of Health, Bethesda, MD), and (2) an approach based on crystallographic studies of antigen-antibody complexes (Al-Lazikani et al., 1997, *J. Mol. Biol.*, 273:927-948). In addition, combinations of these two approaches are sometimes used in the art to determine CDRs.

[063] The term “monoclonal antibody” as used herein refers to a homogeneous antibody population involved in the highly specific recognition and binding of a single antigenic determinant or epitope. This is in contrast to polyclonal antibodies that typically include a mixture of different antibodies directed against a variety of different antigenic determinants. The term “monoclonal antibody” encompasses both intact and full-length monoclonal antibodies as well as antibody fragments (e.g., Fab, Fab', F(ab')<sub>2</sub>, Fv), single chain (scFv) antibodies, bispecific antibodies, fusion proteins comprising an antibody portion, and any other modified immunoglobulin molecule comprising an antigen recognition site (antigen-binding site). Furthermore, “monoclonal antibody” refers to such antibodies made by any number of techniques, including but not limited to, hybridoma production, phage selection, recombinant expression, and transgenic animals.

[064] The term “humanized antibody” as used herein refers to forms of non-human (e.g., murine) antibodies that are specific immunoglobulin chains, chimeric immunoglobulins, or fragments thereof that contain minimal non-human sequences. Typically, humanized antibodies are human immunoglobulins in which residues of the CDRs are replaced by residues from the CDRs of a non-human species (e.g., mouse, rat, rabbit, or hamster) that have the desired specificity, affinity, and/or binding capability (Jones et al., 1986, *Nature*, 321:522-525; Riechmann et al., 1988, *Nature*, 332:323-327; Verhoeyen et al., 1988, *Science*, 239:1534-1536). In some instances, the Fv framework region residues of a human immunoglobulin are replaced with the corresponding residues in an antibody from a non-human species that has the desired specificity, affinity, structural, and/or binding capability. The humanized antibody

can be further modified by the substitution of additional residues either in the Fv framework region and/or within the replaced non-human residues to refine and optimize antibody specificity, affinity, structural, and/or binding capability. In general, the humanized antibody will comprise substantially all of at least one, and typically two or three of the CDRs that correspond to the non-human immunoglobulin whereas all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody can also comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Examples of methods used to generate humanized antibodies are described in, for example, U.S. Pat. 5,225,539.

[065] The term “human antibody” as used herein refers to an antibody produced by a human or an antibody having an amino acid sequence corresponding to an antibody produced by a human. A human antibody may be made using any of the techniques known in the art. This definition of a human antibody specifically excludes a humanized antibody comprising non-human CDRs.

[066] The term “chimeric antibody” as used herein refers to an antibody wherein the amino acid sequence of the immunoglobulin molecule is derived from two or more species. Typically, the variable region of both light and heavy chains corresponds to the variable region of antibodies derived from one species of mammal (e.g., mouse, rat, rabbit, etc.) with the desired specificity, affinity, and/or binding capability, while the constant regions correspond to sequences in antibodies derived from another species (usually human).

[067] The phrase “affinity-matured antibody” as used herein refers to an antibody with one or more alterations in one or more CDRs thereof that result in an improvement in the affinity of the antibody for an antigen, compared to a parent antibody that does not possess those alterations(s). The definition also includes alterations in non-CDR residues made in conjunction with alterations to CDR residues. Preferred affinity-matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity-matured antibodies are produced by procedures known in the art. For example, Marks et al., 1992, *Bio/Technology* 10:779-783, describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by Barbas et al., 1994, *PNAS*, 91:3809-3813; Schier et al., 1995, *Gene*, 169:147-155; Yelton et al., 1995, *J. Immunol.* 155:1994-2004; Jackson et al., 1995, *J. Immunol.*, 154:3310-9; and Hawkins et al., 1992, *J. Mol. Biol.*, 226:889-896. Site-directed mutagenesis may also be used to obtain affinity-matured antibodies.

[068] The terms “epitope” and “antigenic determinant” are used interchangeably herein and refer to that portion of an antigen capable of being recognized and specifically bound by a particular antibody. When the antigen is a polypeptide, epitopes can be formed both from contiguous amino acids and noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids (also referred to as linear epitopes) are typically retained upon protein denaturing, whereas epitopes formed by tertiary folding (also referred to as conformational epitopes) are typically lost upon protein

denaturing. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation.

**[069]** The terms “heteromultimeric molecule” or “heteromultimer” or “heteromultimeric complex” or “heteromultimeric polypeptide” are used interchangeably herein to refer to a molecule comprising at least a first polypeptide and a second polypeptide, wherein the second polypeptide differs in amino acid sequence from the first polypeptide by at least one amino acid residue. The heteromultimeric molecule can comprise a “heterodimer” formed by the first and second polypeptide or can form higher order tertiary structures where additional polypeptides are present.

**[070]** The terms “selectively binds” or “specifically binds” mean that a binding agent or an antibody reacts or associates more frequently, more rapidly, with greater duration, with greater affinity, or with some combination of the above to the epitope, protein or target molecule than with alternative substances, including unrelated proteins. In certain embodiments “specifically binds” means, for instance, that an antibody binds a protein with a  $K_D$  of about 0.1mM or less, but more usually less than about 1 $\mu$ M. In certain embodiments, “specifically binds” means that an antibody binds a target at times with a  $K_D$  of at least about 0.1 $\mu$ M or less, at other times at least about 0.01 $\mu$ M or less, and at other times at least about 1nM or less. Because of the sequence identity between homologous proteins in different species, specific binding can include an antibody that recognizes a protein in more than one species (e.g., human RSPO3 and mouse RSPO3). Likewise, because of homology within certain regions of polypeptide sequences of different proteins, specific binding can include an antibody (or other polypeptide or binding agent) that recognizes more than one protein (e.g., human RSPO3 and human RSPO1). It is understood that, in certain embodiments, an antibody or binding moiety that specifically binds a first target may or may not specifically bind a second target. As such, “specific binding” does not necessarily require (although it can include) exclusive binding, i.e. binding to a single target. Thus, an antibody may, in certain embodiments, specifically bind more than one target. In certain embodiments, multiple targets may be bound by the same antigen-binding site on the antibody. For example, an antibody may, in certain instances, comprise two identical antigen-binding sites, each of which specifically binds the same epitope on two or more proteins (e.g., RSPO3 and RSPO1). In certain alternative embodiments, an antibody may be multispecific and comprise at least two antigen-binding sites with differing specificities. By way of non-limiting example, a bispecific antibody may comprise one antigen-binding site that recognizes an epitope on one protein (e.g., human RSPO3) and further comprise a second, different antigen-binding site that recognizes a different epitope on a second protein (e.g., human RSPO2). Generally, but not necessarily, reference to binding means specific binding.

**[071]** The terms “polypeptide” and “peptide” and “protein” are used interchangeably herein and refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino



acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids), as well as other modifications known in the art. It is understood that, because the polypeptides of this invention may be based upon antibodies, in certain embodiments, the polypeptides can occur as single chains or associated chains.

[072] The terms “polynucleotide” and “nucleic acid” are used interchangeably herein and refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase.

[073] “Conditions of high stringency” may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 15mM NaCl/1.5mM sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 in 5x SSC (0.75M NaCl, 75mM sodium citrate) at 42°C; or (3) employ during hybridization 50% formamide in 5x SSC, 50mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5x Denhardt's solution, sonicated salmon sperm DNA (50µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2x SSC and 50% formamide, followed by a high-stringency wash consisting of 0.1x SSC containing EDTA at 55°C.

[074] The terms “identical” or percent “identity” in the context of two or more nucleic acids or polypeptides, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned (introducing gaps, if necessary) for maximum correspondence, not considering any conservative amino acid substitutions as part of the sequence identity. The percent identity may be measured using sequence comparison software or algorithms or by visual inspection. Various algorithms and software that may be used to obtain alignments of amino acid or nucleotide sequences are well-known in the art. These include, but are not limited to, BLAST, ALIGN, Megalign, BestFit, GCG Wisconsin Package, and variations thereof. In some embodiments, two nucleic acids or polypeptides of the invention are substantially identical, meaning they have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, and in some embodiments at least 95%, 96%, 97%, 98%, 99% nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection. In some embodiments, identity exists over a region of the sequences that is at least about 10, at least about 20, at least about 40-60 residues, at least about 60-80 residues in length or any integral value therebetween. In some embodiments, identity exists over a longer

region than 60-80 residues, such as at least about 80-100 residues, and in some embodiments the sequences are substantially identical over the full length of the sequences being compared, such as the coding region of a nucleotide sequence.

[075] A “conservative amino acid substitution” is one in which one amino acid residue is replaced with another amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). For example, substitution of a phenylalanine for a tyrosine is a conservative substitution. Preferably, conservative substitutions in the sequences of the polypeptides and antibodies of the invention do not abrogate the binding of the polypeptide or antibody containing the amino acid sequence, to the antigen(s), i.e., the one or more RSPO protein(s) to which the polypeptide or antibody binds. Methods of identifying nucleotide and amino acid conservative substitutions which do not eliminate antigen binding are well-known in the art.

[076] The term “vector” as used herein means a construct, which is capable of delivering, and usually expressing, one or more gene(s) or sequence(s) of interest in a host cell. Examples of vectors include, but are not limited to, viral vectors, naked DNA or RNA expression vectors, plasmid, cosmid, or phage vectors, DNA or RNA expression vectors associated with cationic condensing agents, and DNA or RNA expression vectors encapsulated in liposomes.

[077] A polypeptide, antibody, polynucleotide, vector, cell, or composition which is “isolated” is a polypeptide, antibody, polynucleotide, vector, cell, or composition which is in a form not found in nature. Isolated polypeptides, antibodies, polynucleotides, vectors, cells, or compositions include those which have been purified to a degree that they are no longer in a form in which they are found in nature. In some embodiments, a polypeptide, antibody, polynucleotide, vector, cell, or composition which is isolated is substantially pure.

[078] The term “substantially pure” as used herein refers to material which is at least 50% pure (i.e., free from contaminants), at least 90% pure, at least 95% pure, at least 98% pure, or at least 99% pure.

[079] The terms “cancer” and “cancerous” as used herein refer to or describe the physiological condition in mammals in which a population of cells are characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, blastoma, sarcoma, and hematologic cancers such as lymphoma and leukemia.

[080] The terms “tumor” and “neoplasm” as used herein refer to any mass of tissue that results from excessive cell growth or proliferation, either benign (noncancerous) or malignant (cancerous) including pre-cancerous lesions.

[081] The term “metastasis” as used herein refers to the process by which a cancer spreads or transfers from the site of origin to other regions of the body with the development of a similar cancerous lesion at a new location. A “metastatic” or “metastasizing” cell is one that loses adhesive contacts with neighboring cells and migrates via the bloodstream or lymph from the primary site of disease to invade neighboring body structures.

[082] The terms “cancer stem cell” and “CSC” and “tumor stem cell” and “tumor initiating cell” are used interchangeably herein and refer to cells from a cancer or tumor that: (1) have extensive proliferative capacity; 2) are capable of asymmetric cell division to generate one or more types of differentiated cell progeny wherein the differentiated cells have reduced proliferative or developmental potential; and (3) are capable of symmetric cell divisions for self-renewal or self-maintenance. These properties confer on the cancer stem cells the ability to form or establish a tumor or cancer upon serial transplantation into an immunocompromised host (e.g., a mouse) compared to the majority of tumor cells that fail to form tumors. Cancer stem cells undergo self-renewal versus differentiation in a chaotic manner to form tumors with abnormal cell types that can change over time as mutations occur.

[083] The terms “cancer cell” and “tumor cell” refer to the total population of cells derived from a cancer or tumor or pre-cancerous lesion, including both non-tumorigenic cells, which comprise the bulk of the cancer cell population, and tumorigenic stem cells (cancer stem cells). As used herein, the terms “cancer cell” or “tumor cell” will be modified by the term “non-tumorigenic” when referring solely to those cells lacking the capacity to renew and differentiate to distinguish those tumor cells from cancer stem cells.

[084] The term “tumorigenic” as used herein refers to the functional features of a cancer stem cell including the properties of self-renewal (giving rise to additional tumorigenic cancer stem cells) and proliferation to generate all other tumor cells (giving rise to differentiated and thus non-tumorigenic tumor cells).

[085] The term “tumorigenicity” as used herein refers to the ability of a random sample of cells from the tumor to form palpable tumors upon serial transplantation into immunocompromised hosts (e.g., mice). This definition also includes enriched and/or isolated populations of cancer stem cells that form palpable tumors upon serial transplantation into immunocompromised hosts (e.g., mice).

[086] The term “subject” refers to any animal (e.g., a mammal), including, but not limited to, humans, non-human primates, canines, felines, rodents, and the like, which is to be the recipient of a particular treatment. Typically, the terms “subject” and “patient” are used interchangeably herein in reference to a human subject.

[087] The term “pharmaceutically acceptable” refers to a product or compound approved (or approvable) by a regulatory agency of the Federal government or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, including humans.

[088] The terms “pharmaceutically acceptable excipient, carrier or adjuvant” or “acceptable pharmaceutical carrier” refer to an excipient, carrier or adjuvant that can be administered to a subject, together with at least one binding agent (e.g., an antibody) of the present disclosure, and which does not destroy the activity of the binding agent. The excipient, carrier, or adjuvant should be non-toxic when administered with a binding agent in doses sufficient to deliver a therapeutic effect.

[089] The terms “effective amount” or “therapeutically effective amount” or “therapeutic effect” refer to an amount of a binding agent, an antibody, polypeptide, polynucleotide, small organic molecule, or other drug effective to “treat” a disease or disorder in a subject or mammal. In the case of cancer, the therapeutically effective amount of a drug (e.g., an antibody) has a therapeutic effect and as such can reduce the number of cancer cells; decrease tumorigenicity, tumorigenic frequency or tumorigenic capacity; reduce the number or frequency of cancer stem cells; reduce the tumor size; reduce the cancer cell population; inhibit and/or stop cancer cell infiltration into peripheral organs including, for example, the spread of cancer into soft tissue and bone; inhibit and/or stop tumor or cancer cell metastasis; inhibit and/or stop tumor or cancer cell growth; relieve to some extent one or more of the symptoms associated with the cancer; reduce morbidity and mortality; improve quality of life; or a combination of such effects. To the extent the agent, for example an antibody, prevents growth and/or kills existing cancer cells, it can be referred to as cytostatic and/or cytotoxic.

[090] The terms “treating” or “treatment” or “to treat” or “alleviating” or “to alleviate” refer to both 1) therapeutic measures that cure, slow down, lessen symptoms of, and/or halt progression of a diagnosed pathologic condition or disorder and 2) prophylactic or preventative measures that prevent or slow the development of a targeted pathologic condition or disorder. Thus those in need of treatment include those already with the disorder; those prone to have the disorder; and those in whom the disorder is to be prevented. In some embodiments, a subject is successfully “treated” according to the methods of the present invention if the patient shows one or more of the following: a reduction in the number of or complete absence of cancer cells; a reduction in the tumor size; inhibition of or an absence of cancer cell infiltration into peripheral organs including the spread of cancer cells into soft tissue and bone; inhibition of or an absence of tumor or cancer cell metastasis; inhibition or an absence of cancer growth; relief of one or more symptoms associated with the specific cancer; reduced morbidity and mortality; improvement in quality of life; reduction in tumorigenicity; reduction in the number or frequency of cancer stem cells; or some combination of effects.

[091] As used in the present disclosure and claims, the singular forms “a”, “an” and “the” include plural forms unless the context clearly dictates otherwise.

[092] It is understood that wherever embodiments are described herein with the language “comprising” otherwise analogous embodiments described in terms of “consisting of” and/or “consisting essentially of” are also provided. It is also understood that wherever embodiments are described herein with the language “consisting essentially of” otherwise analogous embodiments described in terms of “consisting of” are also provided.

[093] The term “and/or” as used in a phrase such as “A and/or B” herein is intended to include both A and B; A or B; A (alone); and B (alone). Likewise, the term “and/or” as used in a phrase such as “A, B, and/or C” is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

## II. RSPO-binding agents

[094] The present invention provides agents that specifically bind human RSPO proteins. These agents are referred to herein as “RSPO-binding agents”. In some embodiments, the RSPO-binding agent is an antibody. In some embodiments, the RSPO-binding agent is a polypeptide. In certain embodiments, the RSPO-binding agent binds RSPO3 (“RSPO3-binding agents”). In certain embodiments, the RSPO3-binding agent specifically binds at least one other human RSPO. In some embodiments, the at least one other human RSPO bound by a RSPO3-binding agent is selected from the group consisting of RSPO1, RSPO2, and RSPO4. In some embodiments, the RSPO3-binding agent is an antibody that binds a common epitope on RSPO1, RSPO2, and/or RSPO4. In some embodiments, the RSPO3-binding agent is a bispecific antibody that binds a first epitope on RSPO3 and binds a second, different epitope on RSPO1, RSPO2, and/or RSPO4. The full-length amino acid (aa) sequences for human RSPO1, RSPO2, RSPO3, and RSPO4 are known in the art and are provided herein as SEQ ID NO:1 (RSPO1), SEQ ID NO:2 (RSPO2), SEQ ID NO:3 (RSPO3), and SEQ ID NO:4 (RSPO4).

[095] In certain embodiments, the antigen-binding site of a RSPO-binding agent (e.g., an antibody or a bispecific antibody) described herein is capable of binding (or binds) one, two, three, or four RSPOs. In certain embodiments, the antigen-binding site of a RSPO-binding agent (e.g., an antibody or a bispecific antibody) described herein is capable of binding (or binds) RSPO3 as well as one, two, or three other RSPOs. For example, in certain embodiments, the antigen-binding site of a RSPO3-binding agent is capable of specifically binding RSPO3 as well as at least one other RSPO selected from the group consisting of RSPO1, RSPO2, and RSPO4. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3 and RSPO1. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3 and RSPO2. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3 and RSPO4. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3, RSPO1, and RSPO2. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3, RSPO1, and RSPO4. In certain embodiments, the RSPO3-binding agent specifically binds RSPO3, RSPO2, and

RSPO4. In some embodiments, the RSPO3-binding agent specifically binds human RSPO3. In some embodiments, the RSPO3-binding agent (e.g., antibody) specifically binds both human RSPO3 and mouse RSPO3.

[096] In certain embodiments, the agent-binding agent is an antibody that specifically binds within amino acids 22-272 of human RSPO3. In certain embodiments, the agent-binding agent is an antibody that specifically binds within amino acids 22-207 of human RSPO3. In certain embodiments, the antigen-binding agent is an antibody that specifically binds within amino acids 35-135 of human RSPO3. In certain embodiments, the antigen-binding agent is an antibody that specifically binds within amino acids 35-86 of human RSPO3. In certain embodiments, the antigen-binding agent is an antibody that specifically binds within amino acids 92-135 of human RSPO3. In certain embodiments, the RSPO3-binding agent binds within SEQ ID NO:5. In certain embodiments, the RSPO3-binding agent or antibody binds a furin-like cysteine-rich domain of RSPO3. In some embodiments, the agent or antibody binds at least one amino acid within a furin-like cysteine-rich domain of RSPO3. In certain embodiments, the RSPO3-binding agent or antibody binds within sequence SEQ ID NO:6 or SEQ ID NO:7. In certain embodiments, the RSPO3-binding agent or antibody binds within sequence SEQ ID NO:6 and SEQ ID NO:7. In some embodiments, the RSPO3-binding agent binds the thrombospondin domain of RSPO3. In some embodiments, the RSPO3-binding agent or antibody binds at least one amino acid within the thrombospondin domain of RSPO3. In some embodiments, the RSPO3-binding agent or antibody binds within SEQ ID NO:8.

[097] In certain embodiments, the RSPO-binding agent or antibody binds at least one RSPO protein with a dissociation constant ( $K_D$ ) of about 1  $\mu$ M or less, about 100nM or less, about 40nM or less, about 20nM or less, about 10nM or less, about 1nM or less, or about 0.1nM or less. In certain embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a dissociation constant ( $K_D$ ) of about 1  $\mu$ M or less, about 100nM or less, about 40nM or less, about 20nM or less, about 10nM or less, about 1nM or less, or about 0.1nM or less. In some embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a  $K_D$  of about 20nM or less. In some embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a  $K_D$  of about 10nM or less. In some embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a  $K_D$  of about 1nM or less. In some embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a  $K_D$  of about 0.5nM or less. In some embodiments, a RSPO3-binding agent or antibody binds RSPO3 with a  $K_D$  of about 0.1nM or less. In certain embodiments, a RSPO3-binding agent or antibody described herein binds at least one other RSPO. In certain embodiments, a RSPO3-binding agent or antibody described herein that binds at least one other RSPO, binds at least one other RSPO with a  $K_D$  of about 100nM or less, about 20nM or less, about 10nM or less, about 1nM or less or about 0.1nM or less. For example, in some embodiments, a RSPO3-binding agent or antibody also binds RSPO1, RSPO2, and/or RSPO4 with a  $K_D$  of about 10nM or less. In some embodiments, the RSPO-binding agent binds both

human RSPO and mouse RSPO with a  $K_D$  of about 10nM or less. In some embodiments, a RSPO3-binding agent binds both human RSPO3 and mouse RSPO3 with a  $K_D$  of about 1nM or less. In some embodiments, a RSPO3-binding agent binds both human RSPO3 and mouse RSPO3 with a  $K_D$  of about 0.1nM or less. In some embodiments, the dissociation constant of the binding agent (e.g., an antibody) to a RSPO3 protein is the dissociation constant determined using a RSPO3 fusion protein comprising at least a portion of the RSPO3 protein immobilized on a Biacore chip. In some embodiments, the dissociation constant of the binding agent (e.g., an antibody) to a RSPO3 protein is the dissociation constant determined using the binding agent captured by an anti-human IgG antibody on a Biacore chip and a RSPO3 protein.

**[098]** In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first antigen-binding site that specifically binds RSPO3 and a second antigen-binding site that specifically binds a second target. In some embodiments, a RSPO3-binding agent or antibody binds both RSPO3 and the second target with a  $K_D$  of about 100nM or less. In some embodiments, a RSPO3-binding agent or antibody binds both RSPO3 and the second target with a  $K_D$  of about 50nM or less. In some embodiments, a RSPO3-binding agent or antibody binds both RSPO3 and the second target with a  $K_D$  of about 20nM or less. In some embodiments, a RSPO3-binding agent or antibody binds both RSPO3 and the second target with a  $K_D$  of about 10nM or less. In some embodiments, a RSPO3-binding agent or antibody binds both RSPO3 and the second target with a  $K_D$  of about 1nM or less. In some embodiments, the affinity of one of the antigen-binding sites may be weaker than the affinity of the other antigen-binding site. For example, the  $K_D$  of one antigen binding site may be about 1nM and the  $K_D$  of the second antigen-binding site may be about 10nM. In some embodiments, the difference in affinity between the two antigen-binding sites may be about 2-fold or more, about 3-fold or more, about 5-fold or more, about 8-fold or more, about 10-fold or more, about 15-fold or more, about 20-fold or more, about 30-fold or more, about 50-fold or more, or about 100-fold or more. Modulation of the affinities of the two antigen-binding sites may affect the biological activity of the bispecific antibody. For example, decreasing the affinity of the antigen-binding site for RSPO3 or the second target, may have a desirable effect, for example decreased toxicity of the binding agent and/or increased therapeutic index.

**[099]** By way of non-limiting example, the bispecific antibody may comprise (a) a first antigen-binding site that binds human RSPO3 with a  $K_D$  between about 0.1nM and about 10nM, and (b) a second antigen-binding site that specifically binds a second target (e.g., human RSPO2) with a  $K_D$  between about 0.1nM and about 20nM, between about 0.5nM and about 20nM, or between about 1.0nM and 10nM.

**[0100]** In certain embodiments, the RSPO-binding agent (e.g., an antibody) binds to at least one human RSPO protein with a half maximal effective concentration ( $EC_{50}$ ) of about 1 $\mu$ M or less, about 100nM or less, about 40nM or less, about 20nM or less, about 10nM or less, about 1nM or less, or about 0.1nM or less. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds to human RSPO3 with a

half maximal effective concentration ( $EC_{50}$ ) of about  $1\mu\text{M}$  or less, about  $100\text{nM}$  or less, about  $40\text{nM}$  or less, about  $20\text{nM}$  or less, about  $10\text{nM}$  or less, about  $1\text{nM}$  or less, or about  $0.1\text{nM}$  or less. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) also binds to human RSPO1, RSPO2, and/or RSPO4 with an  $EC_{50}$  of about  $40\text{nM}$  or less, about  $20\text{nM}$  or less, about  $10\text{nM}$  or less, about  $1\text{nM}$  or less or about  $0.1\text{nM}$  or less.

**[0101]** In certain embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the antibody is a recombinant antibody. In some embodiments, the antibody is a monoclonal antibody. In some embodiments, the antibody is a chimeric antibody. In some embodiments, the antibody is a humanized antibody. In some embodiments, the antibody is a human antibody. In some embodiments, the antibody is an IgA, IgD, IgE, IgG, or IgM antibody. In certain embodiments, the antibody is an IgG1 antibody. In certain embodiments, the antibody is an IgG2 antibody. In certain embodiments, the antibody is an antibody fragment comprising an antigen-binding site. In some embodiments, the antibody is a bispecific antibody or a multispecific antibody. In some embodiments, the antibody is a monovalent antibody. In some embodiments, the antibody is a monospecific antibody. In some embodiments, the antibody is a bivalent antibody. In some embodiments, the antibody is conjugated to a cytotoxic moiety. In some embodiments, the antibody is isolated. In some embodiments, the antibody is substantially pure.

**[0102]** The RSPO3-binding agents (e.g., antibodies) of the present invention can be assayed for specific binding by any method known in the art. The immunoassays which can be used include, but are not limited to, competitive and non-competitive assay systems using techniques such as Biacore analysis, FACS analysis, immunofluorescence, immunocytochemistry, Western blot analysis, radioimmunoassays, ELISA, "sandwich" immunoassays, immunoprecipitation assays, precipitation reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, and protein A immunoassays. Such assays are routine and well-known in the art (see, e.g., Ausubel et al., Editors, 1994-present, *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., New York, NY).

**[0103]** For example, the specific binding of an antibody to human RSPO3 may be determined using ELISA. An ELISA assay comprises preparing antigen, coating wells of a 96 well microtiter plate with antigen, adding the RSPO3-binding antibody or other RSPO3-binding agent conjugated to a detectable compound such as an enzymatic substrate (e.g. horseradish peroxidase or alkaline phosphatase) to the well, incubating for a period of time and detecting the presence of the antibody bound to the antigen. In some embodiments, the RSPO3-binding antibody or agent is not conjugated to a detectable compound, but instead a second conjugated antibody that recognizes the RSPO3-binding agent or antibody (e.g., an anti-Fc antibody) and is conjugated to a detectable compound is added to the well. In some embodiments, instead of coating the well with the antigen, the RSPO3-binding agent or antibody can be coated to the well and a second antibody conjugated to a detectable compound can be added following the addition of



the antigen to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art.

[0104] In another example, the specific binding of an antibody to human RSPO3 may be determined using FACS. A FACS screening assay may comprise generating a cDNA construct that expresses an antigen as a fusion protein (e.g., RSPO3-Fc or RSPO3-CD4<sup>TM</sup>), transfecting the construct into cells, expressing the antigen on the surface of the cells, mixing the RSPO3-binding agent with the transfected cells, and incubating for a period of time. The cells bound by the RSPO3-binding agent may be identified using a secondary antibody conjugated to a detectable compound (e.g., PE-conjugated anti-Fc antibody) and a flow cytometer. One of skill in the art would be knowledgeable as to the parameters that can be modified to optimize the signal detected as well as other variations of FACS that may enhance screening (e.g., screening for blocking antibodies).

[0105] The binding affinity of an antibody or other binding-agent to an antigen (e.g., RSPO3) and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., <sup>3</sup>H or <sup>125</sup>I), or fragment or variant thereof, with the antibody of interest in the presence of increasing amounts of unlabeled antigen followed by the detection of the antibody bound to the labeled antigen. The affinity of the antibody for the antigen and the binding off-rates can be determined from the data by Scatchard plot analysis. In some embodiments, Biacore kinetic analysis is used to determine the binding on and off rates of antibodies or agents that bind an antigen (e.g., RSPO3). In some embodiments, Biacore kinetic analysis comprises analyzing the binding and dissociation of antibodies from chips with immobilized antigen (e.g., RSPO3) on their surface. In some embodiments, Biacore kinetic analysis comprises analyzing the binding and dissociation of antigen (e.g., RSPO3) from chips with immobilized antibody (e.g., anti-RSPO3 antibody) on their surface.

[0106] In certain embodiments, the invention provides a RSPO3-binding agent (e.g., an antibody) that specifically binds human RSPO3, wherein the RSPO3-binding agent (e.g., an antibody) comprises one, two, three, four, five, and/or six of the CDRs of antibody 131R002, antibody 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011 (see Table 1). In some embodiments, the RSPO3-binding agent comprises one or more of the CDRs of 131R002, 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011; two or more of the CDRs of 131R002, 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011; three or more of the CDRs of 131R002, 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011; four or more of the CDRs of 131R002, 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011; five or more of the

CDRs of 131R002, 131R003, or the humanized variants thereof, including 131R005/131R007, h131R006A, h131R006B, or h131R008, h131R010, or h131R011; or all six of the CDRs of 131R002, 131R003, or the humanized variants thereof, including h131R005/131R007, h131R006A, h131R006B, h131R008, h131R010, or h131R011.

Table 1

	131R002/131R003 and Humanized Variants
HC CDR1	KASGYTFTDYS (SEQ ID NO:9) or KASGYTFTSYTF (SEQ ID NO:34) or DYSIH (SEQ ID NO:78)
HC CDR2	IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79)
HC CDR3	ATYFANYFDY (SEQ ID NO:11) or ATYFANNFDY (SEQ ID NO:35) or TYFANNFD (SEQ ID NO:80)
LC CDR1	QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81)
LC CDR2	AAS (SEQ ID NO: 13) or AASNLES (SEQ ID NO:82)
LC CDR3	QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83)

[0107] In certain embodiments, the invention provides a RSPO3-binding agent (e.g., an antibody) that specifically binds human RSPO3, wherein the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments, the RSPO3-binding agent further comprises a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83). In some embodiments, the RSPO3-binding agent comprises a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), and (b) a light chain CDR1 comprising

QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and (b) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), and (b) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and (b) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and (b) a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLTF (SEQ ID NO:83). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and (b) a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and (b) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14).

**[0108]** In certain embodiments, the invention provides a RSPO3-binding agent (e.g., an antibody or bispecific antibody) that specifically binds human RSPO3, wherein the RSPO3-binding agent comprises: (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), DYSIH (SEQ ID NO:78), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions;

(b) a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), YIYPSNGDSGYNQKFK (SEQ ID NO:79), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (c) a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), TYFANNFD (SEQ ID NO:80), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (d) a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), KASQSVDDYDGDSYMN (SEQ ID NO:81), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; (e) a light chain CDR2 comprising AAS (SEQ ID NO:13), AASNLES (SEQ ID NO:82), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions; and (f) a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14), QQSNEPLTF (SEQ ID NO:83), or a variant thereof comprising 1, 2, 3, or 4 amino acid substitutions. In certain embodiments, the amino acid substitutions are conservative substitutions. In some embodiments, the substitutions are made as part of a germline humanization process.

**[0109]** In certain embodiments, the invention provides a RSPO3-binding agent (e.g., an antibody) that specifically binds RSPO3, wherein the RSPO3-binding agent comprises a heavy chain variable region having at least about 80% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62 and/or a light chain variable region having at least 80% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:15. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:16. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:36. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:37. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:44. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:45. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:62. In certain embodiments, the RSPO3-binding agent comprises a light chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a light chain

variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:72. In certain embodiments, the RSPO3-binding agent comprises a light chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region having at least about 95% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62 and/or a light chain variable region having at least about 95% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region comprising SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and/or a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region comprising SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62 and a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting essentially of SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and a light chain variable region consisting essentially of SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting of SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and a light chain variable region consisting of SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.

**[0110]** In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region comprising SEQ ID NO:44 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region comprising SEQ ID NO:45 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region comprising SEQ ID NO:62 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting essentially of SEQ ID NO:44 and a light chain variable region consisting essentially of SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting essentially of SEQ ID NO:45 and a light chain variable region consisting essentially of SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting essentially of SEQ ID NO:62 and a light chain variable region consisting essentially of SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting of SEQ ID NO:44 and a light chain variable region consisting of SEQ ID NO:17. In certain embodiments, the RSPO3-binding agent comprises a heavy chain variable



certain embodiments, the RSPO3-binding agent comprises a heavy chain variable region consisting of SEQ ID NO:62 and a light chain variable region consisting of SEQ ID NO:86.

**[0113]** In certain embodiments, the invention provides a RSPO3-binding agent (e.g., an antibody) that specifically binds RSPO3, wherein the RSPO3-binding agent comprises: (a) a heavy chain having at least 90% sequence identity to SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and/or (b) a light chain having at least 90% sequence identity to SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises: (a) a heavy chain having at least 95% sequence identity to SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and/or (b) a light chain having at least 95% sequence identity to SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:27 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:28 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:39 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:42 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and/or a light chain comprising SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and a light chain consisting essentially of SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:28, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69 and a light chain consisting of SEQ ID NO:29. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:28, SEQ ID

NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69 and a light chain consisting of SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:27 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:28 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:39 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:42 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and/or a light chain comprising SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and/or a light chain comprising SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and a light chain consisting essentially of SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69 and a light chain consisting of SEQ ID NO:74. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and a light chain consisting essentially of SEQ ID NO:88. In some embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69 and a light chain consisting of SEQ ID NO:88.

**[0114]** In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and a light chain comprising SEQ ID NO:29. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and a light chain variable region comprising SEQ ID NO:29. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and a light chain variable region comprising SEQ ID NO:29. In certain embodiments, the RSPO3-



**[0115]** In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:48 and a light chain comprising SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:49 and a light chain variable region comprising SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and a light chain variable region comprising SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and a light chain variable region comprising SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:48 and a light chain consisting essentially of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:49 and a light chain consisting essentially of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:64 and a light chain consisting essentially of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:69 and a light chain consisting essentially of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:48 and a light chain consisting of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:49 and a light chain consisting of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:64 and a light chain consisting of SEQ ID NO:74. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:69 and a light chain consisting of SEQ ID NO:74.

36

comprises a heavy chain comprising SEQ ID NO:49 and a light chain variable region comprising SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:64 and a light chain variable region comprising SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain comprising SEQ ID NO:69 and a light chain variable region comprising SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:48 and a light chain consisting essentially of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:49 and a light chain consisting essentially of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:64 and a light chain consisting essentially of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting essentially of SEQ ID NO:69 and a light chain consisting essentially of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:48 and a light chain consisting of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:49 and a light chain consisting of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:64 and a light chain consisting of SEQ ID NO:88. In certain embodiments, the RSPO3-binding agent comprises a heavy chain consisting of SEQ ID NO:69 and a light chain consisting of SEQ ID NO:88.

[0117] In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and light chain variable region of the 131R002 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain and light chain of the 131R002 antibody (with or without the leader sequence). In certain embodiments, a RSPO3-binding agent is the 131R002 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R002 antibody in a chimeric form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R002 antibody in a humanized form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain CDRs and/or light chain CDRs of the 131R002 antibody in a humanized form of the antibody. In some embodiments, the humanized version of 131R002 is an IgG1 antibody. In some embodiments, the humanized version of 131R002 is an IgG2 antibody.

[0118] In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody 131R002.

[0119] In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and light chain variable region of the 131R003 antibody. In some embodiments, the RSPO3-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable region from 131R003 has been affinity-matured. In some embodiments, the RSPO3-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable

region comprises at least one modified or altered CDR as compared to the parent 131R003 antibody. In some embodiments, the RSPO-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable region comprises a modified CDR1 as compared to the parent 131R003 antibody. In some embodiments, the RSPO-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable region comprises a modified CDR2 as compared to the parent 131R003 antibody. In some embodiments, the RSPO-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable region comprises a modified CDR3 as compared to the parent 131R003 antibody. In some embodiments, the RSPO-binding agent comprises the heavy chain variable region of the 131R003 antibody wherein the heavy chain variable region comprises a modified CDR1 and CDR3 as compared to the parent 131R003 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain and light chain of the 131R003 antibody (with or without the leader sequence). In certain embodiments, a RSPO3-binding agent is the 131R003 antibody. In certain embodiments, a RSPO3-binding agent is a variant of the 131R003 antibody that comprises a different heavy chain CDR1 as compared to the parent 131R003 antibody. In certain embodiments, a RSPO3-binding agent is a variant of the 131R003 antibody that comprises a different heavy chain CDR3 as compared to the parent 131R003 antibody. In certain embodiments, a RSPO3-binding agent is a variant of the 131R003 antibody that comprises a different heavy chain CDR1 and a different heavy chain CDR3 as compared to the parent 131R003 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R003 antibody or of any of the variants of 131R003 in a chimeric form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R003 antibody or of any of the variants of 131R003 in a humanized form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain CDRs and/or light chain CDRs of the 131R003 antibody or of any of the variants of 131R003 in a humanized form of the antibody. In some embodiments, the humanized version of 131R003 or of 131R003 variants is an IgG1 antibody. In some embodiments, the humanized version of 131R003 or of 131R003 variants is an IgG2 antibody.

**[0120]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody 131R003. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody 131R003.

**[0121]** In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and light chain variable region of the 131R006B antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain and light chain of the 131R006B antibody (with or without the leader sequence). In certain embodiments, a RSPO3-binding agent is the 131R006B antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain

variable region of the 131R006B antibody in a chimeric form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R006B antibody in a humanized form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain CDRs and/or light chain CDRs of the 131R006B antibody in a humanized form of the antibody. In some embodiments, the humanized version of 131R006B is an IgG1 antibody. In some embodiments, the humanized version of 131R006B is an IgG2 antibody.

**[0122]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody 131R006B. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody 131R006B.

**[0123]** In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and light chain variable region of the 131R005/131R007 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain and light chain of the 131R005/131R007 antibody (with or without the leader sequence). In certain embodiments, a RSPO3-binding agent is the 131R005/131R007 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R005/131R007 antibody in a chimeric form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the 131R005/131R007 antibody in a humanized form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain CDRs and/or light chain CDRs of the 131R005/131R007 antibody in a humanized form of the antibody. In some embodiments, the humanized version of 131R005/131R007 is an IgG1 antibody. In some embodiments, the humanized version of 131R005/131R007 is an IgG2 antibody. In some embodiments, the anti-RSPO3 antibody is 131R008.

**[0124]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody 131R005/131R007. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody 131R005/131R007.

**[0125]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody 131R008. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody 131R008.

**[0126]** In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and light chain variable region of the h131R010 or h131R011 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain and light chain of the h131R010 or 131R011 antibody (with or without the leader sequence). In certain embodiments, a RSPO3-binding agent is the h131R010 antibody. In certain embodiments, a RSPO3-binding agent is the h131R011 antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain variable region and/or light chain variable region of the h131R010 or h131R011 antibody in a chimeric form of the antibody. In certain embodiments, a RSPO3-binding agent comprises the heavy chain CDRs and/or light chain CDRs of the h131R010 or h131R011

antibody. In some embodiments, the anti-RSPO3 antibody is h131R010. In some embodiments, the anti-RSPO3 antibody is h131R011.

**[0127]** In some embodiments, the RSPO3-binding agent comprises a heavy chain variable region encoded by the plasmid deposited with American Type Culture Collection (ATCC), and designated PTA-\_\_\_\_\_. In some embodiments, the RSPO3-binding agent comprises a light chain variable region encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_. In some embodiments, the RSPO3-binding agent comprises a heavy chain variable region encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_, and a light chain variable region encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_. In some embodiments, the RSPO3-binding agent comprises a heavy chain encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_. In some embodiments, the RSPO3-binding agent comprises a light chain encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_. In some embodiments, the RSPO3-binding agent comprises a heavy chain encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_, and a light chain encoded by the plasmid deposited with ATCC and designated PTA-\_\_\_\_\_.

**[0128]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody h131R010. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody h131R010.

**[0129]** In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, the antibody h131R011. In certain embodiments, a RSPO3-binding agent comprises, consists essentially of, or consists of, a variant of the antibody h131R011.

**[0130]** In certain embodiments, the invention provides a RSPO3-binding agent that is a bispecific antibody. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first antigen-binding site that specifically binds human RSPO3. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first antigen-binding site that specifically binds human RSPO3 and a second antigen-binding site that binds a second target. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), OR TYFANNFD (SEQ ID NO:80). In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35). In some embodiments, the RSPO3-binding agent is a

bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35). In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11). In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising DYSIH (SEQ ID NO:80), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80). In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising DYSIH (SEQ ID NO:80) or KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80). In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising: a first antigen-binding site that specifically binds human RSPO3, wherein the first antigen-binding site comprises (a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35) or TYFANNFD (SEQ ID NO:80), and a second antigen-binding site, wherein the first antigen-binding site and the second antigen-binding site comprise a common (i.e., identical) light chain. In some embodiments, the bispecific antibody comprises a first antigen-binding site comprising a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEEDPLT (SEQ ID NO:14) or QQSNEEDPLTF (SEQ ID NO:83).

**[0131]** In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region having at least about 80% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62. In certain embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62. In some embodiments, the bispecific antibody comprises a light chain variable region at least about 85%, at least about 90%, at least about 95%, at least about 97%,

or at least about 99% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:44. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:45. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:62. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first light chain variable region comprising SEQ ID NO:17. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first light chain variable region comprising SEQ ID NO:72. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first light chain variable region comprising SEQ ID NO:86.

**[0132]** In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62 and a first heavy chain constant region comprising SEQ ID NO:60 or SEQ ID NO:61. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:44 and a first heavy chain constant region comprising SEQ ID NO:60 or SEQ ID NO:61. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:45 and a first heavy chain constant region comprising SEQ ID NO:60 or SEQ ID NO:61. In some embodiments, the RSPO3-binding agent is a bispecific antibody comprising a first heavy chain variable region comprising SEQ ID NO:62 and a first heavy chain constant region comprising SEQ ID NO:60 or SEQ ID NO:61.

**[0133]** In certain embodiments, the RSPO3-binding agent is a bispecific antibody that specifically binds human RSPO3 and a second target. In some embodiments, the RSPO3-binding agent is a bispecific antibody that specifically binds human RSPO3 and a second human RSPO. In some embodiments, the RSPO3-binding agent is a bispecific antibody that specifically binds human RSPO3 and a second human RSPO selected from the group consisting of RSPO1, RSPO2, and RSPO4. Non-limiting examples of antibodies to human RSPO have been described in, for example, International Patent Pub. No. WO 2013/012747.

**[0134]** In some embodiments, the RSPO3-binding agent is a bispecific antibody that specifically binds human RSPO3 and human RSPO1. In some embodiments, the bispecific antibody comprises: a) a first antigen-binding site that specifically binds human RSPO3, and b) a second antigen-binding site that specifically binds human RSPO1, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11),

ATYFANNFDY (SEQ ID NO:35) or TYFANNFD (SEQ ID NO:80); and wherein both the first and second antigen-binding sites comprise a common light chain.

**[0135]** In some embodiments, the RSPO3-binding agent is a bispecific antibody that specifically binds human RSPO3 and human RSPO2. In some embodiments, the bispecific antibody comprises: a) a first antigen-binding site that specifically binds human RSPO3, and b) a second antigen-binding site that specifically binds human RSPO2, wherein the first antigen-binding site comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35) or TYFANNFD (SEQ ID NO:80); and wherein both the first and second antigen-binding sites comprise a common light chain.

**[0136]** In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises a heavy chain variable region from the anti-RSPO3 antibody 131R003. In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a heavy chain variable region from a variant of the anti-RSPO3 antibody 131R003. In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises a heavy chain variable region from the anti-RSPO3 antibody 131R006B. In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises a heavy chain variable region from the anti-RSPO3 antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises a heavy chain variable region from the anti-RSPO3 antibody h131R010 or h131R011.

**[0137]** In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises a first CH3 domain and a second CH3 domain, each of which is modified to promote formation of heteromultimers. In some embodiments, the first and second CH3 domains are modified using a knobs-into-holes technique. In some embodiments, the first and second CH3 domains comprise changes in amino acids that result in altered electrostatic interactions. In some embodiments, the first and second CH3 domains comprise changes in amino acids that result in altered hydrophobic/hydrophilic interactions.

**[0138]** In some embodiments, the RSPO3-binding agent is a bispecific antibody that comprises heavy chain constant regions selected from the group consisting of: (a) a first human IgG1 constant region, wherein the amino acids corresponding to positions 253 and 292 of IgG1 (SEQ ID NO:56) are replaced with glutamate or aspartate, and a second human IgG1 constant region, wherein the amino acids corresponding to positions 240 and 282 of IgG1 (SEQ ID NO:56) are replaced with lysine; (b) a first human IgG2 constant region, wherein the amino acids corresponding to positions 249 and 288 of IgG2 (SEQ ID NO:57) are replaced with glutamate or aspartate, and a second human IgG2 constant region wherein the amino acids corresponding to positions 236 and 278 of IgG2 (SEQ ID NO:57) are replaced with lysine; (c) a first human IgG3 constant region, wherein the amino acids corresponding to positions



300 and 339 of IgG3 (SEQ ID NO:58) are replaced with glutamate or aspartate, and a second human IgG3 constant region wherein the amino acids corresponding to positions 287 and 329 of IgG3 (SEQ ID NO:58) are replaced with lysine; and (d) a first human IgG4 constant region, wherein the amino acids corresponding to positions 250 and 289 of IgG4 (SEQ ID NO:59) are replaced with glutamate or aspartate, and a second IgG4 constant region wherein the amino acids corresponding to positions 237 and 279 of IgG4 (SEQ ID NO:59) are replaced with lysine.

**[0139]** In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG1 constant region with amino acid substitutions at positions corresponding to positions 253 and 292 of IgG1 (SEQ ID NO:56), wherein the amino acids at positions corresponding to positions 253 and 292 of IgG1 (SEQ ID NO:56) are replaced with glutamate or aspartate, and a second human IgG1 constant region with amino acid substitutions at positions corresponding to positions 240 and 282 of IgG1 (SEQ ID NO:56), wherein the amino acids at positions corresponding to positions 240 and 282 of IgG1 (SEQ ID NO:56) are replaced with lysine. In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG2 constant region with amino acid substitutions at positions corresponding to positions 249 and 288 of IgG2 (SEQ ID NO:57), wherein the amino acids at positions corresponding to positions 249 and 288 of IgG2 (SEQ ID NO:57) are replaced with glutamate or aspartate, and a second human IgG2 constant region with amino acid substitutions at positions corresponding to positions 236 and 278 of IgG2 (SEQ ID NO:57), wherein the amino acids at positions corresponding to positions 236 and 278 of IgG2 (SEQ ID NO:57) are replaced with lysine. In some embodiments, the RSPO-binding agent is a bispecific antibody which comprises a first human IgG3 constant region with amino acid substitutions at positions corresponding to positions 300 and 339 of IgG3 (SEQ ID NO:58), wherein the amino acids at positions corresponding to positions 300 and 339 of IgG3 (SEQ ID NO:58) are replaced with glutamate or aspartate, and a second human IgG3 constant region with amino acid substitutions at positions corresponding to positions 287 and 329 of IgG3 (SEQ ID NO:58), wherein the amino acids at positions corresponding to positions 287 and 329 of IgG3 (SEQ ID NO:58) are replaced with lysine. In some embodiments, the RSPO-binding agent is a bispecific antibody which comprises a first human IgG4 constant region with amino acid substitutions at positions corresponding to positions 250 and 289 of IgG4 (SEQ ID NO:59), wherein the amino acids at positions corresponding to positions 250 and 289 of IgG4 (SEQ ID NO:59) are replaced with glutamate or aspartate, and a second human IgG4 constant region with amino acid substitutions at positions corresponding to positions 237 and 279 of IgG4 (SEQ ID NO:59), wherein the amino acids at positions corresponding to positions 237 and 279 of IgG4 (SEQ ID NO:59) are replaced with lysine.

**[0140]** In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG1 constant region with amino acid substitutions at positions corresponding to positions 253 and 292 of IgG1 (SEQ ID NO:56), wherein the amino acids are replaced with glutamate, and a second human

IgG1 constant region with amino acid substitutions at positions corresponding to positions 240 and 282 of IgG1 (SEQ ID NO:56), wherein the amino acids are replaced with lysine. In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG1 constant region with amino acid substitutions at positions corresponding to positions 253 and 292 of IgG1 (SEQ ID NO:56), wherein the amino acids are replaced with aspartate, and a second human IgG1 constant region with amino acid substitutions at positions corresponding to positions 240 and 282 of IgG1 (SEQ ID NO:56), wherein the amino acids are replaced with lysine.

[0141] In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG2 constant region with amino acid substitutions at positions corresponding to positions 249 and 288 of IgG2 (SEQ ID NO:57), wherein the amino acids are replaced with glutamate, and a second human IgG2 constant region with amino acid substitutions at positions corresponding to positions 236 and 278 of IgG2 (SEQ ID NO:57), wherein the amino acids are replaced with lysine. In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first human IgG2 constant region with amino acid substitutions at positions corresponding to positions 249 and 288 of IgG2 (SEQ ID NO:57), wherein the amino acids are replaced with aspartate, and a second human IgG2 constant region with amino acid substitutions at positions corresponding to positions 236 and 278 of IgG2 (SEQ ID NO:57), wherein the amino acids are replaced with lysine.

[0142] In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a heavy chain constant region of SEQ ID NO:60. In some embodiments, the RSPO-binding agent is a bispecific antibody which comprises a heavy chain constant region of SEQ ID NO:61. In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises a first heavy chain constant region of SEQ ID NO:60 and a second heavy chain constant region of SEQ ID NO:61.

[0143] In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds RSPO3 with a  $K_D$  of about 50nM or less, about 25nM or less, about 10nM or less, about 1nM or less, or about 0.1nM or less. In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds a second target (e.g., RSPO2) with a  $K_D$  of about 50nM or less, about 25nM or less, about 10nM or less, about 1nM or less, or about 0.1nM or less. In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds RSPO3 with a  $K_D$  of about 50nM or less and binds a second target (e.g., RSPO2) with a  $K_D$  of about 50nM or less. In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds RSPO3 with a  $K_D$  of about 25nM or less and binds a second target (e.g., RSPO2) with a  $K_D$  of about 25nM or less. In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds RSPO3 with a  $K_D$  of about 10nM or less and binds a second target (e.g., RSPO2) with a  $K_D$  of about 10nM or less. In some embodiments, the RSPO3-binding agent is a bispecific antibody which binds RSPO3 with a  $K_D$  of about 1nM or less and binds a second target (e.g., RSPO2) with a  $K_D$  of about 1nM or less.

[0144] In some embodiments, the RSPO3-binding agent is a bispecific antibody which comprises one antigen-binding site with a binding affinity that is weaker than the binding affinity of the second antigen-binding site. For example, in some embodiments, the bispecific antibody may bind RSPO3 with a  $K_D$  ranging from about 0.1nM to 1nM and may bind a second target (e.g., RSPO2) with a  $K_D$  ranging from about 1nM to 10nM. Or the bispecific antibody may bind RSPO3 with a  $K_D$  ranging from about 1nM to 10nM and may bind a second target (e.g., RSPO2) with a  $K_D$  ranging from about 0.1nM to 1nM. In some embodiments, the bispecific antibody may bind RSPO3 with a  $K_D$  ranging from about 0.1nM to 1nM and may bind a second target (e.g., RSPO2) with a  $K_D$  ranging from about 1nM to 10nM. Or the bispecific antibody may bind RSPO3 with a  $K_D$  ranging from about 1nM to 10nM and may bind a second target (e.g., RSPO2) with a  $K_D$  ranging from about 0.1nM to 1nM. In some embodiments, the difference in affinity between the two antigen-binding sites may be about 2-fold or more, about 3-fold or more, about 5-fold or more, about 8-fold or more, about 10-fold or more, about 15-fold or more, about 30-fold or more, about 50-fold or more, or about 100-fold or more. In some embodiments, at least one amino acid residue in at least one CDR of the antigen-binding site for RSPO3 is substituted with a different amino acid so that the affinity of the RSPO3-binding site is altered. In some embodiments, the affinity of the RSPO3-binding site is increased. In some embodiments, the affinity of the RSPO3-binding site is decreased. In some embodiments, at least one amino acid residue in at least one CDR of the antigen-binding site for the second target (e.g., RSPO2) is substituted with a different amino acid so that the affinity of the second antigen-binding site is altered. In some embodiments, the affinity of the second antigen-binding site is increased. In some embodiments, the affinity of the second antigen-binding site is decreased. In some embodiments, the affinities of both the RSPO3 and the second antigen-binding sites are altered.

[0145] The invention provides polypeptides, including, but not limited to, antibodies that specifically bind human RSPO proteins. In some embodiments, the polypeptides bind human RSPO3. In some embodiments, the polypeptides bind human RSPO3 and at least one additional human RSPO selected from the group consisting of RSPO1, RSPO2, and RSPO4.

[0146] In certain embodiments, the polypeptide comprises one, two, three, four, five, and/or six of the CDRs of antibody 131R002, 131R003, or variants of 131R003 including h131R005/131R007, h131R006A, h131R006B, h131R010, and h131R011 (see Table 1 herein). In some embodiments, the polypeptide comprises CDRs with up to four (i.e., 0, 1, 2, 3, or 4) amino acid substitutions per CDR. In certain embodiments, the heavy chain CDR(s) are contained within a heavy chain variable region. In certain embodiments, the light chain CDR(s) are contained within a light chain variable region.

[0147] In some embodiments, the invention provides a polypeptide that specifically binds human RSPO3, wherein the polypeptide comprises an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID

NO:45, or SEQ ID NO:62, and/or an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and/or an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:15 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:16 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:36 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:37 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:44 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:45 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:62 and/or an amino acid sequence comprising SEQ ID NO:17. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:15 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:16 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:36 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:37 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:44 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:45 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:62 and/or an amino acid sequence comprising SEQ ID NO:72. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:44 and/or an amino acid sequence comprising SEQ ID NO:86. In certain embodiments, the

polypeptide comprises an amino acid sequence comprising SEQ ID NO:45 and/or an amino acid sequence comprising SEQ ID NO:86. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:62 and/or an amino acid sequence comprising SEQ ID NO:86.

**[0148]** In some embodiments, the invention provides a polypeptide that specifically binds human RSPO3, wherein the polypeptide comprises an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:63, or SEQ ID NO:68, and/or an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:23, SEQ ID NO:73, or SEQ ID NO:87. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:63, or SEQ ID NO:68. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:23, SEQ ID NO:73, or SEQ ID NO:87. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:63, or SEQ ID NO:68, and/or an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:23, SEQ ID NO:73, or SEQ ID NO:87. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:63, or SEQ ID NO:68, and/or an amino acid sequence comprising SEQ ID NO:23, SEQ ID NO:73, or SEQ ID NO:87. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:63, or SEQ ID NO:68, and/or SEQ ID NO:23, SEQ ID NO:73, or SEQ ID NO:87.

**[0149]** In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69, and/or an amino acid sequence comprising SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:48 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:49 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:64 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:69 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:48 and/or an amino acid sequence comprising SEQ ID NO:74. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID

[illegible]

the polypeptide comprises an amino acid sequence consisting essentially of SEQ ID NO:69 and/or an amino acid sequence consisting essentially of SEQ ID NO:88.

**[0150]** In some embodiments, the invention provides a polypeptide that specifically binds human RSPO3, wherein the polypeptide comprises an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69, and/or an amino acid sequence having at least about 80% sequence identity to SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% sequence identity to SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69, and/or an amino acid sequence having at least about 95% sequence identity to SEQ ID NO:29, SEQ ID NO:74, SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69, and/or an amino acid sequence comprising SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:48 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:49 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:64 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:69 and/or an amino acid sequence comprising SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:27 and/or SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:28 and/or SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:48 and/or SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:49 and/or SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:64 and/or SEQ ID NO:29. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:69 and/or SEQ ID NO:29. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:48 and/or an amino acid sequence comprising SEQ ID NO:74. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:49 and/or an amino acid sequence comprising SEQ ID NO:74. In certain

embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:64 and/or an amino acid sequence comprising SEQ ID NO:74. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:69 and/or an amino acid sequence comprising SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:27 and/or SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:28 and/or SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:48 and/or SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:49 and/or SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:64 and/or SEQ ID NO:74. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:69 and/or SEQ ID NO:74. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:48 and/or an amino acid sequence comprising SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:49 and/or an amino acid sequence comprising SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:64 and/or an amino acid sequence comprising SEQ ID NO:88. In certain embodiments, the polypeptide comprises an amino acid sequence comprising SEQ ID NO:69 and/or an amino acid sequence comprising SEQ ID NO:88. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:48 and/or SEQ ID NO:88. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:49 and/or SEQ ID NO:88. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:64 and/or SEQ ID NO:88. In certain embodiments, the polypeptide consists essentially of SEQ ID NO:69 and/or SEQ ID NO:88.

**[0151]** In some embodiments, a RSPO3-binding agent comprises a polypeptide comprising a sequence selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:86, SEQ ID NO:87, and SEQ ID NO:88.

**[0152]** Many proteins, including antibodies, contain a signal sequence that directs the transport of the proteins to various locations. Signal sequences (also referred to as signal peptides or leader sequences) are located at the N-terminus of nascent polypeptides. They target the polypeptide to the endoplasmic reticulum and the proteins are sorted to their destinations, for example, to the inner space of an organelle, to an interior membrane, to the cell's outer membrane, or to the cell exterior via secretion. Most signal sequences are cleaved from the protein by a signal peptidase after the proteins are transported to the endoplasmic reticulum. The cleavage of the signal sequence from the polypeptide usually occurs at a specific site in the amino acid sequence and is dependent upon amino acid residues within the signal



sequence. Although there is usually one specific cleavage site, more than one cleavage site may be recognized and/or may be used by a signal peptidase resulting in a non-homogenous N-terminus of the polypeptide. For example, the use of different cleavage sites within a signal sequence can result in a polypeptide expressed with different N-terminal amino acids. Accordingly, in some embodiments, the polypeptides as described herein may comprise a mixture of polypeptides with different N-termini. In some embodiments, the N-termini differ in length by 1, 2, 3, 4, or 5 amino acids. In some embodiments, the polypeptide is substantially homogeneous, i.e., the polypeptides have the same N-terminus. In some embodiments, the signal sequence of the polypeptide comprises one or more (e.g., one, two, three, four, five, six, seven, eight, nine, ten, etc.) amino acid substitutions and/or deletions as compared to a “native” or “parental” signal sequence. In some embodiments, the signal sequence of the polypeptide comprises amino acid substitutions and/or deletions that allow one cleavage site to be dominant, thereby resulting in a substantially homogeneous polypeptide with one N-terminus. In some embodiments, a signal sequence of the polypeptide affects the expression level of the polypeptide, e.g., increased expression or decreased expression.

**[0153]** In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62, and a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:44 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:45 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:62 and a light chain variable region comprising SEQ ID NO:17. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:44 and a light chain variable region comprising SEQ ID NO:72. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:45 and a light chain variable region comprising SEQ ID NO:72. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:62 and a light chain variable region comprising SEQ ID NO:72. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:44 and a light chain

variable region comprising SEQ ID NO:86. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:45 and a light chain variable region comprising SEQ ID NO:86. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain variable region comprising SEQ ID NO:62 and a light chain variable region comprising SEQ ID NO:86.

[0154] In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69 and a light chain comprising SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:48 and a light chain comprising SEQ ID NO:29. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:49 and a light chain comprising SEQ ID NO:29. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:64 and a light chain comprising SEQ ID NO:29. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:69 and a light chain comprising SEQ ID NO:29. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:48 and a light chain comprising SEQ ID NO:74. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:49 and a light chain comprising SEQ ID NO:74. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:64 and a light chain comprising SEQ ID NO:74. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:69 and a light chain comprising SEQ ID NO:74. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:48 and a light chain comprising SEQ ID NO:88. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:49 and a light chain comprising SEQ ID NO:88. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3 with an antibody that comprises a heavy chain comprising SEQ ID NO:64 and a light chain comprising SEQ ID NO:88. In certain embodiments, a RSPO3-binding agent (e.g., antibody) competes for specific binding to RSPO3

with an antibody that comprises a heavy chain comprising SEQ ID NO:69 and a light chain comprising SEQ ID NO:88.

**[0155]** In certain embodiments, a RSPO3-binding agent competes with antibody 131R002 or antibody 131R003 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with a variant of antibody 131R003 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with a humanized version of antibody 131R003 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with antibody h131R005/131R007 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with antibody h131R008 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with antibody h131R006A or antibody h131R006B for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with antibody h131R010 for specific binding to human RSPO3. In certain embodiments, a RSPO3-binding agent competes with antibody h131R011 for specific binding to human RSPO3. In some embodiments, a RSPO3-binding agent or antibody competes for specific binding to RSPO3 in an *in vitro* competitive binding assay. In some embodiments, the RSPO3 is human RSPO3. In some embodiments, the RSPO3 is mouse RSPO3.

**[0156]** In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as an antibody of the invention. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody 131R002 or antibody 131R003. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as a variant of antibody 131R003. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as a humanized version of antibody 131R003. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R008. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R010. In certain embodiments, a RSPO3-binding agent (e.g., an antibody) binds the same epitope, or essentially the same epitope, on RSPO3 as antibody h131R011.

**[0157]** In another embodiment, a RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by an antibody of the invention. In some embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody 131R002 or antibody 131R003. In another embodiment, the RSPO3-binding

agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by a humanized version of antibody 131R003. In certain embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R006A or antibody h131R006B. In certain embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R005/131R007. In certain embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R008. In certain embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R010. In certain embodiments, the RSPO3-binding agent is an antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by antibody h131R011.

**[0158]** In certain embodiments, the RSPO-binding agent (e.g., an antibody) described herein binds at least one human RSPO protein and modulates RSPO activity. In some embodiments, the RSPO-binding agent is a RSPO antagonist and decreases RSPO activity. In some embodiments, the RSPO-binding agent is a RSPO antagonist and decreases  $\beta$ -catenin activity.

**[0159]** In certain embodiments, a RSPO3-binding agent (e.g., an antibody) described herein binds human RSPO3 and modulates RSPO3 activity. In some embodiments, a RSPO3-binding agent is a RSPO3 antagonist and decreases RSPO3 activity. In some embodiments, a RSPO3-binding agent is a RSPO3 antagonist and decreases  $\beta$ -catenin activity.

**[0160]** In certain embodiments, the RSPO-binding agent (e.g., an antibody) is an antagonist of at least one human RSPO protein. In some embodiments, the RSPO-binding agent is an antagonist of at least one RSPO and inhibits RSPO activity. In certain embodiments, the RSPO-binding agent inhibits RSPO activity by at least about 10%, at least about 20%, at least about 30%, at least about 50%, at least about 75%, at least about 90%, or about 100%. In some embodiments, the RSPO-binding agent inhibits activity of one, two, three, or four RSPO proteins. In some embodiments, the RSPO-binding agent inhibits activity of human RSPO1, RSPO2, RSPO3, and/or RSPO4. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent inhibits RSPO3 activity. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is antibody h131R008. In certain embodiments, a

RSPO3-binding agent that inhibits human RSPO3 activity is antibody h131R010. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 activity is antibody h131R011.

[0161] In certain embodiments, the RSPO-binding agent (e.g., antibody) is an antagonist of at least one human RSPO protein. In certain embodiments, the RSPO-binding agent inhibits RSPO signaling by at least about 10%, at least about 20%, at least about 30%, at least about 50%, at least about 75%, at least about 90%, or about 100%. In some embodiments, the RSPO-binding agent inhibits signaling by one, two, three, or four RSPO proteins. In some embodiments, the RSPO-binding agent inhibits signaling of human RSPO1, RSPO2, RSPO3, and/or RSPO4. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent inhibits human RSPO3 signaling. In certain embodiments, a RSPO3-binding agent that inhibits RSPO3 signaling is antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits RSPO3 signaling is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 signaling is antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 signaling is antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 signaling is antibody h131R008. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 signaling is antibody h131R010. In certain embodiments, a RSPO3-binding agent that inhibits human RSPO3 signaling is antibody h131R011.

[0162] In certain embodiments, the RSPO-binding agent (e.g., antibody) is an antagonist of  $\beta$ -catenin signaling. In certain embodiments, the RSPO-binding agent inhibits  $\beta$ -catenin signaling by at least about 10%, at least about 20%, at least about 30%, at least about 50%, at least about 75%, at least about 90%, or about 100%. In some embodiments, the RSPO-binding agent that inhibits  $\beta$ -catenin signaling is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent inhibits  $\beta$ -catenin signaling. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R008. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R010. In certain embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R011.

[0163] In certain embodiments, the RSPO-binding agent (e.g., antibody) inhibits binding of at least one RSPO protein to a receptor. In certain embodiments, the RSPO-binding agent inhibits binding of a human

RSPO protein to one or more of its receptors. In some embodiments, the RSPO-binding agent inhibits binding of a RSPO protein to at least one LGR protein. In some embodiments, the RSPO-binding agent inhibits binding of a RSPO protein to LGR4, LGR5, and/or LGR6. In some embodiments, a RSPO3-binding agent inhibits binding of RSPO3 to LGR4. In some embodiments, a RSPO3-binding agent inhibits binding of RSPO3 to LGR5. In some embodiments, a RSPO3-binding agent inhibits binding of RSPO3 to LGR6. In certain embodiments, the inhibition of binding of a RSPO-binding agent to at least one LGR protein is at least about 10%, at least about 25%, at least about 50%, at least about 75%, at least about 90%, or at least about 95%. In certain embodiments, a RSPO-binding agent that inhibits binding of at least one RSPO to at least one LGR protein further inhibits  $\beta$ -catenin signaling. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody h131R008. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody h131R010. In certain embodiments, a RSPO3-binding agent that inhibits binding of human RSPO3 to at least one LGR protein is antibody h131R011.

**[0164]** In certain embodiments, the RSPO-binding agent (e.g., antibody) blocks binding of at least one RSPO to a receptor. In certain embodiments, the RSPO-binding agent blocks binding of a human RSPO protein to one or more of its receptors. In some embodiments, the RSPO-binding agent blocks binding of a RSPO to at least one LGR protein. In some embodiments, the RSPO-binding agent blocks binding of at least one RSPO protein to LGR4, LGR5, and/or LGR6. In some embodiments, a RSPO3-binding agent blocks binding of RSPO3 to LGR4. In some embodiments, a RSPO3-binding agent blocks binding of RSPO3 to LGR5. In some embodiments, a RSPO3-binding agent blocks binding of RSPO3 to LGR6. In certain embodiments, the blocking of binding of a RSPO-binding agent to at least one LGR protein is at least about 10%, at least about 25%, at least about 50%, at least about 75%, at least about 90%, or at least about 95%. In certain embodiments, a RSPO-binding agent that blocks binding of at least one RSPO protein to at least one LGR protein further inhibits  $\beta$ -catenin signaling. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In certain embodiments, a RSPO3-binding agent

that blocks binding of human RSPO3 to at least one LGR protein is antibody h131R006A or antibody h131R006B. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is antibody h131R005/131R007. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is antibody h131R008. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is antibody h131R010. In certain embodiments, a RSPO3-binding agent that blocks binding of human RSPO3 to at least one LGR protein is antibody h131R011.

[0165] In certain embodiments, the RSPO-binding agent (e.g., an antibody) inhibits  $\beta$ -catenin signaling. It is understood that a RSPO-binding agent that inhibits  $\beta$ -catenin signaling may, in certain embodiments, inhibit signaling by one or more receptors in the  $\beta$ -catenin signaling pathway but not necessarily inhibit signaling by all receptors. In certain alternative embodiments,  $\beta$ -catenin signaling by all human receptors may be inhibited. In certain embodiments,  $\beta$ -catenin signaling by one or more receptors selected from the group consisting of LGR4, LGR5, and LGR6 is inhibited. In certain embodiments, the inhibition of  $\beta$ -catenin signaling by a RSPO-binding agent is a reduction in the level of  $\beta$ -catenin signaling of at least about 10%, at least about 25%, at least about 50%, at least about 75%, at least about 90%, or at least about 95%. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody 131R002, antibody 131R003, or a variant of 131R003. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is a humanized version of antibody 131R002, antibody 131R003, or a variant of 131R003. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R006A or antibody h131R006B. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R005/131R007. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R008. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R010. In some embodiments, a RSPO3-binding agent that inhibits  $\beta$ -catenin signaling is antibody h131R011.

[0166] In certain embodiments, the RSPO-binding agent (e.g., an antibody) inhibits activation of  $\beta$ -catenin. It is understood that a RSPO-binding agent that inhibits activation of  $\beta$ -catenin may, in certain embodiments, inhibit activation of  $\beta$ -catenin by one or more receptors, but not necessarily inhibit activation of  $\beta$ -catenin by all receptors. In certain alternative embodiments, activation of  $\beta$ -catenin by all human receptors may be inhibited. In certain embodiments, activation of  $\beta$ -catenin by one or more receptors selected from the group consisting of LGR4, LGR5, and LGR6 is inhibited. In certain embodiments, the inhibition of activation of  $\beta$ -catenin by a RSPO-binding agent is a reduction in the level of activation of  $\beta$ -catenin of at least about 10%, at least about 25%, at least about 50%, at least about 75%, at least about 90%, or at least about 95%. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody 131R002, antibody 131R003, or a variant of 131R003. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is a humanized version of

antibody 131R002, antibody 131R003, or a variant of 131R003. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody h131R006A or antibody h131R006B. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody h131R005/131R007. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody h131R008. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody h131R010. In some embodiments, a RSPO3-binding agent that inhibits activation of  $\beta$ -catenin is antibody h131R011.

[0167] *In vivo* and *in vitro* assays for determining whether a RSPO-binding agent (or candidate RSPO-binding agent) inhibits  $\beta$ -catenin signaling are known in the art. For example, cell-based, luciferase reporter assays utilizing a TCF/Luc reporter vector containing multiple copies of the TCF-binding domain upstream of a firefly luciferase reporter gene may be used to measure  $\beta$ -catenin signaling levels *in vitro* (Gazit et al., 1999, *Oncogene*, 18; 5959-66; TOPflash, Millipore, Billerica MA). The level of  $\beta$ -catenin signaling in the presence of one or more Wnts (e.g., Wnt(s) expressed by transfected cells or provided by Wnt-conditioned media) with or without a RSPO protein or RSPO-conditioned media in the presence of a RSPO-binding agent is compared to the level of signaling without the RSPO-binding agent present. In addition to the TCF/Luc reporter assay, the effect of a RSPO-binding agent (or candidate agent) on  $\beta$ -catenin signaling may be measured *in vitro* or *in vivo* by measuring the effect of the agent on the level of expression of  $\beta$ -catenin-regulated genes, such as c-myc (He et al., 1998, *Science*, 281:1509-12), cyclin D1 (Tetsu et al., 1999, *Nature*, 398:422-6) and/or fibronectin (Gradl et al. 1999, *Mol. Cell Biol.*, 19:5576-87). In certain embodiments, the effect of a RSPO-binding agent on  $\beta$ -catenin signaling may also be assessed by measuring the effect of the agent on the phosphorylation state of Dishevelled-1, Dishevelled-2, Dishevelled-3, LRP5, LRP6, and/or  $\beta$ -catenin.

[0168] In certain embodiments, the RSPO3-binding agents have one or more of the following effects: inhibit proliferation of tumor cells, inhibit tumor growth, reduce the tumorigenicity of a tumor, reduce the tumorigenicity of a tumor by reducing the frequency of cancer stem cells in the tumor, inhibit tumor growth, trigger cell death of tumor cells, induce cells in a tumor to differentiate, differentiate tumorigenic cells to a non-tumorigenic state, induce expression of differentiation markers in the tumor cells, prevent metastasis of tumor cells, decrease survival of tumor cells, or modulate angiogenesis.

[0169] In certain embodiments, the RSPO3-binding agents are capable of inhibiting tumor growth. In certain embodiments, the RSPO3-binding agents are capable of inhibiting tumor growth *in vivo* (e.g., in a xenograft mouse model, and/or in a human having cancer). In certain embodiments, tumor growth is inhibited at least about two-fold, about three-fold, about five-fold, about ten-fold, about 50-fold, about 100-fold, or about 1000-fold as compared to a untreated tumor.

[0170] In certain embodiments, the RSPO3-binding agents are capable of reducing the tumorigenicity of a tumor. In certain embodiments, the RSPO3-binding agent or antibody is capable of reducing the



tumorigenicity of a tumor comprising cancer stem cells in an animal model, such as a mouse xenograft model. In certain embodiments, the RSPO3-binding agent or antibody is capable of reducing the tumorigenicity of a tumor by decreasing the number or frequency of cancer stem cells in the tumor. In certain embodiments, the number or frequency of cancer stem cells in a tumor is reduced by at least about two-fold, about three-fold, about five-fold, about ten-fold, about 50-fold, about 100-fold, or about 1000-fold. In certain embodiments, the reduction in the number or frequency of cancer stem cells is determined by limiting dilution assay using an animal model. Additional examples and guidance regarding the use of limiting dilution assays to determine a reduction in the number or frequency of cancer stem cells in a tumor can be found, e.g., in International Publication Number WO 2008/042236, U.S. Patent Publication No. 2008/0064049, and U.S. Patent Publication No. 2008/0178305.

[0171] In certain embodiments, the RSPO3-binding agents described herein have a circulating half-life in mice, cynomolgus monkeys, or humans of at least about 2 hours, at least about 5 hours, at least about 10 hours, at least about 24 hours, at least about 3 days, at least about 1 week, or at least about 2 weeks. In certain embodiments, the RSPO3-binding agent is an IgG (e.g., IgG1 or IgG2) antibody that has a circulating half-life in mice, cynomolgus monkeys, or humans of at least about 2 hours, at least about 5 hours, at least about 10 hours, at least about 24 hours, at least about 3 days, at least about 1 week, or at least about 2 weeks. Methods of increasing (or decreasing) the half-life of agents such as polypeptides and antibodies are known in the art. For example, known methods of increasing the circulating half-life of IgG antibodies include the introduction of mutations in the Fc region which increase the pH-dependent binding of the antibody to the neonatal Fc receptor (FcRn) at pH 6.0 (see, e.g., U.S. Patent Publication Nos. 2005/0276799, 2007/0148164, and 2007/0122403). Known methods of increasing the circulating half-life of antibody fragments lacking the Fc region include such techniques as PEGylation.

[0172] In some embodiments, the RSPO3-binding agents are polyclonal antibodies. Polyclonal antibodies can be prepared by any known method. In some embodiments, polyclonal antibodies are produced by immunizing an animal (e.g., a rabbit, rat, mouse, goat, donkey) with an antigen of interest (e.g., a purified peptide fragment, full-length recombinant protein, or fusion protein) using multiple subcutaneous or intraperitoneal injections. The antigen can be optionally conjugated to a carrier such as keyhole limpet hemocyanin (KLH) or serum albumin. The antigen (with or without a carrier protein) is diluted in sterile saline and usually combined with an adjuvant (e.g., Complete or Incomplete Freund's Adjuvant) to form a stable emulsion. After a sufficient period of time, polyclonal antibodies are recovered from the immunized animal, usually from blood or ascites. The polyclonal antibodies can be purified from serum or ascites according to standard methods in the art including, but not limited to, affinity chromatography, ion-exchange chromatography, gel electrophoresis, and dialysis.

[0173] In some embodiments, the RSPO3-binding agents are monoclonal antibodies. Monoclonal antibodies can be prepared using hybridoma methods known to one of skill in the art (see e.g., Kohler and

Milstein, 1975, *Nature*, 256:495-497). In some embodiments, using the hybridoma method, a mouse, hamster, or other appropriate host animal, is immunized as described above to elicit from lymphocytes the production of antibodies that specifically bind the immunizing antigen. In some embodiments, lymphocytes can be immunized *in vitro*. In some embodiments, the immunizing antigen can be a human protein or a portion thereof. In some embodiments, the immunizing antigen can be a mouse protein or a portion thereof.

[0174] Following immunization, lymphocytes are isolated and fused with a suitable myeloma cell line using, for example, polyethylene glycol. The hybridoma cells are selected using specialized media as known in the art and unfused lymphocytes and myeloma cells do not survive the selection process. Hybridomas that produce monoclonal antibodies directed specifically against a chosen antigen may be identified by a variety of methods including, but not limited to, immunoprecipitation, immunoblotting, and *in vitro* binding assays (e.g., flow cytometry, FACS, ELISA, and radioimmunoassay). The hybridomas can be propagated either in *in vitro* culture using standard methods (J.W. Goding, 1996, *Monoclonal Antibodies: Principles and Practice*, 3<sup>rd</sup> Edition, Academic Press, San Diego, CA) or *in vivo* as ascites tumors in an animal. The monoclonal antibodies can be purified from the culture medium or ascites fluid according to standard methods in the art including, but not limited to, affinity chromatography, ion-exchange chromatography, gel electrophoresis, and dialysis.

[0175] In certain embodiments, monoclonal antibodies can be made using recombinant DNA techniques as known to one skilled in the art. The polynucleotides encoding a monoclonal antibody are isolated from mature B-cells or hybridoma cells, such as by RT-PCR using oligonucleotide primers that specifically amplify the genes encoding the heavy and light chains of the antibody, and their sequence is determined using standard techniques. The isolated polynucleotides encoding the heavy and light chains are then cloned into suitable expression vectors which produce the monoclonal antibodies when transfected into host cells such as *E. coli*, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin proteins.

[0176] In certain other embodiments, recombinant monoclonal antibodies, or fragments thereof, can be isolated from phage display libraries expressing variable domains or CDRs of a desired species (see e.g., McCafferty et al., 1990, *Nature*, 348:552-554; Clackson et al., 1991, *Nature*, 352:624-628; and Marks et al., 1991, *J. Mol. Biol.*, 222:581-597).

[0177] The polynucleotide(s) encoding a monoclonal antibody can be modified, for example, by using recombinant DNA technology to generate alternative antibodies. In some embodiments, the constant domains of the light and heavy chains of, for example, a mouse monoclonal antibody can be substituted for those regions of, for example, a human antibody to generate a chimeric antibody, or for a non-immunoglobulin polypeptide to generate a fusion antibody. In some embodiments, the constant regions are truncated or removed to generate the desired antibody fragment of a monoclonal antibody. Site-

directed or high-density mutagenesis of the variable region can be used to optimize specificity, affinity, etc. of a monoclonal antibody.

**[0178]** In some embodiments, a monoclonal antibody against human RSPO3 is a humanized antibody. Typically, humanized antibodies are human immunoglobulins in which residues from the CDRs are replaced by residues from a CDR of a non-human species (e.g., mouse, rat, rabbit, hamster, etc.) that have the desired specificity, affinity, and/or binding capability using methods known to one skilled in the art. In some embodiments, the Fv framework region residues of a human immunoglobulin are replaced with the corresponding residues in an antibody from a non-human species that has the desired specificity, affinity, and/or binding capability. In some embodiments, a humanized antibody can be further modified by the substitution of additional residues either in the Fv framework region and/or within the replaced non-human residues to refine and optimize antibody specificity, affinity, and/or capability. In general, a humanized antibody will comprise substantially all of at least one, and typically two or three, variable domain regions containing all, or substantially all, of the CDRs that correspond to the non-human immunoglobulin whereas all, or substantially all, of the framework regions are those of a human immunoglobulin consensus sequence. In some embodiments, a humanized antibody can also comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. In certain embodiments, such humanized antibodies are used therapeutically because they may reduce antigenicity and HAMA (human anti-mouse antibody) responses when administered to a human subject. One skilled in the art would be able to obtain a functional humanized antibody with reduced immunogenicity following known techniques (see e.g., U.S. Patent Nos. 5,225,539; 5,585,089; 5,693,761; and 5,693,762).

**[0179]** In certain embodiments, the RSPO3-binding agent is a human antibody. Human antibodies can be directly prepared using various techniques known in the art. In some embodiments, human antibodies may be generated from immortalized human B lymphocytes immunized *in vitro* or from lymphocytes isolated from an immunized individual. In either case, cells that produce an antibody directed against a target antigen can be generated and isolated (see, e.g., Cole et al., 1985, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77; Boemer et al., 1991, *J. Immunol.*, 147:86-95; and U.S. Patent Nos. 5,750,373; 5,567,610 and 5,229,275). In some embodiments, the human antibody can be selected from a phage library, where that phage library expresses human antibodies (Vaughan et al., 1996, *Nature Biotechnology*, 14:309-314; Sheets et al., 1998, *PNAS*, 95:6157-6162; Hoogenboom and Winter, 1991, *J. Mol. Biol.*, 227:381; Marks et al., 1991, *J. Mol. Biol.*, 222:581). Alternatively, phage display technology can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable domain gene repertoires from unimmunized donors. Techniques for the generation and use of antibody phage libraries are also described in U.S. Patent Nos. 5,969,108; 6,172,197; 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915; 6,593,081; 6,300,064; 6,653,068; 6,706,484; and 7,264,963; and Rothe

et al., 2008, *J. Mol. Bio.*, 376:1182-1200. Once antibodies are identified, affinity maturation strategies known in the art, including but not limited to, chain shuffling (Marks et al., 1992, *Bio/Technology*, 10:779-783) and site-directed mutagenesis, may be employed to generate high affinity human antibodies.

**[0180]** In some embodiments, human antibodies can be made in transgenic mice that contain human immunoglobulin loci. Upon immunization these mice are capable of producing the full repertoire of human antibodies in the absence of endogenous immunoglobulin production. This approach is described in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016.

**[0181]** This invention also encompasses bispecific antibodies that specifically recognize at least one human RSPO protein. Bispecific antibodies are capable of specifically recognizing and binding at least two different antigens or epitopes. The different epitopes can either be within the same molecule (e.g., two epitopes on human RSPO3) or on different molecules (e.g., one epitope on RSPO3 and one epitope on RSPO2). In some embodiments, a bispecific antibody has enhanced potency as compared to an individual antibody or to a combination of more than one antibody. In some embodiments, a bispecific antibody has reduced toxicity as compared to an individual antibody or to a combination of more than one antibody. It is known to those of skill in the art that any binding agent (e.g., antibody) may have unique pharmacokinetics (PK) (e.g., circulating half-life). In some embodiments, a bispecific antibody has the ability to synchronize the PK of two active binding agents wherein the two individual binding agents have different PK profiles. In some embodiments, a bispecific antibody has the ability to concentrate the actions of two binding agents (e.g., antibodies) in a common area (e.g., a tumor and/or tumor microenvironment). In some embodiments, a bispecific antibody has the ability to concentrate the actions of two binding agents (e.g., antibodies) to a common target (e.g., a tumor or a tumor cell). In some embodiments, a bispecific antibody has the ability to target the actions of two binding agents (e.g., antibodies) to more than one biological pathway or function.

**[0182]** In certain embodiments, the bispecific antibody specifically binds RSPO3 and a second target. In certain embodiments, the bispecific antibody specifically binds RSPO3 and a second human RSPO (e.g., RSPO1, RSPO2, or RSPO4). In certain embodiments, the bispecific antibody specifically binds RSPO3 and RSPO2. In some embodiments, the bispecific antibody is a monoclonal antibody. In some embodiments, the bispecific antibody is a humanized antibody. In some embodiments, the bispecific antibody is a human antibody. In some embodiments, the bispecific antibody is an IgG1 antibody. In some embodiments, the bispecific antibody is an IgG2 antibody. In some embodiments, the bispecific antibody has decreased toxicity and/or side effects. In some embodiments, the bispecific antibody has decreased toxicity and/or side effects as compared to a mixture of the two individual antibodies or the antibodies as single agents. In some embodiments, the bispecific antibody has an increased therapeutic index. In some embodiments, the bispecific antibody has an increased therapeutic index as compared to a mixture of the two individual antibodies or the antibodies as single agents.

**[0183]** In some embodiments, the antibodies can specifically recognize and bind a first antigen target, (e.g., RSPO3) as well as a second antigen target, such as an effector molecule on a leukocyte (e.g., CD2, CD3, CD28, CTLA-4, CD80, or CD86) or a Fc receptor (e.g., CD64, CD32, or CD16) so as to focus cellular defense mechanisms to the cell expressing and/or producing the first antigen target. In some embodiments, the antibodies can be used to direct cytotoxic agents to cells which express a particular target antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA.

**[0184]** Techniques for making bispecific antibodies are known by those skilled in the art, see for example, Millstein et al., 1983, *Nature*, 305:537-539; Brennan et al., 1985, *Science*, 229:81; Suresh et al., 1986, *Methods in Enzymol.*, 121:120; Traunecker et al., 1991, *EMBO J.*, 10:3655-3659; Shalaby et al., 1992, *J. Exp. Med.*, 175:217-225; Kostelny et al., 1992, *J. Immunol.*, 148:1547-1553; Gruber et al., 1994, *J. Immunol.*, 152:5368; U.S. Patent No. 5,731,168; International Publication No. WO 2009/089004; and U.S. Patent Publication No. 2011/0123532. In some embodiments, the bispecific antibodies comprise heavy chain constant regions with modifications in the amino acids which are part of the interface between the two heavy chains. In some embodiments, the bispecific antibodies can be generated using a “knobs-into-holes” strategy (see, e.g., U.S. Patent No. 5,731,168; Ridgway et. al., 1996, *Prot. Engin.*, 9:617-621). In some cases, the “knobs” and “holes” terminology is replaced with the terms “protuberances” and “cavities”. In some embodiments, the bispecific antibodies may comprise variant hinge regions incapable of forming disulfide linkages between the heavy chains (see, e.g., WO 2006/028936). In some embodiments, the modifications may comprise changes in amino acids that result in altered electrostatic interactions. In some embodiments, the modifications may comprise changes in amino acids that result in altered hydrophobic/hydrophilic interactions.

**[0185]** Bispecific antibodies can be intact antibodies or antibody fragments comprising antigen-binding sites. Antibodies with more than two valencies are also contemplated. For example, trispecific antibodies can be prepared (Tutt et al., 1991, *J. Immunol.*, 147:60). Thus, in certain embodiments the antibodies to RSPO3 are multispecific.

**[0186]** In certain embodiments, the antibodies (or other polypeptides) described herein may be monospecific. In certain embodiments, each of the one or more antigen-binding sites that an antibody contains is capable of binding (or binds) a homologous epitope on RSPO proteins. In certain embodiments, an antigen-binding site of a monospecific antibody described herein is capable of binding (or binds), for example, RSPO3 and RSPO2 (i.e., the same epitope is found on both RSPO3 and RSPO2 proteins).

**[0187]** In certain embodiments, the RSPO3-binding agent is an antibody fragment. Antibody fragments may have different functions or capabilities than intact antibodies; for example, antibody fragments can have increased tumor penetration. Various techniques are known for the production of antibody

fragments including, but not limited to, proteolytic digestion of intact antibodies. In some embodiments, antibody fragments include a F(ab')<sub>2</sub> fragment produced by pepsin digestion of an antibody molecule. In some embodiments, antibody fragments include a Fab fragment generated by reducing the disulfide bridges of an F(ab')<sub>2</sub> fragment. In other embodiments, antibody fragments include a Fab fragment generated by the treatment of the antibody molecule with papain and a reducing agent. In certain embodiments, antibody fragments are produced recombinantly. In some embodiments, antibody fragments include Fv or single chain Fv (scFv) fragments. Fab, Fv, and scFv antibody fragments can be expressed in and secreted from *E. coli* or other host cells, allowing for the production of large amounts of these fragments. In some embodiments, antibody fragments are isolated from antibody phage libraries as discussed herein. For example, methods can be used for the construction of Fab expression libraries (Huse et al., 1989, *Science*, 246:1275-1281) to allow rapid and effective identification of monoclonal Fab fragments with the desired specificity for a RSPO protein or derivatives, fragments, analogs or homologs thereof. In some embodiments, antibody fragments are linear antibody fragments. In certain embodiments, antibody fragments are monospecific or bispecific. In certain embodiments, the RSPO3-binding agent is a scFv. Various techniques can be used for the production of single-chain antibodies specific to one or more human RSPOs (see, e.g., U.S. Patent No. 4,946,778).

**[0188]** It can further be desirable, especially in the case of antibody fragments, to modify an antibody in order to alter (e.g., increase or decrease) its serum half-life. This can be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle (e.g., by DNA or peptide synthesis).

**[0189]** Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune cells to unwanted cells (see, e.g., U.S. Patent No. 4,676,980). It is also contemplated that the heteroconjugate antibodies can be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

**[0190]** For the purposes of the present invention, it should be appreciated that modified antibodies can comprise any type of variable region that provides for the association of the antibody with the target (i.e., human RSPO3). In this regard, the variable region may comprise or be derived from any type of mammal that can be induced to mount a humoral response and generate immunoglobulins against the desired antigen. As such, the variable region of the modified antibodies can be, for example, of human, murine, non-human primate (e.g. cynomolgus monkeys, macaques, etc.) or rabbit origin. In some embodiments, both the variable and constant regions of the modified immunoglobulins are human. In other

embodiments, the variable regions of compatible antibodies (usually derived from a non-human source) can be engineered or specifically tailored to improve the binding properties or reduce the immunogenicity of the molecule. In this respect, variable regions useful in the present invention can be humanized or otherwise altered through the inclusion of imported amino acid sequences.

**[0191]** In certain embodiments, the variable domains in both the heavy and light chains are altered by at least partial replacement of one or more CDRs and, if necessary, by partial framework region replacement and sequence modification and/or alteration. Although the CDRs may be derived from an antibody of the same class or even subclass as the antibody from which the framework regions are derived, it is envisaged that the CDRs may be derived from an antibody of different class and often from an antibody from a different species. It may not be necessary to replace all of the CDRs with all of the CDRs from the donor variable region to transfer the antigen binding capacity of one variable domain to another. Rather, it may only be necessary to transfer those residues that are required to maintain the activity of the antigen-binding site.

**[0192]** Alterations to the variable region notwithstanding, those skilled in the art will appreciate that the modified antibodies of this invention will comprise antibodies (e.g., full-length antibodies or immunoreactive fragments thereof) in which at least a fraction of one or more of the constant region domains has been deleted or otherwise altered so as to provide desired biochemical characteristics such as increased tumor localization or increased serum half-life when compared with an antibody of approximately the same immunogenicity comprising a native or unaltered constant region. In some embodiments, the constant region of the modified antibodies will comprise a human constant region. Modifications to the constant region compatible with this invention comprise additions, deletions or substitutions of one or more amino acids in one or more domains. The modified antibodies disclosed herein may comprise alterations or modifications to one or more of the three heavy chain constant domains (CH1, CH2 or CH3) and/or to the light chain constant domain (CL). In some embodiments, one or more domains are partially or entirely deleted from the constant regions of the modified antibodies. In some embodiments, the modified antibodies will comprise domain deleted constructs or variants wherein the entire CH2 domain has been removed ( $\Delta$ CH2 constructs). In some embodiments, the omitted constant region domain is replaced by a short amino acid spacer (e.g., 10 amino acid residues) that provides some of the molecular flexibility typically imparted by the absent constant region.

**[0193]** In some embodiments, the modified antibodies are engineered to fuse the CH3 domain directly to the hinge region of the antibody. In other embodiments, a peptide spacer is inserted between the hinge region and the modified CH2 and/or CH3 domains. For example, constructs may be expressed wherein the CH2 domain has been deleted and the remaining CH3 domain (modified or unmodified) is joined to the hinge region with a 5-20 amino acid spacer. Such a spacer may be added to ensure that the regulatory elements of the constant domain remain free and accessible or that the hinge region remains flexible.

However, it should be noted that amino acid spacers may, in some cases, prove to be immunogenic and elicit an unwanted immune response against the construct. Accordingly, in certain embodiments, any spacer added to the construct will be relatively non-immunogenic so as to maintain the desired biological qualities of the modified antibodies.

[0194] In some embodiments, the modified antibodies may have only a partial deletion of a constant domain or substitution of a few or even a single amino acid. For example, the mutation of a single amino acid in selected areas of the CH2 domain may be enough to substantially reduce Fc binding and thereby increase cancer cell localization and/or tumor penetration. Similarly, it may be desirable to simply delete the part of one or more constant region domains that control a specific effector function (e.g. complement C1q binding) to be modulated. Such partial deletions of the constant regions may improve selected characteristics of the antibody (serum half-life) while leaving other desirable functions associated with the subject constant region domain intact. Moreover, as alluded to above, the constant regions of the disclosed antibodies may be modified through the mutation or substitution of one or more amino acids that enhances the profile of the resulting construct. In this respect it may be possible to disrupt the activity provided by a conserved binding site (e.g., Fc binding) while substantially maintaining the configuration and immunogenic profile of the modified antibody. In certain embodiments, the modified antibodies comprise the addition of one or more amino acids to the constant region to enhance desirable characteristics such as decreasing or increasing effector function or provide for more cytotoxin or carbohydrate attachment sites.

[0195] It is known in the art that the constant region mediates several effector functions. For example, binding of the C1 component of complement to the Fc region of IgG or IgM antibodies (bound to antigen) activates the complement system. Activation of complement is important in the opsonization and lysis of cell pathogens. The activation of complement also stimulates the inflammatory response and can also be involved in autoimmune hypersensitivity. In addition, the Fc region of an antibody can bind a cell expressing a Fc receptor (FcR). There are a number of Fc receptors which are specific for different classes of antibody, including IgG (gamma receptors), IgE (epsilon receptors), IgA (alpha receptors) and IgM (mu receptors). Binding of antibody to Fc receptors on cell surfaces triggers a number of important and diverse biological responses including engulfment and destruction of antibody-coated particles, clearance of immune complexes, lysis of antibody-coated target cells by killer cells (called antibody-dependent cell cytotoxicity or ADCC), release of inflammatory mediators, placental transfer, and control of immunoglobulin production.

[0196] In certain embodiments, the modified antibodies provide for altered effector functions that, in turn, affect the biological profile of the administered antibody. For example, in some embodiments, the deletion or inactivation (through point mutations or other means) of a constant region domain may reduce Fc receptor binding of the circulating modified antibody thereby increasing cancer cell localization and/or



tumor penetration. In other embodiments, the constant region modifications increase the serum half-life of the antibody. In other embodiments, the constant region modifications reduce the serum half-life of the antibody. In some embodiments, the constant region is modified to eliminate disulfide linkages or oligosaccharide moieties. Modifications to the constant region in accordance with this invention may easily be made using well known biochemical or molecular engineering techniques.

**[0197]** In certain embodiments, a RSPO3-binding agent that is an antibody does not have one or more effector functions. For instance, in some embodiments, the antibody has no ADCC activity, and/or no complement-dependent cytotoxicity (CDC) activity. In certain embodiments, the antibody does not bind an Fc receptor, and/or complement factors. In certain embodiments, the antibody has no effector function.

**[0198]** The present invention further embraces variants and equivalents which are substantially homologous to the chimeric, humanized, and human antibodies, or antibody fragments thereof, set forth herein. These can contain, for example, conservative substitution mutations, i.e. the substitution of one or more amino acids by similar amino acids. For example, conservative substitution refers to the substitution of an amino acid with another amino acid within the same general class such as, for example, one acidic amino acid with another acidic amino acid, one basic amino acid with another basic amino acid or one neutral amino acid by another neutral amino acid. What is intended by a conservative amino acid substitution is well known in the art and described herein.

**[0199]** Thus, the present invention provides methods for producing an antibody that binds RSPO3, including bispecific antibodies that specifically bind both RSPO3 and a second target (e.g., a human RSPO). In some embodiments, the method for producing an antibody that binds RSPO3 comprises using hybridoma techniques. In some embodiments, a method for producing an antibody that binds human RSPO3 is provided. In some embodiments, the method comprises using amino acids 22-272 of human RSPO3. In some embodiments, the method comprises using amino acids 22-272 of SEQ ID NO:3. In some embodiments, the method of generating an antibody that binds RSPO3 comprises screening a human phage library. The present invention further provides methods of identifying an antibody that binds RSPO3. In some embodiments, the antibody is identified by FACS screening for binding to RSPO3 or a portion thereof. In some embodiments, the antibody is identified by FACS screening for binding to RSPO3 and a second RSPO or a portion thereof. In some embodiments, the antibody is identified by FACS screening for binding to both RSPO3 and RSPO2 or a portion thereof. In some embodiments, the antibody is identified by screening using ELISA for binding to RSPO3. In some embodiments, the antibody is identified by screening using ELISA for binding to RSPO3 and a second RSPO. In some embodiments, the antibody is identified by screening using ELISA for binding to both RSPO3 and RSPO2. In some embodiments, the antibody is identified by screening by FACS for blocking of binding of RSPO3 to a human LGR protein. In some embodiments, the antibody is identified by screening for inhibition or blocking of  $\beta$ -catenin signaling.

[0200] In some embodiments, a method of generating an antibody to human RSPO3 protein comprises immunizing a mammal with a polypeptide comprising amino acids 22-272 of human RSPO3. In some embodiments, a method of generating an antibody to human RSPO3 protein comprises immunizing a mammal with a polypeptide comprising at least a portion of amino acids 22-272 of human RSPO3. In some embodiments, the method further comprises isolating antibodies or antibody-producing cells from the mammal. In some embodiments, a method of generating a monoclonal antibody which binds RSPO3 protein comprises: (a) immunizing a mammal with a polypeptide comprising at least a portion of amino acids 22-272 of human RSPO3; (b) isolating antibody producing cells from the immunized mammal; (c) fusing the antibody-producing cells with cells of a myeloma cell line to form hybridoma cells. In some embodiments, the method further comprises (d) selecting a hybridoma cell expressing an antibody that binds RSPO3 protein. In some embodiments, the at least a portion of amino acids 22-272 of human RSPO3 is selected from the group consisting of SEQ ID NOs:5-8. In some embodiments, the at least a portion of amino acids 22-272 of human RSPO3 is SEQ ID NO:5. In some embodiments, the at least a portion of amino acids 22-272 of human RSPO3 is SEQ ID NO:6 or SEQ ID NO:7. In some embodiments, the at least a portion of amino acids 22-272 of human RSPO3 is SEQ ID NO:6 and SEQ ID NO:7. In certain embodiments, the mammal is a mouse. In some embodiments, the antibody is selected using a polypeptide comprising at least a portion of amino acid 22-272 of human RSPO3. In certain embodiments, the polypeptide used for selection comprising at least a portion of amino acids 22-272 of human RSPO3 is selected from the group consisting of SEQ ID NOs:5-8. In some embodiments, the antibody binds RSPO3 and at least one other RSPO protein. In certain embodiments, the at least one other RSPO protein is selected from the group consisting of RSPO1, RSPO2, and RSPO4. In certain embodiments, the antibody binds RSPO3 and RSPO1. In certain embodiments, the antibody binds RSPO3 and RSPO2. In certain embodiments, the antibody binds RSPO3 and RSPO4. In certain embodiments, the antibody binds RSPO3, RSPO1, and RSPO2. In certain embodiments, the antibody binds RSPO3, RSPO1, and RSPO4. In certain embodiments, the antibody binds RSPO3, RSPO2, and RSPO4. In some embodiments, the antibody binds both human RSPO3 and mouse RSPO3.

[0201] In some embodiments, the antibody generated by the methods described herein is a RSPO antagonist, particularly a RSPO3 antagonist. In some embodiments, the antibody generated by the methods described herein inhibits  $\beta$ -catenin signaling.

[0202] In some embodiments, a method of producing an antibody to at least one human RSPO protein comprises identifying an antibody using a membrane-bound heterodimeric molecule comprising a single antigen-binding site. In some non-limiting embodiments, the antibody is identified using methods and polypeptides described in International Publication WO 2011/100566.

[0203] In some embodiments, a method of producing an antibody to at least one human RSPO protein comprises screening an antibody-expressing library for antibodies that bind a human RSPO protein. In

some embodiments, the antibody-expressing library is a phage library. In some embodiments, the screening comprises panning. In some embodiments, the antibody-expressing library is a phage library. In some embodiments, the antibody-expressing library is a mammalian cell library. In some embodiments, the antibody-expressing library is screened using at least a portion of amino acids 22-272 of human RSPO3. In some embodiments, antibodies identified in the first screening, are screened again using a different RSPO protein thereby identifying an antibody that binds RSPO3 and a second RSPO protein. In certain embodiments, the polypeptide used for screening comprises at least a portion of amino acids 22-272 of human RSPO3 selected from the group consisting of SEQ ID NOs:5-8. In some embodiments, the antibody identified in the screening binds RSPO3 and at least one other RSPO protein. In certain embodiments, the at least one other RSPO protein is selected from the group consisting of RSPO1, RSPO2, and RSPO4. In certain embodiments, the antibody identified in the screening binds RSPO3 and RSPO1. In certain embodiments, the antibody identified in the screening binds RSPO3 and RSPO2. In certain embodiments, the antibody identified in the screening binds RSPO3 and RSPO4. In some embodiments, the antibody identified in the screening binds both human RSPO3 and mouse RSPO3. In some embodiments, the antibody identified in the screening is a RSPO3 antagonist. In some embodiments, the antibody identified in the screening inhibits  $\beta$ -catenin signaling induced by RSPO3.

[0204] In certain embodiments, the antibodies described herein are isolated. In certain embodiments, the antibodies described herein are substantially pure.

[0205] In some embodiments of the present invention, the RSPO3-binding agents are polypeptides. The polypeptides can be recombinant polypeptides, natural polypeptides, or synthetic polypeptides comprising an antibody, or fragment thereof, that bind RSPO3. It will be recognized in the art that some amino acid sequences of the invention can be varied without significant effect of the structure or function of the protein. Thus, the invention further includes variations of the polypeptides which show substantial activity or which include regions of an antibody, or fragment thereof, against human RSPO3. In some embodiments, amino acid sequence variations of RSPO-binding polypeptides include deletions, insertions, inversions, repeats, and/or other types of substitutions.

[0206] In certain embodiments, the polypeptides described herein are isolated. In certain embodiments, the polypeptides described herein are substantially pure.

[0207] The polypeptides, analogs and variants thereof, can be further modified to contain additional chemical moieties not normally part of the polypeptide. The derivatized moieties can improve or otherwise modulate the solubility, the biological half-life, and/or absorption of the polypeptide. The moieties can also reduce or eliminate undesirable side effects of the polypeptides and variants. An overview for chemical moieties can be found in *Remington: The Science and Practice of Pharmacy*, 22<sup>nd</sup> Edition, 2012, Pharmaceutical Press, London.

[0208] The polypeptides described herein can be produced by any suitable method known in the art. Such methods range from direct protein synthesis methods to constructing a DNA sequence encoding polypeptide sequences and expressing those sequences in a suitable host. In some embodiments, a DNA sequence is constructed using recombinant technology by isolating or synthesizing a DNA sequence encoding a wild-type protein of interest. Optionally, the sequence can be mutagenized by site-specific mutagenesis to provide functional analogs thereof. See, e.g., Zoeller et al., 1984, *PNAS*, 81:5662-5066 and U.S. Patent No. 4,588,585.

[0209] In some embodiments, a DNA sequence encoding a polypeptide of interest may be constructed by chemical synthesis using an oligonucleotide synthesizer. Oligonucleotides can be designed based on the amino acid sequence of the desired polypeptide and selecting those codons that are favored in the host cell in which the recombinant polypeptide of interest will be produced. Standard methods can be applied to synthesize a polynucleotide sequence encoding an isolated polypeptide of interest. For example, a complete amino acid sequence can be used to construct a back-translated gene. Further, a DNA oligomer containing a nucleotide sequence coding for the particular isolated polypeptide can be synthesized. For example, several small oligonucleotides coding for portions of the desired polypeptide can be synthesized and then ligated. The individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

[0210] Once assembled (by synthesis, site-directed mutagenesis, or another method), the polynucleotide sequences encoding a particular polypeptide of interest can be inserted into an expression vector and operatively linked to an expression control sequence appropriate for expression of the protein in a desired host. Proper assembly can be confirmed by nucleotide sequencing, restriction enzyme mapping, and/or expression of a biologically active polypeptide in a suitable host. As is well-known in the art, in order to obtain high expression levels of a transfected gene in a host, the gene must be operatively linked to transcriptional and translational expression control sequences that are functional in the chosen expression host.

[0211] In certain embodiments, recombinant expression vectors are used to amplify and express DNA encoding antibodies, or fragments thereof, against human RSPO3. For example, recombinant expression vectors can be replicable DNA constructs which have synthetic or cDNA-derived DNA fragments encoding a polypeptide chain of a RSPO-binding agent, such as an anti-RSPO antibody, or fragment thereof, operatively linked to suitable transcriptional and/or translational regulatory elements derived from mammalian, microbial, viral or insect genes. A transcriptional unit generally comprises an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, transcriptional promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. Regulatory elements can include an operator sequence to control transcription. The ability to

replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants can additionally be incorporated. DNA regions are "operatively linked" when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operatively linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operatively linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operatively linked to a coding sequence if it is positioned so as to permit translation. In some embodiments, structural elements intended for use in yeast expression systems include a leader sequence enabling extracellular secretion of translated protein by a host cell. In other embodiments, in situations where recombinant protein is expressed without a leader or transport sequence, it can include an N-terminal methionine residue. This residue can optionally be subsequently cleaved from the expressed recombinant protein to provide a final product.

[0212] The choice of an expression control sequence and an expression vector depends upon the choice of host. A wide variety of expression host/vector combinations can be employed. Useful expression vectors for eukaryotic hosts include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus, and cytomegalovirus. Useful expression vectors for bacterial hosts include known bacterial plasmids, such as plasmids from *E. coli*, including pCR1, pBR322, pMB9 and their derivatives, and wider host range plasmids, such as M13 and other filamentous single-stranded DNA phages.

[0213] The RSPO-binding agents (e.g., polypeptides or antibodies) of the present invention can be expressed from one or more vectors. For example, in some embodiments, one heavy chain polypeptide is expressed by one vector, a second heavy chain polypeptide is expressed by a second vector and a light chain polypeptide is expressed by a third vector. In some embodiments, a first heavy chain polypeptide and a light chain polypeptide is expressed by one vector and a second heavy chain polypeptide is expressed by a second vector. In some embodiments, two heavy chain polypeptides are expressed by one vector and a light chain polypeptide is expressed by a second vector. In some embodiments, three polypeptides are expressed from one vector. Thus, in some embodiments, a first heavy chain polypeptide, a second heavy chain polypeptide, and a light chain polypeptide are expressed by a single vector.

[0214] Suitable host cells for expression of a RSPO3-binding polypeptide or antibody (or a RSPO protein to use as an antigen) include prokaryotes, yeast cells, insect cells, or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram-negative or gram-positive organisms, for example *E. coli* or *Bacillus*. Higher eukaryotic cells include established cell lines of mammalian origin as described below. Cell-free translation systems may also be employed. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described in Pouwels et al., 1985, *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, NY. Additional information regarding methods of protein production, including antibody production, can be found, e.g.,

in U.S. Patent Publication No. 2008/0187954, U.S. Patent Nos. 6,413,746, 6,660,501; and International Patent Publication No. WO 04/009823.

**[0215]** Various mammalian culture systems may be used to express recombinant polypeptides.

Expression of recombinant proteins in mammalian cells may be desirable because these proteins are generally correctly folded, appropriately modified, and biologically functional. Examples of suitable mammalian host cell lines include, but are not limited to, COS-7 (monkey kidney-derived), L-929 (murine fibroblast-derived), C127 (murine mammary tumor-derived), 3T3 (murine fibroblast-derived), CHO (Chinese hamster ovary-derived), HeLa (human cervical cancer-derived), BHK (hamster kidney fibroblast-derived), HEK-293 (human embryonic kidney-derived) cell lines and variants thereof.

Mammalian expression vectors can comprise non-transcribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking non-transcribed sequences, and 5' or 3' non-translated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

**[0216]** Expression of recombinant proteins in insect cell culture systems (e.g., baculovirus) also offers a robust method for producing correctly folded and biologically functional proteins. Baculovirus systems for production of heterologous proteins in insect cells are well-known to those of skill in the art (see, e.g., Luckow and Summers, 1988, *Bio/Technology*, 6:47).

**[0217]** Thus, the present invention provides cells comprising the RSPO3-binding agents described herein. In some embodiments, the cells produce the RSPO3-binding agents described herein. In certain embodiments, the cells produce an antibody. In some embodiments, the cells produce an antibody that binds human RSPO3. In certain embodiments, the cells produce antibody 131R002. In certain embodiments, the cells produce antibody 131R003. In certain embodiments, the cells produce variants of antibody 131R003. In certain embodiments, the cells produce a humanized version of antibody 131R002, antibody 131R003, or variants of antibody 131R003. In some embodiments, the cells produce a chimeric version of antibody 131R002, antibody 131R003, or variants of antibody 131R003. In some embodiments, the cells produce antibody h131R006A or antibody h131R006B. In some embodiments, the cells produce antibody h131R005/131R007. In some embodiments, the cells produce antibody h131R008. In some embodiments, the cells produce antibody h131R010. In some embodiments, the cells produce antibody h131R011. In some embodiments, the cells produce a bispecific antibody that binds RSPO3. In some embodiments, the cells produce a bispecific antibody that binds RSPO3 and RSPO2. In some embodiments, the cell is a hybridoma cell. In some embodiments, the cell is a mammalian cell. In some embodiments, the cell is a prokaryotic cell. In some embodiments, the cell is an eukaryotic cell.

**[0218]** The proteins produced by a transformed host can be purified according to any suitable method. Standard methods include chromatography (e.g., ion exchange, affinity, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for protein

purification. Affinity tags such as hexa-histidine, maltose binding domain, influenza coat sequence, and glutathione-S-transferase can be attached to the protein to allow easy purification by passage over an appropriate affinity column. Affinity chromatography used for purifying immunoglobulins can include Protein A, Protein G, and Protein L chromatography. Isolated proteins can be physically characterized using such techniques as proteolysis, size exclusion chromatography (SEC), mass spectrometry (MS), nuclear magnetic resonance (NMR), isoelectric focusing (IEF), high performance liquid chromatography (HPLC), and x-ray crystallography. The purity of isolated proteins can be determined using techniques known to those of skill in the art, including but not limited to, SDS-PAGE, SEC, capillary gel electrophoresis, IEF, and capillary isoelectric focusing (cIEF).

**[0219]** In some embodiments, supernatants from expression systems which secrete recombinant protein into culture media can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix. In some embodiments, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose, or other types commonly employed in protein purification. In some embodiments, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. In some embodiments, a hydroxyapatite media can be employed, including but not limited to, ceramic hydroxyapatite (CHT). In certain embodiments, one or more reverse-phase HPLC steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify a recombinant protein (e.g., a RSPO3-binding agent). Some or all of the foregoing purification steps, in various combinations, can be employed to provide a homogeneous recombinant protein.

**[0220]** In some embodiments, heterodimeric proteins such as bispecific antibodies are purified according to any of the methods described herein. In some embodiments, anti-RSPO bispecific antibodies are isolated and/or purified using at least one chromatography step. In some embodiments, the at least one chromatography step comprises affinity chromatography. In some embodiments, the at least one chromatography step further comprises anion exchange chromatography. In some embodiments, the isolated and/or purified antibody product comprises at least 90% heterodimeric antibody. In some embodiments, the isolated and/or purified antibody product comprises at least 95%, 96%, 97%, 98% or 99% heterodimeric antibody. In some embodiments, the isolated and/or purified antibody product comprises about 100% heterodimeric antibody.

**[0221]** In some embodiments, recombinant protein produced in bacterial culture can be isolated, for example, by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange, or size exclusion chromatography steps. HPLC can be employed for final

purification steps. Microbial cells employed in expression of a recombinant protein can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

[0222] Methods known in the art for purifying antibodies and other proteins also include, for example, those described in U.S. Patent Publication Nos. 2008/0312425, 2008/0177048, and 2009/0187005.

[0223] In certain embodiments, the RSPO3-binding agent is a polypeptide that is not an antibody. A variety of methods for identifying and producing non-antibody polypeptides that bind with high affinity to a protein target are known in the art. See, e.g., Skerra, 2007, *Curr. Opin. Biotechnol.*, 18:295-304; Hosse et al., 2006, *Protein Science*, 15:14-27; Gill et al., 2006, *Curr. Opin. Biotechnol.*, 17:653-658; Nygren, 2008, *FEBS J.*, 275:2668-76; and Skerra, 2008, *FEBS J.*, 275:2677-83. In certain embodiments, phage or mammalian display technology may be used to produce and/or identify a RSPO3-binding polypeptide. In certain embodiments, the polypeptide comprises a protein scaffold of a type selected from the group consisting of protein A, protein G, a lipocalin, a fibronectin domain, an ankyrin consensus repeat domain, and thioredoxin.

[0224] In certain embodiments, the RSPO3-binding agents or antibodies can be used in any one of a number of conjugated (i.e. an immunoconjugate or radioconjugate) or non-conjugated forms. In certain embodiments, the antibodies can be used in a non-conjugated form to harness the subject's natural defense mechanisms including complement-dependent cytotoxicity and antibody dependent cellular toxicity to eliminate malignant or cancer cells.

[0225] In some embodiments, the RSPO3-binding agent (e.g., an antibody or polypeptide) is conjugated to a cytotoxic agent. In some embodiments, the cytotoxic agent is a chemotherapeutic agent including, but not limited to, methotrexate, adriamycin, doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents. In some embodiments, the cytotoxic agent is an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof, including, but not limited to, diphtheria A chain, non-binding active fragments of diphtheria toxin, exotoxin A chain, ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), Momordica charantia inhibitor, curcin, crotin, Sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. In some embodiments, the cytotoxic agent is a radioisotope to produce a radioconjugate or a radioconjugated antibody. A variety of radionuclides are available for the production of radioconjugated antibodies including, but not limited to,  $^{90}\text{Y}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{111}\text{In}$ ,  $^{131}\text{In}$ ,  $^{105}\text{Rh}$ ,  $^{153}\text{Sm}$ ,  $^{67}\text{Cu}$ ,  $^{67}\text{Ga}$ ,  $^{166}\text{Ho}$ ,  $^{177}\text{Lu}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$  and  $^{212}\text{Bi}$ . Conjugates of an antibody and one or more small molecule toxins, such as calicheamicins, maytansinoids, trichothenes, and CC1065, and the derivatives of these toxins that have toxin activity, can also be used. Conjugates of an antibody and cytotoxic agent may be made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate



(SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis(p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene).

### III. Polynucleotides

**[0226]** In certain embodiments, the invention encompasses polynucleotides comprising polynucleotides that encode a polypeptide (or a fragment of a polypeptide) that specifically binds RSPO3. The term “polynucleotides that encode a polypeptide” encompasses a polynucleotide which includes only coding sequences for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequences. For example, in some embodiments, the invention provides a polynucleotide comprising a polynucleotide sequence that encodes an antibody to human RSPO3 or encodes a fragment of such an antibody (e.g., a fragment comprising the antigen-binding site). The polynucleotides of the invention can be in the form of RNA or in the form of DNA. DNA includes cDNA, genomic DNA, and synthetic DNA; and can be double-stranded or single-stranded, and if single stranded can be the coding strand or non-coding (anti-sense) strand.

**[0227]** In certain embodiments, the polynucleotide comprises a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:86, SEQ ID NO:87, and SEQ ID NO:88. In some embodiments, the polynucleotide comprises a polynucleotide sequence selected from the group consisting of: SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, and SEQ ID NO:95.

**[0228]** In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:18. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:19. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:20. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:24. In some embodiments, a plasmid

comprises a polynucleotide comprising SEQ ID NO:25. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:26. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:30. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:31. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:32. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:40. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:43. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:50. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:51. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:52. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:53. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:54. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:55. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:65. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:66. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:67. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:70. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:71. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:75. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:76. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:77. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:84. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:85. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:89. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:90. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:91. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:92. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:93. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:94. In some embodiments, a plasmid comprises a polynucleotide comprising SEQ ID NO:95.

**[0229]** In certain embodiments, the polynucleotide comprises a polynucleotide having a nucleotide sequence at least about 80% identical, at least about 85% identical, at least about 90% identical, at least about 95% identical, and in some embodiments, at least about 96%, 97%, 98% or 99% identical to a polynucleotide comprising a sequence selected from the group consisting of SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID

NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, and SEQ ID NO:95. Also provided is a polynucleotide that comprises a polynucleotide that hybridizes to SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:990, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, or SEQ ID NO:95. Also provided is a polynucleotide that comprises a polynucleotide that hybridizes to the complement of SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, or SEQ ID NO:95. In certain embodiments, the hybridization is under conditions of high stringency.

**[0230]** In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:18 and SEQ ID NO:20. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:19 and SEQ ID NO:20. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:50 and SEQ ID NO:20. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:51 and SEQ ID NO:20. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:65 and SEQ ID NO:20. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:18 and SEQ ID NO:75. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:19 and SEQ ID NO:75. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:50 and SEQ ID NO:75. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:51 and SEQ ID NO:75. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:65 and SEQ ID NO:75. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:95 and SEQ ID NO:89. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:92 and SEQ ID NO:89.

**[0231]** In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:24 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:25 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:40 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:43 and SEQ ID NO:26. In some embodiments, an antibody is

encoded by a polynucleotide comprising SEQ ID NO:52 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:53 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:66 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:70 and SEQ ID NO:26. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:24 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:25 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:40 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:43 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:52 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:53 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:66 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:70 and SEQ ID NO:76. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:84 and SEQ ID NO:90. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:93 and SEQ ID NO:90.

[0232] In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:30 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:31 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:54 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:55 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:67 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:71 and SEQ ID NO:32. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:30 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:31 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:54 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:55 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:67 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:71 and SEQ ID NO:77. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:85 and SEQ ID NO:91. In some embodiments, an antibody is encoded by a polynucleotide comprising SEQ ID NO:94 and SEQ ID NO:91.

[0233] In certain embodiments, the polynucleotides comprise the coding sequence for the mature polypeptide fused in the same reading frame to a polynucleotide which aids, for example, in expression

and secretion of a polypeptide from a host cell (e.g., a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell). The polypeptide having a leader sequence is a preprotein and can have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides can also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

**[0234]** In certain embodiments, the polynucleotides comprise the coding sequence for the mature polypeptide fused in the same reading frame to a marker sequence that allows, for example, for purification of the encoded polypeptide. For example, the marker sequence can be a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or the marker sequence can be a hemagglutinin (HA) tag derived from the influenza hemagglutinin protein when a mammalian host (e.g., COS-7 cells) is used. In some embodiments, the marker sequence is a FLAG-tag, a peptide of sequence DYKDDDDK (SEQ ID NO:33) which can be used in conjunction with other affinity tags.

**[0235]** The present invention further relates to variants of the hereinabove described polynucleotides encoding, for example, fragments, analogs, and/or derivatives.

**[0236]** In certain embodiments, the present invention provides polynucleotides comprising polynucleotides having a nucleotide sequence at least about 80% identical, at least about 85% identical, at least about 90% identical, at least about 95% identical, and in some embodiments, at least about 96%, 97%, 98% or 99% identical to a polynucleotide encoding a polypeptide comprising a RSPO3-binding agent (e.g., an antibody), or fragment thereof, described herein.

**[0237]** As used herein, the phrase a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence is intended to mean that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence can include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence can be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence can be inserted into the reference sequence. These mutations of the reference sequence can occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

**[0238]** The polynucleotide variants can contain alterations in the coding regions, non-coding regions, or both. In some embodiments, a polynucleotide variant contains alterations which produce silent substitutions, additions, or deletions, but does not alter the properties or activities of the encoded

polypeptide. In some embodiments, a polynucleotide variant comprises silent substitutions that result in no change to the amino acid sequence of the polypeptide (due to the degeneracy of the genetic code). In some embodiments, nucleotide variants comprise nucleotide sequences which result in expression differences (e.g., increased or decreased expression) at the transcript level. Polynucleotide variants can be produced for a variety of reasons, for example, to optimize codon expression for a particular host (i.e., change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*). In some embodiments, a polynucleotide variant comprises at least one silent mutation in a non-coding or a coding region of the sequence.

[0239] In some embodiments, a polynucleotide variant is produced to modulate or alter expression (or expression levels) of the encoded polypeptide. In some embodiments, a polynucleotide variant is produced to increase expression of the encoded polypeptide. In some embodiments, a polynucleotide variant is produced to decrease expression of the encoded polypeptide. In some embodiments, a polynucleotide variant has increased expression of the encoded polypeptide as compared to a parental polynucleotide sequence. In some embodiments, a polynucleotide variant has decreased expression of the encoded polypeptide as compared to a parental polynucleotide sequence.

[0240] In some embodiments, at least one polynucleotide variant is produced (without changing the amino acid sequence of the encoded polypeptide) to increase production of a heteromultimeric molecule. In some embodiments, at least one polynucleotide variant is produced (without changing the amino acid sequence of the encoded polypeptide) to increase production of a bispecific antibody.

[0241] In certain embodiments, the polynucleotides are isolated. In certain embodiments, the polynucleotides are substantially pure.

[0242] Vectors comprising the polynucleotides described herein are also provided. Cells comprising the polynucleotides described herein are also provided. In some embodiments, an expression vector comprises a polynucleotide molecule. In some embodiments, a host cell comprises an expression vector comprising the polynucleotide molecule. In some embodiments, a host cell comprises a polynucleotide molecule.

#### IV. Methods of use and pharmaceutical compositions

[0243] The RSPO3-binding agents (including polypeptides and antibodies) of the invention are useful in a variety of applications including, but not limited to, therapeutic treatment methods, such as the treatment of cancer. In certain embodiments, the agents are useful for inhibiting  $\beta$ -catenin signaling, inhibiting tumor growth, modulating angiogenesis, inhibiting angiogenesis, inducing differentiation, reducing tumor volume, reducing the frequency of cancer stem cells in a tumor, and/or reducing the tumorigenicity of a tumor. The methods of use may be *in vitro*, *ex vivo*, or *in vivo* methods. In certain embodiments, a RSPO3-binding agent or polypeptide or antibody is an antagonist of human RSPO3.

[0244] In certain embodiments, the RSPO3-binding agents are used in the treatment of a disease associated with activation of  $\beta$ -catenin, increased  $\beta$ -catenin signaling, and/or aberrant  $\beta$ -catenin signaling. In certain embodiments, the disease is a disease dependent upon  $\beta$ -catenin signaling. In certain embodiments, the RSPO3-binding agents are used in the treatment of disorders characterized by increased angiogenesis. In certain embodiments, the RSPO3-binding agents are used in the treatment of disorders characterized by increased levels of stem cells and/or progenitor cells. In some embodiments, the methods comprise administering a therapeutically effective amount of a RSPO3-binding agent (e.g., antibody) to a subject. In some embodiments, the subject is human.

[0245] The present invention provides methods for inhibiting growth of a tumor using the RSPO3-binding agents or antibodies described herein. In certain embodiments, the method of inhibiting growth of a tumor comprises contacting a cell with a RSPO3-binding agent (e.g., an antibody) *in vitro*. For example, an immortalized cell line or a cancer cell line is cultured in medium to which is added an anti-RSPO3 antibody or other agent to inhibit tumor growth. In some embodiments, tumor cells are isolated from a patient sample such as, for example, a tissue biopsy, pleural effusion, or blood sample and cultured in medium to which is added a RSPO3-binding agent to inhibit tumor growth.

[0246] In some embodiments, the method of inhibiting growth of a tumor comprises contacting the tumor or tumor cells with a RSPO3-binding agent (e.g., an antibody) *in vivo*. In certain embodiments, contacting a tumor or tumor cell with a RSPO3-binding agent is undertaken in an animal model. For example, a RSPO3-binding agent may be administered to immunocompromised mice (e.g. NOD/SCID mice) which have xenografts. In some embodiments, cancer cells or cancer stem cells are isolated from a patient sample such as, for example, a tissue biopsy, pleural effusion, or blood sample and injected into immunocompromised mice that are then administered a RSPO3-binding agent to inhibit tumor cell growth. In some embodiments, a RSPO3-binding agent is administered to the animal. In some embodiments, the RSPO3-binding agent is administered at the same time or shortly after introduction of tumorigenic cells into the animal to prevent tumor growth ("preventative model"). In some embodiments, the RSPO3-binding agent is administered as a therapeutic after tumors have grown to a specified size ("therapeutic model"). In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is an anti-RSPO3 antibody. In some embodiments, the anti-RSPO3 antibody is antibody 131R002. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or antibody h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is

antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

[0247] In certain embodiments, the method of inhibiting growth of a tumor comprises administering to a subject a therapeutically effective amount of a RSPO3-binding agent, wherein the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments of the method, the RSPO3-binding agent further comprises a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14) or QQSNEPLTF (SEQ ID NO:83). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLTF (SEQ ID NO:83). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ



ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14).

[0248] In certain embodiments, the method of inhibiting growth of a tumor comprises administering to a subject a therapeutically effective amount of a RSPO3-binding agent. In certain embodiments, the subject is a human. In certain embodiments, the subject has a tumor or has had a tumor which was removed. In some embodiments, the subject has a tumor with an elevated expression level of at least one RSPO protein (e.g., RSPO1, RSPO2, RSPO3, or RSPO4). In some embodiments, the subject has a tumor with a high expression level of at least one RSPO protein (e.g., RSPO1, RSPO2, RSPO3, or RSPO4). In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R01.

[0249] In certain embodiments, the tumor is a tumor in which  $\beta$ -catenin signaling is active. In some embodiments, the tumor is a tumor in which  $\beta$ -catenin signaling is aberrant. In certain embodiments, the tumor comprises an inactivating mutation (e.g., a truncating mutation) in the APC tumor suppressor gene. In certain embodiments, the tumor does not comprise an inactivating mutation in the APC tumor suppressor gene. In some embodiments, the tumor comprises a wild-type APC gene. In some embodiments, the tumor does not comprise an activating mutation in the  $\beta$ -catenin gene. In certain embodiments, a cancer for which a subject is being treated involves such a tumor.

[0250] In some embodiments, the tumor comprises a RSPO gene fusion. In some embodiments, the tumor comprises a RSPO2 gene fusion. In some embodiments, the tumor comprises a RSPO3 gene fusion.

[0251] In certain embodiments, the tumor expresses RSPO3 to which a RSPO3-binding agent or antibody binds. In certain embodiments, the tumor has elevated expression levels of RSPO1 or over-expresses RSPO1. In certain embodiments, the tumor has elevated expression levels of RSPO2 or over-

expresses RSPO2. In certain embodiments, the tumor has elevated expression levels of RSPO3 or over-expresses RSPO3. The phrase “a tumor has elevated expression levels of” may refer to expression levels of a protein or expression levels of a nucleic acid. In general, the phrase “a tumor has elevated expression levels of” a protein or a gene (or similar phrases) refers to expression levels of a protein or a gene in a tumor as compared to expression levels of the same protein or the same gene in a reference sample or to a pre-determined expression level. In some embodiments, the reference sample is normal tissue of the same tissue type. In some embodiments, the reference sample is normal tissue of a group of tissue types. In some embodiments, the reference sample is a tumor or group of tumors of the same tissue type. In some embodiments, the reference sample is a tumor or group of tumors of a different tissue type. Thus in some embodiments, the expression levels of a protein or a gene in a tumor are “elevated” or “high” as compared to the average expression level of the protein or the gene within a group of tissue types. In some embodiments, the expression levels of a protein or a gene in a tumor are “elevated” or “high” as compared to the expression level of the protein or the gene in other tumors of the same tissue type or a different tissue type. In some embodiments, the tumor expresses “elevated” or “high” levels of RSPO1, RSPO2, RSPO3, and/or RSPO4 as compared to the RSPO levels expressed in normal tissue of the same tissue type. In some embodiments, the tumor expresses “elevated” or “high” levels of RSPO1, RSPO2, RSPO3, and/or RSPO4 as compared to a predetermined level.

[0252] In addition, the invention provides a method of inhibiting growth of a tumor in a subject, comprising administering a therapeutically effective amount of a RSPO3-binding agent to the subject. In certain embodiments, the tumor comprises cancer stem cells. In certain embodiments, the frequency of cancer stem cells in the tumor is reduced by administration of the RSPO3-binding agent. The invention also provides a method of reducing the frequency of cancer stem cells in a tumor, comprising contacting the tumor with an effective amount of a RSPO3-binding agent (e.g., an anti-RSPO3 antibody). In some embodiments, a method of reducing the frequency of cancer stem cells in a tumor in a subject, comprising administering to the subject a therapeutically effective amount of a RSPO3-binding agent (e.g., an anti-RSPO3 antibody) is provided. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is an anti-RSPO3 antibody. In some embodiments, the anti-RSPO3 antibody is 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or antibody h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

[0253] In some embodiments, the tumor is a solid tumor. In certain embodiments, the tumor is a tumor selected from the group consisting of colorectal tumor, pancreatic tumor, lung tumor, ovarian tumor, liver tumor, breast tumor, kidney tumor, prostate tumor, gastrointestinal tumor, melanoma, cervical tumor, bladder tumor, glioblastoma, and head and neck tumor. As used herein, "lung cancer" includes but is not limited to, small cell lung carcinoma and non-small cell lung carcinoma (NSCLC). In certain embodiments, the tumor is a colorectal tumor. In certain embodiments, the tumor is an ovarian tumor. In some embodiments, the tumor is a lung tumor. In certain embodiments, the tumor is a pancreatic tumor. In some embodiments, the tumor is a colorectal tumor that comprises an inactivating mutation in the APC gene. In some embodiments, the tumor is a colorectal tumor that does not comprise an inactivating mutation in the APC gene. In some embodiments, the tumor is a colorectal tumor that contains a RSPO gene fusion. In some embodiments, the tumor is a colorectal tumor that contains a RSPO2 gene fusion. In some embodiments, the tumor is a colorectal tumor that contains a RSPO3 gene fusion. In some embodiments, the tumor is an ovarian tumor with an elevated expression level of RSPO1. In some embodiments, the tumor is a pancreatic tumor with an elevated expression level of RSPO2. In some embodiments, the tumor is a colon tumor with an elevated expression level of RSPO2. In some embodiments, the tumor is a lung tumor with an elevated expression level of RSPO2. In some embodiments, the tumor is a lung tumor with an elevated expression level of RSPO3. In some embodiments, the tumor is an ovarian tumor with an elevated expression level of RSPO3. In some embodiments, the tumor is a breast tumor with an elevated expression level of RSPO3. In some embodiments, the tumor is a colorectal tumor with an elevated expression level of RSPO3.

[0254] The present invention further provides methods for treating cancer comprising administering a therapeutically effective amount of a RSPO3-binding agent to a subject. In certain embodiments, the cancer is characterized by cells expressing elevated levels of at least one RSPO protein as compared to expression levels of the same RSPO protein in a reference sample. As used herein, a "reference sample" includes but is not limited to, normal tissue, non-cancerous tissue of the same tissue type, tumor tissue of the same tissue type, and tumor tissue of a different tissue type. In certain embodiments, the cancer is characterized by cells expressing elevated levels of at least one RSPO protein as compared to a pre-determined level of the same RSPO protein. In some embodiments, determining the expression level of at least one RSPO is done prior to treatment. In some embodiments, determining the expression level of at least one RSPO is by immunohistochemistry. Thus, in certain embodiments, the cancer is characterized by cells expressing elevated levels of at least one RSPO protein as compared to expression levels of the same RSPO protein in normal tissue. In certain embodiments, the cancer is characterized by cells over-expressing RSPO1. In certain embodiments, the cancer is characterized by cells over-expressing RSPO2. In certain embodiments, the cancer is characterized by cells over-expressing RSPO3. In certain embodiments, the cancer over-expresses at least one RSPO protein selected from the group consisting of

RSPO1, RSPO2, RSPO3, and/or RSPO4. In certain embodiments, the cancer is characterized by cells expressing  $\beta$ -catenin, wherein the RSPO3-binding agent (e.g., an antibody) interferes with RSPO3-induced  $\beta$ -catenin signaling and/or activation.

[0255] In some embodiments, the RSPO-binding agent binds RSPO3, and inhibits or reduces growth of the cancer. In some embodiments, the RSPO-binding agent binds RSPO3, interferes with RSPO3/LGR interactions, and inhibits or reduces growth of the cancer. In some embodiments, the RSPO-binding agent binds RSPO3, inhibits  $\beta$ -catenin activation, and inhibits or reduces growth of the cancer. In some embodiments, the RSPO-binding agent binds RSPO3, and reduces the frequency of cancer stem cells in the cancer. In some embodiments, the RSPO-binding agent is an antibody. In some embodiments, the RSPO-binding agent is an anti-RSPO3 antibody. In some embodiments, the anti-RSPO3 antibody is antibody 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or antibody h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

[0256] The present invention provides for methods of treating cancer comprising administering a therapeutically effective amount of a RSPO3-binding agent to a subject (e.g., a subject in need of treatment). In certain embodiments, the method of treating cancer comprises administering to a subject a therapeutically effective amount of a RSPO3-binding agent, wherein the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80). In some embodiments of the method, the RSPO3-binding agent further comprises a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12) or KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14) or QQSNEDPLTF (SEQ ID NO:83). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising

QQSNEDPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANNFDY (SEQ ID NO:35), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLTF (SEQ ID NO:83). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising KASQSVDDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In some embodiments, the RSPO3-binding agent comprises a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80), and/or a light chain CDR1 comprising QSVDDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEDPLT (SEQ ID NO:14). In certain embodiments, the subject is a human. In certain embodiments, the subject has a cancerous tumor. In certain embodiments, the subject has had a tumor removed. In some embodiments, a method of treating cancer comprises administering a therapeutically effective amount of a RSPO3-binding agent to a subject, wherein the subject has a tumor that has elevated expression of at least one RSPO protein as compared to a reference sample or a pre-determined level. In some embodiments, the subject has a lung tumor that has elevated expression of RSPO3 and is administered an anti-RSPO3 antibody.

[0257] The invention also provides a RSPO3-binding agent for use in a method of treating cancer, wherein the RSPO3-binding agent is an antibody described herein. The invention also provides the use of an RSPO3-binding agent (e.g., an antibody) described herein for the manufacture of a medicament for the treatment of cancer.

[0258] In certain embodiments, the cancer is a cancer selected from the group consisting of colorectal cancer, pancreatic cancer, lung cancer, ovarian cancer, liver cancer, breast cancer, kidney cancer, prostate cancer, gastrointestinal cancer, melanoma, cervical cancer, bladder cancer, glioblastoma, and head and neck cancer. In certain embodiments, the cancer is pancreatic cancer. In certain embodiments, the cancer is ovarian cancer. In certain embodiments, the cancer is colorectal cancer. In certain embodiments, the cancer is breast cancer. In certain embodiments, the cancer is prostate cancer. In certain embodiments, the cancer is lung cancer.

[0259] In addition, the invention provides a method of reducing the tumorigenicity of a tumor in a subject, comprising administering to a subject a therapeutically effective amount of a RSPO3-binding agent. In certain embodiments, the tumor comprises cancer stem cells. In some embodiments, the tumorigenicity of a tumor is reduced by reducing the frequency of cancer stem cells in the tumor. In some embodiments, the methods comprise using the RSPO3-binding agents described herein. In certain embodiments, the frequency of cancer stem cells in the tumor is reduced by administration of a RSPO3-binding agent.

[0260] In certain embodiments, the methods further comprise a step of determining the expression level of at least one RSPO (i.e., protein or nucleic acid) in the tumor or cancer. In some embodiments, the step of determining the expression level of a RSPO in the tumor or cancer comprises determining the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4. In some embodiments, the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a tumor or cancer is compared to the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a reference sample. In some embodiments, the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a tumor or cancer is compared to the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4 in normal tissue. In some embodiments, the level of expression of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a tumor or cancer is compared to a pre-determined level of expression of RSPO1, RSPO2, RSPO3, and/or RSPO4. In some embodiments, the level of expression of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a tumor or cancer is compared to a pre-determined level of expression of RSPO1, RSPO2, RSPO3, and/or RSPO4 in normal tissue. In some embodiments, the tumor has a high expression level of RSPO1. In some embodiments, the tumor has a high expression level of RSPO3. In general, the expression level of a RSPO (i.e., protein or nucleic acid) is compared to the expression level of the RSPO (i.e., protein or nucleic acid) in normal tissue of the same tissue type. However, in some embodiments, the expression level of a RSPO (i.e., protein or nucleic acid) is compared to the average expression level of the RSPO (i.e., protein or nucleic acid) within a group of tissue types. In some embodiments, the expression levels of a RSPO (i.e., protein or nucleic acid) in a tumor is compared to the expression level of the RSPO (i.e., protein or nucleic acid) in other tumors of the same tissue type or a different tissue type.

[0261] In some embodiments, determining the level of RSPO expression is done prior to treatment. In some embodiments, the subject is administered a RSPO3-binding agent or antibody describe herein if the tumor or cancer has an elevated expression level of RSPO as compared to the expression level of the same RSPO in a reference sample (e.g., normal tissue) or a pre-determined level. For example, in some embodiments, the subject is administered a RSPO3-binding agent (e.g., anti-RSPO3 antibody) if the tumor or cancer has an elevated expression level of RSPO3 (i.e., protein or nucleic acid) as compared to the expression level of RSPO3 in normal or control tissue.

[0262] In certain embodiments, the methods further comprise a step of determining if the tumor or cancer has an inactivating mutation in the APC gene. In some embodiments, the methods further comprise a step of determining if the tumor or cancer has an activating mutation in the  $\beta$ -catenin gene. In some embodiments, the methods further comprise a step of determining if the tumor or cancer has a RSPO gene fusion.

[0263] In addition, the invention provides a method of modulating angiogenesis, comprising administering to a subject a therapeutically effective amount of a RSPO3-binding agent. In some embodiments, the modulating angiogenesis comprises inhibiting angiogenesis. In some embodiments, the methods comprise using the RSPO3-binding agents described herein. In certain embodiments, the RSPO3-binding agent binds RSPO3 and inhibits or reduces angiogenesis. In certain embodiments, the inhibition and/or reduction of angiogenesis inhibits or reduces growth of a tumor or cancer. In some embodiments, the RSPO3-binding agent binds RSPO3 and promotes aberrant angiogenesis. In some embodiments, the RSPO3-binding agent binds RSPO3 and promotes unproductive angiogenesis. In certain embodiments, the aberrant angiogenesis or the unproductive angiogenesis inhibits or reduces growth of a tumor or cancer.

[0264] In addition, the present invention provides methods of identifying a human subject for treatment with a RSPO-binding agent, comprising determining if the subject has a tumor that has an elevated expression level of RSPO (i.e., protein or nucleic acid) as compared to expression of the same RSPO (i.e., protein or nucleic acid) in normal tissue, in a reference sample, or to a pre-determined level of the RSPO protein. In some embodiments, a method of identifying a human subject for treatment with a RSPO3-binding agent comprises determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3. In some embodiments, a method of identifying a human subject for treatment with a RSPO3-binding agent comprises: obtaining a tumor sample from the subject, and determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3. In some embodiments, if the tumor has an elevated expression level of RSPO3, the subject is selected for treatment with an antibody that specifically binds RSPO3. In some embodiments, if selected for treatment, the subject is administered a RSPO3-binding agent or antibody describe herein. In some embodiments, if the tumor has

an elevated expression level of more than one RSPO (i.e., protein or nucleic acid), the subject is administered a RSPO-binding agent that binds the RSPO with the highest level of expression. In certain embodiments, the subject has had a tumor removed. For example, in some embodiments, the expression level of RSPO1, RSPO2, RSPO3, and/or RSPO4 in a tumor is determined, if the tumor has an elevated level of RSPO3 expression as compared to the level of RSPO3 in normal tissue, the subject is selected for treatment with an antibody that specifically binds RSPO3. If selected for treatment, the subject is administered an anti-RSPO3 antibody describe herein. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized form of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized form of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R011.

**[0265]** The present invention provides methods of selecting a human subject for treatment with a RSPO-binding agent, comprising determining if the subject has a tumor that has an elevated expression level of at least one RSPO (i.e., protein or nucleic acid), as compared to expression of the same RSPO in normal tissue or as compared to a predetermined level, wherein if the tumor has an elevated expression level of at least one RSPO, the subject is selected for treatment with an antibody that specifically binds the RSPO with the elevated expression level. In some embodiments, if selected for treatment, the subject is administered a RSPO-binding agent or antibody describe herein. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3 comprises: determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3 comprises obtaining a tumor sample from the subject, and determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3, wherein if the tumor has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises obtaining a tumor sample from the subject, and determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3, wherein if the tumor



has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody. In certain embodiments, the subject has had a tumor removed. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an anti-RSPO3 antibody. In some embodiments, the anti-RSPO3 antibody is antibody 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

**[0266]** The present invention also provides methods of treating cancer in a human subject, comprising: (a) selecting a subject for treatment based, at least in part, on the subject having a cancer that has an elevated level of a RSPO, and (b) administering to the subject a therapeutically effective amount of a RSPO3-binding agent described herein. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

**[0267]** Methods for determining the level of RSPO expression in a cell, tumor or cancer are known by those of skill in the art. For nucleic acid expression these methods include, but are not limited to, PCR-based assays, microarray analyses and nucleotide sequencing (e.g., NextGen sequencing). For protein expression these methods include, but are not limited to, Western blot analysis, protein arrays, ELISAs, immunohistochemistry (IHC) assays, and FACS.

**[0268]** The present invention provides methods of identifying a human subject for treatment with a RSPO3-binding agent, comprising obtaining a tumor sample from the subject, and determining if the tumor has a RSPO gene fusion. In some embodiments, a method of identifying a human subject for treatment with a RSPO3-binding agent comprises: determining if the subject has a tumor that has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with

the antibody. In some embodiments, a method of identifying a human subject for treatment with a RSPO3-binding agent comprises: (a) obtaining a tumor sample from the subject, and (b) determining if the tumor has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with the antibody. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises determining if the subject has a tumor that has a RSPO gene fusion.

[0269] The present invention also provides methods of selecting a human subject for treatment with a RSPO-binding agent, comprising determining if the subject has a tumor that has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion, the subject is selected for treatment with an antibody that specifically binds a RSPO protein. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3 comprises determining if the subject has a tumor that has a RSPO gene fusion. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises obtaining a tumor sample from the subject, and determining if the tumor has a RSPO gene fusion. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises determining if the subject has a tumor that has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion the subject is selected for treatment with the antibody. In some embodiments, a method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprises obtaining a tumor sample from the subject, and determining if the tumor has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion the subject is selected for treatment with the antibody. In some embodiments, the RSPO gene fusion is a RSPO2 gene fusion. In some embodiments, the RSPO gene fusion is a RSPO3 gene fusion. In some embodiments, if selected for treatment, the subject is administered a RSPO-binding agent or antibody describe herein. In certain embodiments, the subject has had a tumor removed. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an anti-RSPO3 antibody. In some embodiments, the anti-RSPO3 antibody is antibody 131R002. In some embodiments, the anti-RSPO3 antibody is antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

**[0270]** The present invention also provides methods of treating cancer in a human subject, comprising: (a) selecting a subject for treatment based, at least in part, on the subject having a cancer that has a RSPO gene fusion, and (b) administering to the subject a therapeutically effective amount of a RSPO3-binding agent described herein. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R002. In some embodiments, the anti-RSPO3 antibody is a humanized version of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is a humanized version of a variant of antibody 131R003. In some embodiments, the anti-RSPO3 antibody is antibody h131R006A or h131R006B. In some embodiments, the anti-RSPO3 antibody is antibody h131R005/131R007. In some embodiments, the anti-RSPO3 antibody is antibody h131R008. In some embodiments, the anti-RSPO3 antibody is antibody h131R010. In some embodiments, the anti-RSPO3 antibody is antibody h131R011.

**[0271]** Methods for determining whether a tumor has a RSPO gene fusion are known by those of skill in the art. Methods may include but are not limited to, PCR-based assays, microarray analyses, and nucleotide sequencing (e.g., NextGen sequencing, whole-genome sequencing (WGS)).

**[0272]** Methods for determining whether a tumor or cancer has an elevated level of RSPO expression or has a RSPO gene fusion can use a variety of samples. In some embodiments, the sample is taken from a subject having a tumor or cancer. In some embodiments, the sample is a fresh tumor/cancer sample. In some embodiments, the sample is a frozen tumor/cancer sample. In some embodiments, the sample is a formalin-fixed paraffin-embedded sample. In some embodiments, the sample is processed to a cell lysate. In some embodiments, the sample is processed to DNA or RNA.

**[0273]** Methods of treating a disease or disorder in a subject, wherein the disease or disorder is associated with aberrant (e.g., increased levels)  $\beta$ -catenin signaling are further provided. Methods of treating a disease or disorder in a subject, wherein the disease or disorder is characterized by an increased level of stem cells and/or progenitor cells are further provided. In some embodiments, the treatment methods comprise administering a therapeutically effective amount of a RSPO-binding agent, polypeptide, or antibody to the subject. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R002. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the

RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R011.

[0274] The invention also provides a method of inhibiting  $\beta$ -catenin signaling in a cell comprising contacting the cell with an effective amount of a RSPO-binding agent. In certain embodiments, the cell is a tumor cell. In certain embodiments, the method is an *in vivo* method wherein the step of contacting the cell with the RSPO3-binding agent comprises administering a therapeutically effective amount of the RSPO3-binding agent to the subject. In some embodiments, the method is an *in vitro* or *ex vivo* method. In certain embodiments, the RSPO-binding agent inhibits  $\beta$ -catenin signaling. In some embodiments, the RSPO-binding agent inhibits activation of  $\beta$ -catenin. In certain embodiments, the RSPO-binding agent interferes with a RSPO/LGR interaction. In certain embodiments, the LGR is LGR4, LGR5, and/or LGR6. In certain embodiments, the LGR is LGR4. In certain embodiments, the LGR is LGR5. In certain embodiments, the LGR is LGR6. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R002. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R011.

[0275] The use of the RSPO-binding agents, polypeptides, or antibodies described herein to induce the differentiation of cells, including, but not limited to tumor cells, is also provided. In some embodiments, methods of inducing cells to differentiate comprise contacting the cells with an effective amount of a RSPO-binding agent (e.g., an anti-RSPO antibody) described herein. In certain embodiments, methods of inducing cells in a tumor in a subject to differentiate comprise administering a therapeutically effective amount of a RSPO-binding agent, polypeptide, or antibody to the subject. In some embodiments, methods for inducing differentiation markers on tumor cells comprise administering a therapeutically effective amount of a RSPO-binding agent, polypeptide, or antibody. In some embodiments, the tumor is a solid tumor. In some embodiments, the tumor is selected from the group consisting of colorectal tumor, pancreatic tumor, lung tumor, ovarian tumor, liver tumor, breast tumor, kidney tumor, prostate tumor, gastrointestinal tumor, melanoma, cervical tumor, bladder tumor, glioblastoma, and head and neck tumor. In certain embodiments, the tumor is an ovarian tumor. In certain other embodiments, the tumor is a

colon tumor. In some embodiments, the tumor is a lung tumor. In certain embodiments, the method is an *in vivo* method. In certain embodiments, the method is an *in vitro* method. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R002. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R011.

[0276] The invention further provides methods of differentiating tumorigenic cells into non-tumorigenic cells comprising contacting the tumorigenic cells with a RSPO-binding agent. In some embodiments, the method comprises administering the RSPO-binding agent to a subject that has a tumor comprising tumorigenic cells or that has had such a tumor removed. In certain embodiments, the tumorigenic cells are ovarian tumor cells. In certain embodiments, the tumorigenic cells are colon tumor cells. In some embodiments, the tumorigenic cells are lung tumor cells. In some embodiments, the RSPO-binding agent is a RSPO3-binding agent. In some embodiments, the RSPO3-binding agent is an antibody. In some embodiments, the RSPO3-binding agent is antibody 131R002. In some embodiments, the RSPO3-binding agent is antibody 131R003. In some embodiments, the RSPO3-binding agent is a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R002. In some embodiments, the RSPO3-binding agent is a humanized version of antibody 131R003. In some embodiments, the RSPO3-binding agent is a humanized version of a variant of antibody 131R003. In some embodiments, the RSPO3-binding agent is antibody h131R006A or antibody h131R006B. In some embodiments, the RSPO3-binding agent is antibody h131R005/131R007. In some embodiments, the RSPO3-binding agent is antibody h131R008. In some embodiments, the RSPO3-binding agent is antibody h131R010. In some embodiments, the RSPO3-binding agent is antibody h131R011.

[0277] In certain embodiments, the disease treated with the RSPO3-binding agents described herein is not a cancer. For example, the disease may be a metabolic disorder such as obesity or diabetes (e.g., type II diabetes) (Jin T., 2008, *Diabetologia*, 51:1771-80). Alternatively, the disease may be a bone disorder such as osteoporosis, osteoarthritis, or rheumatoid arthritis (Corr M., 2008, *Nat. Clin. Pract. Rheumatol.*, 4:550-6; Day et al., 2008, *Bone Joint Surg. Am.*, 90 Suppl 1:19-24). The disease may also be a kidney

disorder, such as a polycystic kidney disease (Harris et al., 2009, *Ann. Rev. Med.*, 60:321-337; Schmidt-Ott et al., 2008, *Kidney Int.*, 74:1004-8; Benzing et al., 2007, *J. Am. Soc. Nephrol.*, 18:1389-98).

Alternatively, eye disorders including, but not limited to, macular degeneration and familial exudative vitreoretinopathy may be treated (Lad et al., 2009, *Stem Cells Dev.*, 18:7-16). Cardiovascular disorders, including myocardial infarction, atherosclerosis, and valve disorders, may also be treated (Al-Aly Z., 2008, *Transl. Res.*, 151:233-9; Kobayashi et al., 2009, *Nat. Cell Biol.*, 11:46-55; van Gijn et al., 2002, *Cardiovasc. Res.*, 55:16-24; Christman et al., 2008, *Am. J. Physiol. Heart Circ. Physiol.*, 294:H2864-70).

In some embodiments, the disease is a pulmonary disorder such as idiopathic pulmonary arterial hypertension or pulmonary fibrosis (Laumanns et al., 2008, *Am. J. Respir. Cell Mol. Biol.*, 2009, 40:683-691; Königshoff et al., 2008, *PLoS ONE*, 3:e2142). In some embodiments, the disease treated with the RSPO3-binding agent is a liver disease, such as cirrhosis or liver fibrosis (Cheng et al., 2008, *Am. J. Physiol. Gastrointest. Liver Physiol.*, 294:G39-49).

[0278] The present invention further provides pharmaceutical compositions comprising the RSPO3-binding agents described herein. In certain embodiments, the pharmaceutical compositions further comprise a pharmaceutically acceptable vehicle. In some embodiments, these pharmaceutical compositions find use in inhibiting tumor growth and treating cancer in a subject (e.g., a human patient).

[0279] In certain embodiments, formulations are prepared for storage and use by combining a purified antibody or agent of the present invention with a pharmaceutically acceptable vehicle (e.g., a carrier or excipient). Suitable pharmaceutically acceptable vehicles include, but are not limited to, nontoxic buffers such as phosphate, citrate, and other organic acids; salts such as sodium chloride; antioxidants including ascorbic acid and methionine; preservatives such as octadecyldimethylbenzyl ammonium chloride, hexamethonium chloride, benzalkonium chloride, benzethonium chloride, phenol, butyl or benzyl alcohol, alkyl parabens, such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, 3-pentanol, and m-cresol; low molecular weight polypeptides (e.g., less than about 10 amino acid residues); proteins such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; carbohydrates such as monosaccharides, disaccharides, glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes such as Zn-protein complexes; and non-ionic surfactants such as TWEEN or polyethylene glycol (PEG). (*Remington: The Science and Practice of Pharmacy*, 22<sup>nd</sup> Edition, 2012, Pharmaceutical Press, London.)

[0280] The pharmaceutical compositions of the present invention can be administered in any number of ways for either local or systemic treatment. Administration can be topical by epidermal or transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders; pulmonary by inhalation or insufflation of powders or aerosols, including by nebulizer, intratracheal, and intranasal;

oral; or parenteral including intravenous, intraarterial, intratumoral, subcutaneous, intraperitoneal, intramuscular (e.g., injection or infusion), or intracranial (e.g., intrathecal or intraventricular).

[0281] The therapeutic formulation can be in unit dosage form. Such formulations include tablets, pills, capsules, powders, granules, solutions or suspensions in water or non-aqueous media, or suppositories. In solid compositions such as tablets the principal active ingredient is mixed with a pharmaceutical carrier. Conventional tableting ingredients include corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and diluents (e.g., water). These can be used to form a solid pre-formulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. The solid pre-formulation composition is then subdivided into unit dosage forms of a type described above. The tablets, pills, etc. of the formulation or composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner composition covered by an outer component. Furthermore, the two components can be separated by an enteric layer that serves to resist disintegration and permits the inner component to pass intact through the stomach or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials include a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

[0282] The RSPO3-binding agents or antibodies described herein can also be entrapped in microcapsules. Such microcapsules are prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules) or in macroemulsions as described in *Remington: The Science and Practice of Pharmacy*, 22<sup>nd</sup> Edition, 2012, Pharmaceutical Press, London.

[0283] In certain embodiments, pharmaceutical formulations include a RSPO3-binding agent (e.g., an antibody) of the present invention complexed with liposomes. Methods to produce liposomes are known to those of skill in the art. For example, some liposomes can be generated by reverse phase evaporation with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes can be extruded through filters of defined pore size to yield liposomes with the desired diameter.

[0284] In certain embodiments, sustained-release preparations can be produced. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing a RSPO3-binding agent (e.g., an antibody), where the matrices are in the form of shaped articles (e.g., films or microcapsules). Examples of sustained-release matrices include polyesters, hydrogels such as poly(2-hydroxyethyl-methacrylate) or poly(vinyl alcohol), polylactides, copolymers of L-glutamic acid

and 7 ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), sucrose acetate isobutyrate, and poly-D-(-)-3-hydroxybutyric acid.

**[0285]** In certain embodiments, in addition to administering a RSPO3-binding agent (e.g., an antibody), the method or treatment further comprises administering at least one additional therapeutic agent. An additional therapeutic agent can be administered prior to, concurrently with, and/or subsequently to, administration of the RSPO3-binding agent. Pharmaceutical compositions comprising a RSPO3-binding agent and the additional therapeutic agent(s) are also provided. In some embodiments, the at least one additional therapeutic agent comprises 1, 2, 3, or more additional therapeutic agents.

**[0286]** Combination therapy with two or more therapeutic agents often uses agents that work by different mechanisms of action, although this is not required. Combination therapy using agents with different mechanisms of action may result in additive or synergetic effects. Combination therapy may allow for a lower dose of each agent than is used in monotherapy, thereby reducing toxic side effects and/or increasing the therapeutic index of the agent(s). Combination therapy may decrease the likelihood that resistant cancer cells will develop. In some embodiments, combination therapy comprises a therapeutic agent that affects (e.g., inhibits or kills) non-tumorigenic cells and a therapeutic agent that affects (e.g., inhibits or kills) tumorigenic CSCs.

**[0287]** In some embodiments, the combination of a RSPO3-binding agent and at least one additional therapeutic agent results in additive or synergistic results. In some embodiments, the combination therapy results in an increase in the therapeutic index of the RSPO3-binding agent. In some embodiments, the combination therapy results in an increase in the therapeutic index of the additional agent(s). In some embodiments, the combination therapy results in a decrease in the toxicity and/or side effects of the RSPO3-binding agent. In some embodiments, the combination therapy results in a decrease in the toxicity and/or side effects of the additional agent(s).

**[0288]** Useful classes of therapeutic agents include, for example, antitubulin agents, auristatins, DNA minor groove binders, DNA replication inhibitors, alkylating agents (e.g., platinum complexes such as cisplatin, mono(platinum), bis(platinum) and tri-nuclear platinum complexes and carboplatin), anthracyclines, antibiotics, antifolates, antimetabolites, chemotherapy sensitizers, duocarmycins, etoposides, fluorinated pyrimidines, ionophores, lexitropsins, nitrosoureas, platinols, purine antimetabolites, puromycins, radiation sensitizers, steroids, taxanes, topoisomerase inhibitors, vinca alkaloids, or the like. In certain embodiments, the second therapeutic agent is an alkylating agent, an antimetabolite, an antimetabolic, a topoisomerase inhibitor, or an angiogenesis inhibitor. In some embodiments, the second therapeutic agent is a platinum complex such as carboplatin or cisplatin. In some embodiments, the additional therapeutic agent is a platinum complex in combination with a taxane.



[0289] Therapeutic agents that may be administered in combination with the RSPO3-binding agents include chemotherapeutic agents. Thus, in some embodiments, the method or treatment involves the administration of a RSPO3-binding agent or antibody of the present invention in combination with a chemotherapeutic agent or cocktail of multiple different chemotherapeutic agents. Treatment with a RSPO3-binding agent (e.g., an antibody) can occur prior to, concurrently with, or subsequent to administration of chemotherapies. Combined administration can include co-administration, either in a single pharmaceutical formulation or using separate formulations, or consecutive administration in either order but generally within a time period such that all active agents can exert their biological activities simultaneously. Preparation and dosing schedules for such chemotherapeutic agents can be used according to manufacturers' instructions or as determined empirically by the skilled practitioner.

Preparation and dosing schedules for such chemotherapy are also described in *The Chemotherapy Source Book, 4<sup>th</sup> Edition*, 2008, M. C. Perry, Editor, Lippincott, Williams & Wilkins, Philadelphia, PA.

[0290] Chemotherapeutic agents useful in the instant invention include, but are not limited to, alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, caminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytosine arabinoside, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenishers such as folinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK; razoxane;

sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside (Ara-C); taxoids, e.g. paclitaxel (TAXOL) and docetaxel (TAXOTERE); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; ibandronate; CPT11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine (XELODA); and pharmaceutically acceptable salts, acids or derivatives of any of the above. Chemotherapeutic agents also include anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (FARESTON); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above. In certain embodiments, the additional therapeutic agent is cisplatin. In certain embodiments, the additional therapeutic agent is carboplatin. In certain embodiments, the additional therapeutic agent is paclitaxel (taxol). In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, 131R006A, 131R006B, 131R005/131R007, or 131R008 in combination with cisplatin.

**[0291]** In certain embodiments, the chemotherapeutic agent is a topoisomerase inhibitor. Topoisomerase inhibitors are chemotherapy agents that interfere with the action of a topoisomerase enzyme (e.g., topoisomerase I or II). Topoisomerase inhibitors include, but are not limited to, doxorubicin HCl, daunorubicin citrate, mitoxantrone HCl, actinomycin D, etoposide, topotecan HCl, teniposide (VM-26), and irinotecan, as well as pharmaceutically acceptable salts, acids, or derivatives of any of these. In some embodiments, the additional therapeutic agent is irinotecan. Thus, in some embodiments, a method comprises administering a RSPO3-binding agent in combination with a topoisomerase inhibitor. In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with irinotecan.

**[0292]** In certain embodiments, the chemotherapeutic agent is an anti-metabolite. An anti-metabolite is a chemical with a structure that is similar to a metabolite required for normal biochemical reactions, yet different enough to interfere with one or more normal functions of cells, such as cell division. Anti-metabolites include, but are not limited to, gemcitabine, fluorouracil, capecitabine, methotrexate sodium, ralitrexed, pemetrexed, tegafur, cytosine arabinoside, thioguanine, 5-azacytidine, 6-mercaptopurine, azathioprine, 6-thioguanine, pentostatin, fludarabine phosphate, and cladribine, as well as pharmaceutically acceptable salts, acids, or derivatives of any of these. In certain embodiments, the additional therapeutic agent is gemcitabine. Thus, in some embodiments, a method comprises

administering a RSPO3-binding agent in combination with an anti-metabolite. In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with gemcitabine. In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with pemetrexed.

**[0293]** In certain embodiments, the chemotherapeutic agent is an antimitotic agent, including, but not limited to, agents that bind tubulin. In some embodiments, the agent is a taxane. In certain embodiments, the agent is paclitaxel or docetaxel, or a pharmaceutically acceptable salt, acid, or derivative of paclitaxel or docetaxel. In certain embodiments, the agent is paclitaxel (TAXOL), docetaxel (TAXOTERE), albumin-bound paclitaxel (nab-paclitaxel; ABRAXANE), DHA-paclitaxel, or PG-paclitaxel. In certain alternative embodiments, the antimitotic agent comprises a vinca alkaloid, such as vincristine, binblastine, vinorelbine, or vindesine, or pharmaceutically acceptable salts, acids, or derivatives thereof. In some embodiments, the antimitotic agent is an inhibitor of kinesin Eg5 or an inhibitor of a mitotic kinase such as Aurora A or Plk1. In certain embodiments, where the chemotherapeutic agent administered in combination with a RSPO-binding agent is an anti-mitotic agent, the cancer or tumor being treated is breast cancer or a breast tumor. In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with paclitaxel. In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with nab-paclitaxel (ABRAXANE). In some embodiments, a method comprises administering anti-RSPO3 antibody 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with gemcitabine and nab-paclitaxel (ABRAXANE).

**[0294]** In some embodiments, an additional therapeutic agent comprises an agent such as a small molecule. For example, treatment can involve the combined administration of a RSPO3-binding agent (e.g. an antibody) of the present invention with a small molecule that acts as an inhibitor against additional tumor-associated antigens including, but not limited to, EGFR, ErbB2, HER2, and/or VEGF. In certain embodiments, the additional therapeutic agent is a small molecule that inhibits a cancer stem cell pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the Notch pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the Wnt pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the BMP pathway. In some embodiments, the additional therapeutic agent is a molecule that inhibits  $\beta$ -catenin signaling.

**[0295]** In some embodiments, an additional therapeutic agent comprises a biological molecule, such as an antibody. For example, treatment can involve the combined administration of a RSPO3-binding agent (e.g. an antibody) of the present invention with other antibodies against additional tumor-associated antigens including, but not limited to, antibodies that bind EGFR, ErbB2, HER2, and/or VEGF. In some embodiments, the additional therapeutic agent is an antibody that binds a second RSPO, e.g., RSPO1, RSPO2, and/or RSPO4. In some embodiments, the additional therapeutic agent is an anti-RSPO2 antibody. In some embodiments, the additional therapeutic agent is an anti-RSPO1 antibody. In certain embodiments, the additional therapeutic agent is an antibody specific for an anti-cancer stem cell marker. In some embodiments, the additional therapeutic agent is an antibody that binds a component of the Notch pathway. In some embodiments, the additional therapeutic agent is an antibody that binds a component of the Wnt pathway. In certain embodiments, the additional therapeutic agent is an antibody that inhibits a cancer stem cell pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the Notch pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the Wnt pathway. In some embodiments, the additional therapeutic agent is an inhibitor of the BMP pathway. In some embodiments, the additional therapeutic agent is an antibody that inhibits  $\beta$ -catenin signaling. In certain embodiments, the additional therapeutic agent is an antibody that is an angiogenesis inhibitor (e.g., an anti-VEGF or VEGF receptor antibody). In certain embodiments, the additional therapeutic agent is bevacizumab (AVASTIN), trastuzumab (HERCEPTIN), panitumumab (VECTIBIX), or cetuximab (ERBITUX).

**[0296]** In some embodiments, the methods described herein comprise administering a therapeutically effective amount of a RSPO3-binding agent in combination with Wnt pathway inhibitors. In some embodiments, the Wnt pathway inhibitors are frizzled (FZD) protein binding agents, "FZD-binding agents". Non-limiting examples of FZD-binding agents can be found in U.S. Patent No. 7,982,013, which is incorporated by reference herein in its entirety. FZD-binding agents may include, but are not limited to, anti-FZD antibodies. In some embodiments, a method comprises administering a RSPO-binding agent in combination with an anti-FZD antibody. In some embodiments, a method comprises administering a RSPO-binding agent in combination with the anti-FZD antibody 18R5. In some embodiments, the Wnt pathway inhibitors are Wnt protein binding agents, "Wnt-binding agents". Nonlimiting examples of Wnt-binding agents can be found in U.S. Patent Nos. 7,723,477 and 7,947,277; and International Publications WO 2011/088127 and WO 2011/088123, which are incorporated by reference herein in their entirety. Wnt-binding agents may include, but are not limited to, anti-Wnt antibodies and FZD-Fc soluble receptors. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with a FZD-Fc soluble receptor. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with a FZD8-Fc soluble receptor. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with an anti-FZD antibody. In some

embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with an anti-FZD antibody. In some embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with anti-FZD antibody 18R5. In some embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with a FZD-Fc soluble receptor. In some embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with a FZD8-Fc soluble receptor.

[0297] In some embodiments, the methods described herein comprise administering a therapeutically effective amount of a RSPO-binding agent in combination with more than one additional therapeutic agent. Thus, in some embodiments, a method comprises administering a RSPO-binding agent in combination with a chemotherapeutic agent and a Wnt pathway inhibitor. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with a chemotherapeutic agent and a Wnt pathway inhibitor. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with a chemotherapeutic agent and anti-FZD antibody 18R5. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with a chemotherapeutic agent and a FZD8-Fc soluble receptor. In some embodiments, a method comprises administering a RSPO3-binding agent in combination with gemcitabine and a Wnt pathway inhibitor. In some embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with gemcitabine and anti-FZD antibody 18R5. In some embodiments, a method comprises administering anti-RSPO3 antibodies 131R002, 131R003, a variant of 131R003, a humanized version of 131R003, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, or h131R011 in combination with gemcitabine and FZD8-Fc soluble receptor.

[0298] Furthermore, treatment with a RSPO3-binding agent described herein can include combination treatment with other biologic molecules, such as one or more cytokines (e.g., lymphokines, interleukins, tumor necrosis factors, and/or growth factors) or can be accompanied by surgical removal of tumors, cancer cells or any other therapy deemed necessary by a treating physician.

[0299] In certain embodiments, the treatment involves the administration of a RSPO3-binding agent (e.g. an antibody) of the present invention in combination with radiation therapy. Treatment with a RSPO3-

binding agent can occur prior to, concurrently with, or subsequent to administration of radiation therapy. Dosing schedules for such radiation therapy can be determined by the skilled medical practitioner.

**[0300]** Combined administration can include co-administration, either in a single pharmaceutical formulation or using separate formulations, or consecutive administration in either order but generally within a time period such that all active agents can exert their biological activities simultaneously.

**[0301]** It will be appreciated that the combination of a RSPO3-binding agent and at least one additional therapeutic agent may be administered in any order or concurrently. In some embodiments, the RSPO3-binding agent will be administered to patients that have previously undergone treatment with a second therapeutic agent. In certain other embodiments, the RSPO3-binding agent and a second therapeutic agent will be administered substantially simultaneously or concurrently. For example, a subject may be given a RSPO3-binding agent (e.g., an antibody) while undergoing a course of treatment with a second therapeutic agent (e.g., chemotherapy). In certain embodiments, a RSPO3-binding agent will be administered within 1 year of the treatment with a second therapeutic agent. In certain alternative embodiments, a RSPO3-binding agent will be administered within 10, 8, 6, 4, or 2 months of any treatment with a second therapeutic agent. In certain other embodiments, a RSPO3-binding agent will be administered within 4, 3, 2, or 1 weeks of any treatment with a second therapeutic agent. In some embodiments, a RSPO3-binding agent will be administered within 5, 4, 3, 2, or 1 days of any treatment with a second therapeutic agent. It will further be appreciated that the two (or more) agents or treatments may be administered to the subject within a matter of hours or minutes (i.e., substantially simultaneously).

**[0302]** For the treatment of a disease, the appropriate dosage of an RSPO3-binding agent (e.g., an antibody) of the present invention depends on the type of disease to be treated, the severity and course of the disease, the responsiveness of the disease, whether the RSPO3-binding agent or antibody is administered for therapeutic or preventative purposes, previous therapy, the patient's clinical history, and so on, all at the discretion of the treating physician. The RSPO3-binding agent or antibody can be administered one time or over a series of treatments lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved (e.g., reduction in tumor size). Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient and will vary depending on the relative potency of an individual antibody or agent. The administering physician can easily determine optimum dosages, dosing methodologies, and repetition rates. In certain embodiments, dosage is from 0.01  $\mu$ g to 100mg/kg of body weight, from 0.1  $\mu$ g to 100mg/kg of body weight, from 1  $\mu$ g to 100mg/kg of body weight, from 1mg to 100mg/kg of body weight, 1mg to 80mg/kg of body weight from 10mg to 100mg/kg of body weight, from 10mg to 75mg/kg of body weight, or from 10mg to 50mg/kg of body weight. In certain embodiments, the dosage of the antibody or other RSPO3-binding agent is from about 0.1mg to about 20mg/kg of body weight. In certain embodiments, dosage can

be given once or more daily, weekly, monthly, or yearly. In certain embodiments, the antibody or other RSPO3-binding agent is given once every week, once every two weeks or once every three weeks.

**[0303]** In some embodiments, a RSPO3-binding agent (e.g., an antibody) may be administered at an initial higher “loading” dose, followed by one or more lower doses. In some embodiments, the frequency of administration may also change. In some embodiments, a dosing regimen may comprise administering an initial dose, followed by additional doses (or “maintenance” doses) once a week, once every two weeks, once every three weeks, or once every month. For example, a dosing regimen may comprise administering an initial loading dose, followed by a weekly maintenance dose of, for example, one-half of the initial dose. Or a dosing regimen may comprise administering an initial loading dose, followed by maintenance doses of, for example one-half of the initial dose every other week. Or a dosing regimen may comprise administering three initial doses for 3 weeks, followed by maintenance doses of, for example, the same amount every other week.

**[0304]** As is known to those of skill in the art, administration of any therapeutic agent may lead to side effects and/or toxicities. In some cases, the side effects and/or toxicities are so severe as to preclude administration of the particular agent at a therapeutically effective dose. In some cases, drug therapy must be discontinued, and other agents may be tried. However, many agents in the same therapeutic class often display similar side effects and/or toxicities, meaning that the patient either has to stop therapy, or if possible, suffer from the unpleasant side effects associated with the therapeutic agent.

**[0305]** Thus, the present invention provides methods of treating cancer in a subject comprising using an intermittent dosing strategy for administering one or more agents, which may reduce side effects and/or toxicities associated with administration of a RSPO3-binding agent, chemotherapeutic agent, etc. In some embodiments, a method for treating cancer in a human subject comprises administering to the subject a therapeutically effective dose of a RSPO3-binding agent in combination with a therapeutically effective dose of a chemotherapeutic agent, wherein one or both of the agents are administered according to an intermittent dosing strategy. In some embodiments, the intermittent dosing strategy comprises administering an initial dose of a RSPO3-binding agent to the subject, and administering subsequent doses of the RSPO3-binding agent about once every 2 weeks. In some embodiments, the intermittent dosing strategy comprises administering an initial dose of a RSPO3-binding agent to the subject, and administering subsequent doses of the RSPO3-binding agent about once every 3 weeks. In some embodiments, the intermittent dosing strategy comprises administering an initial dose of a RSPO3-binding agent to the subject, and administering subsequent doses of the RSPO3-binding agent about once every 4 weeks. In some embodiments, the RSPO3-binding agent is administered using an intermittent dosing strategy and the chemotherapeutic agent is administered weekly.

## V. Kits comprising RSPO-binding agents

[0306] The present invention provides kits that comprise the RSPO3-binding agents (e.g., antibodies) described herein and that can be used to perform the methods described herein. In certain embodiments, a kit comprises at least one purified antibody against at least one human RSPO protein in one or more containers. In some embodiments, the kits contain all of the components necessary and/or sufficient to perform a detection assay, including all controls, directions for performing assays, and any necessary software for analysis and presentation of results. One skilled in the art will readily recognize that the disclosed RSPO3-binding agents of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

[0307] Further provided are kits comprising a RSPO3-binding agent (e.g., an anti-RSPO3 antibody), as well as at least one additional therapeutic agent. In certain embodiments, the second (or more) therapeutic agent is a chemotherapeutic agent. In certain embodiments, the second (or more) therapeutic agent is a Wnt pathway inhibitor. In certain embodiments, the second (or more) therapeutic agent is an angiogenesis inhibitor.

[0308] Embodiments of the present disclosure can be further defined by reference to the following non-limiting examples, which describe in detail preparation of certain antibodies of the present disclosure and methods for using antibodies of the present disclosure. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the scope of the present disclosure.

## EXAMPLES

### Example 1

#### Expression of RSPO and LGR in human tumors

[0309] mRNA from normal tissue, benign tumor and malignant tumor samples of a large number of human patients was analyzed by microarray analysis (Genologic BioExpress Datasuite). This data revealed elevated expression levels of RSPO1 in malignant tissue relative to normal tissue in several tumor types including kidney, endometrial, and ovarian. RSPO1 was noted to be frequently over-expressed in ovarian cancer (Fig. 1A). In addition, this data suggested elevated expression levels of RSPO3 in malignant tissue relative to normal tissue in several tumor types including ovarian, pancreas, and lung (Fig. 1C). In addition, it was found that LGR5 and LGR6 were over-expressed in malignant breast tumors, colon tumors, lung tumors, and ovarian tumors relative to normal tissue, while LGR4 was over-expressed in lung tumors. LGR5 and LGR6 over-expression appeared to be restricted to triple-negative (ER<sup>neg</sup>PR<sup>neg</sup>HER2<sup>neg</sup>) breast tumors relative to other breast tumor subtypes.



[0310] RNA was isolated from a series of human tumors grown in murine xenografts. The RNA samples were prepared and processed using established Affymetrix protocols for the generation of labeled cRNA. The processed RNA was hybridized to Affymetrix HG-U133 plus 2.0 microarrays (Affymetrix, Santa Clara, CA) as outlined in the manufacturer's technical manuals. After hybridization, the microarrays were washed, scanned, and analyzed. Scanned array background adjustment and signal intensity normalization were performed using the GCRMA algorithm (Bioconductor, [www.bioconductor.org](http://www.bioconductor.org)).

[0311] Particular human RSPOs and human LGRs were evaluated -- RSPO1 (241450\_at), RSPO2 (1554012\_at), RSPO3 (228186\_s\_at), RSPO4 (237423\_at), LGR4 (218326\_s\_at), LGR5 (210393\_at) and LGR6 (227819\_at). Microarray analysis showed that, while LGR4 and LGR6 were broadly expressed in almost all tumors, many tumors were found to greatly over-express only particular RSPO family members and LGR5 (Table 2), although these expression levels were not compared to expression levels in normal tissue. Generally there is only a single RSPO family member that is highly expressed in a given tumor, suggesting that there may be functional redundancy within the RSPO family.

Table 2

Tumor	RSPO1	RSPO2	RSPO3	RSPO4	LGR4	LGR5	LGR6
Breast tumor							
B34	4.79	4.93	<b>303.31</b>	4.41			
B39	20.59	<b>588.88</b>	22.60	4.40			
B60	4.60	4.92	10.89	<b>64.79</b>			
B02	4.60	4.92	<b>692.34</b>	4.41	2678.95	4.28	50.88
B03	5.56	4.89	<b>1870.42</b>	4.41	686.47	<b>30.78</b>	73.49
B06	4.60	4.91	4.51	<b>120.72</b>	274.54	4.26	20.77
B59	4.60	4.91	4.53	<b>1158.11</b>	200.48	4.26	6467.15
Colon tumors							
C11	4.63	4.98	4.56	4.43	3852.26	6.22	11.31
C17	4.64	5.00	4.57	4.44	2822.46	<b>62.34</b>	43.94
C18	4.63	4.95	<b>13.83</b>	4.42	2454.15	4.29	723.15
C27	6.66	<b>980.49</b>	4.75	4.40	5083.84	4.30	20.82
Lung tumors							
LU02	4.62	<b>15190.40</b>	4.55	4.43	13.95	4.29	14.56
LU11	4.60	4.92	<b>4.53</b>	4.41	999.55	4.27	146.67
LU25	4.64	5.56	<b>11123.06</b>	4.44	1208.92	4.29	41089
LU33	4.64	5.01	12.02	<b>62.98</b>	329.62	4.30	20.96
LU45	4.64	4.99	4.62	4.44	3877.47	4.29	4.86
Melanoma tumors							

M06	4.73	<b>21.80</b>	4.65	4.50	1077.93	4.34	3.90
Ovarian tumors							
OV12	4.72	5.12	4.64	<b>460.40</b>	5383.63	<b>1152.73</b>	115.04
OV19	<b>960.19</b>	4.74	69.77	20.90	494.67	5.72	4302.78
OV22	4.66	5.10	<b>132.85</b>	37.43	3743.91	<b>482.33</b>	812.05
OV27	4.55	4.86	<b>125.78</b>	4.92			
OV38	9.19	4.83	<b>3439.88</b>	16.35	1528.12	4.24	19.49
Pancreatic tumors							
PN07	4.58	<b>689.52</b>	4.51	4.40	6777.41	4.28	746.38
PN18	4.72	<b>2508.47</b>	4.65	4.50	6750.73	<b>51.15</b>	564.94

## Example 2

### Binding of RSPO proteins to LGR5

[0312] A cell surface LGR5 protein was generated by ligating amino acids 22-564 of human LGR5 to an N-terminal FLAG tag and to the transmembrane domain of CD4 and a C-terminal GFP protein tag using standard recombinant DNA techniques (FLAG-LGR5-CD4TM-GFP). RSPO-Fc constructs were generated using standard recombinant DNA techniques. Specifically, full-length human RSPO1, RSPO2, RSPO3 and RSPO4 were ligated in-frame to a human Fc region and the recombinant RSPO-Fc proteins were expressed in insect cells using baculovirus. The fusion proteins were purified from the insect medium using protein A chromatography.

[0313] HEK-293 cells were transiently transfected with the FLAG-LGR5-CD4TM-GFP construct. After 48 hours, transfected cells were suspended in ice cold PBS containing 2% FBS and heparin and incubated on ice in the presence of 10µg/ml RSPO1-Fc, RSPO2-Fc, RSPO3-Fc, RSPO4-Fc, or FZD8-Fc fusion proteins for 15 minutes. A second incubation with 100µl PE-conjugated anti-human Fc secondary antibody was performed to detect cells bound by the Fc fusion proteins. Cells were incubated with an anti-FLAG antibody (Sigma-Aldrich, St. Louis, MO) as a positive control and with an anti-PE antibody as a negative control. The cells were analyzed on a FACSCalibur instrument (BD Biosciences, San Jose, CA) and the data was processed using FlowJo software.

[0314] As shown in Figure 2, RSPO1, RSPO2, RSPO3 and RSPO4 all bound to LGR5 expressed on the surface of the HEK-293 cells, while FZD8, the negative control, did not bind LGR5.

[0315] Binding affinities between RSPO proteins and LGR5 were analyzed by surface plasmon resonance. A soluble LGR5-Fc construct was generated using standard recombinant DNA techniques. Specifically, amino acids 1-564 of human LGR5 were ligated in frame to human Fc and the recombinant LGR5-Fc fusion protein was expressed in insect cells using baculovirus. The LGR5-Fc fusion protein was purified from the insect medium using protein A chromatography. Cleavage of the LGR5 signal

sequence results in a mature LGR5-Fc fusion protein containing amino acids 22-564 of LGR5. Recombinant RSPO1-Fc, RSPO2-Fc, RSPO3-Fc and RSPO4-Fc fusion proteins were immobilized on CM5 chips using standard amine-based chemistry (NHS/EDC). Two-fold dilutions of soluble LGR5-Fc were injected over the chip surface (100nM to 0.78nM). Kinetic data were collected over time using a Biacore 2000 system from Biacore Life Sciences (GE Healthcare) and the data were fit using the simultaneous global fit equation to yield affinity constants ( $K_D$  values) for each RSPO protein (Table 3).

Table 3

	LGR5 (nM)
RSPO1	110
RSPO2	14
RSPO3	<1.0
RSPO4	73

**[0316]** Human RSPO1, RSPO2, RSPO3 and RSPO4 all bound to LGR5, demonstrating that RSPO proteins may be ligands for LGR proteins.

### Example 3

#### Identification of anti-RSPO3 antibodies

**[0317]** A mammalian cell antibody library was screened and two anti-RSPO3 antibodies, 131R002 and 131R003, were identified. Sequence data subsequently demonstrated that antibodies 131R002 and 131R003 have the same light chain sequence but different heavy chain sequences.

**[0318]** The  $K_{DS}$  of antibodies 131R002 and 131R003 were determined using a Biacore 2000 system from Biacore LifeSciences (GE Healthcare). Recombinant human RSPO3 protein was biotinylated and captured on streptavidin-coated chips (GE Healthcare) with coating densities of 400-700ru. The antibodies were serially diluted 2-fold from 100nM to 0.78nM in HBS-P (0.01M HEPES pH 7.4, 0.15M NaCl, 0.005% v/v Surfactant P20) and were injected over the chip surface. Kinetic data were collected over time and were fit using the simultaneous global fit equation to yield affinity constants ( $K_D$  values) for each antibody.

**[0319]** Antibody 131R002 had an affinity constant ( $K_D$ ) for human RSPO3 of 8.2nM and antibody 131R003 had a  $K_D$  for human RSPO3 of 7.3nM.

### Example 4

#### *In vitro* testing for inhibition of $\beta$ -catenin activity by anti-RSPO3 antibodies

[0320] HEK-293 cells were transfected with a 6xTCF-luciferase reporter vector (TOPflash, Millipore, Billerica, MA). After 24-48 hrs, the transfected HEK-293 cells were incubated with a combination of WNT3a (5ng/ml) and human RSPO3 (10ng/ml, R&D BioSystems, Minneapolis, MN) in the presence of anti-RSPO3 antibodies 131R002 and 131R003. Antibodies 131R002 and 131R003 were added to the cells in 4-fold serial dilutions from 20µg/ml to 0.02µg/ml. As controls, cells were incubated with a combination of WNT3a and RSPO3, WNT3a only, RSPO3 only, or with no addition. The cells were incubated for 16 hours and luciferase activity was measured using Steady-Glo® Luciferase Assay System according to the manufacturer's instructions (Promega, Madison, WI).

[0321] As shown in Figure 3, anti-RSPO3 antibodies 131R002 and 131R003 each reduced RSPO3-induced  $\beta$ -catenin signaling in a dose-dependent manner. These results demonstrated that antibodies 131R002 and 131R003 are specific inhibitors of RSPO3 and are capable of reducing and/or blocking RSPO3-induced  $\beta$ -catenin signaling.

#### Example 5

##### Affinity maturation and humanization of RSPO3 antibodies

[0322] Anti-RSPO3 antibody 131R003 was affinity matured and several variants were identified. One 131R003 variant had an altered heavy chain CDR1 (SEQ ID NO:34) as compared to parental 131R003 antibody. A second variant had an altered heavy chain CDR3 (SEQ ID NO:35) as compared to parental 131R003. An additional variant was generated that comprised both the altered heavy chain CDR1 and CDR3 as compared to parental 131R003.

[0323] HEK-293 cells were transfected with a 6xTCF-luciferase reporter vector (TOPflash, Millipore, Billerica, MA). After 24-48 hrs, the transfected HEK-293 cells were incubated with a combination of WNT3a and human RSPO3 in the presence of anti-RSPO3 antibodies 131R003, 131R003 CDR1 variant and 131R003 CDR3 variant. 131R003, 131R003 CDR1 variant, and 131R003 CDR3 variant were added to the cells in 5-fold serial dilutions from 20µg/ml to 0.006µg/ml. As controls, cells were incubated with a combination of WNT3a and RSPO3, WNT3a only, RSPO3 only, a control antibody, or with no addition. The cells were incubated for 16 hours and luciferase activity was measured using Steady-Glo® Luciferase Assay System according to the manufacturer's instructions (Promega, Madison, WI).

[0324] As shown in Figure 4, anti-RSPO3 antibodies 131R003 CDR1 variant and 131R003 CDR3 variant each reduced RSPO3-induced  $\beta$ -catenin signaling in a dose-dependent manner and at lower concentrations than parental 131R003. These results demonstrated that the 131R003 variants retained the characteristics of parental 131R003, i.e., they were specific inhibitors of RSPO3 and were capable of reducing and/or blocking RSPO3-induced  $\beta$ -catenin signaling. In addition, these results demonstrated that the 131R003 variants had better activity than parental 131R003.

[0325] Humanized forms of 131R003 variants were generated using standard techniques. Humanized antibodies h131R005, h131R007, h131R008, h131R010, h131R011 comprise an altered heavy chain CDR3 as compared to parental 131R003 antibody. Humanized 131R006B comprises an altered heavy chain CDR3 as compared to parental 131R003 antibody. Antibodies h131R005/131R007, h131R010, and h131R011 comprise several amino acid substitutions in framework region 3 as compared to antibody 131R006B. Antibodies h131R005/131R007, h131R006, and h131R011 are IgG2 antibodies. Antibodies h131R008 and h131R010 are IgG1 antibodies. Antibodies h131R005/131R007, h131R010, and h131R011 comprise the same heavy chain variable region. Antibodies h131R010 and h131R011 comprise the same light chain variable region, which is different than the light chain variable region of h131R005/131R007.

[0326] A plasmid encoding the heavy chain of the 131R010 antibody was deposited with American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA, USA, under the conditions of the Budapest Treaty on June 18, 2013, and assigned ATCC deposit designation number PTA-\_\_\_\_\_. A plasmid encoding the light chain of the 131R010 antibody was deposited with ATCC, 10801 University Boulevard, Manassas, VA, USA, under the conditions of the Budapest Treaty on June 18, 2013, and assigned ATCC deposit designation number PTA-\_\_\_\_\_.

#### Example 6

Inhibition of ovarian tumor growth *in vivo* by anti-RSPO antibodies

[0327] Dissociated OMP-OV38 ovarian tumor cells ( $1 \times 10^5$  cells) were injected in to 6-8 week old NOD/SCID mice. Tumors were allowed to grow for 39 days until they reached an average volume of 150 mm<sup>3</sup>. The mice were randomized (n = 8 per group) and treated with a combination of anti-RSPO1 antibody 89M5 and anti-RSPO3 antibody 131R003, a combination of anti-RSPO1 antibody 89M5, anti-RSPO3 antibody 131R003, and taxol, taxol as a single agent, or a control antibody. Antibodies were dosed at 20 mg/kg once a week, and taxol was dosed at 15 mg/ml once a week. Administration of the antibodies and taxol was performed via injection into the intraperitoneal cavity. Tumor growth was monitored and tumor volumes were measured with electronic calipers at the indicated time points. Data are expressed as mean  $\pm$  S.E.M.

[0328] As shown in Figure 5, a combination of anti-RSPO1 and anti-RSPO3 antibodies inhibited OMP-OV38 ovarian tumor growth. Surprisingly, a combination of anti-RSPO1 antibody 89M5, anti-RSPO3 antibody 131R002, and taxol inhibited tumor growth to a significantly greater level than taxol alone or the antibody combination alone.

[0329] Dissociated OMP-OV38 ovarian tumor cells ( $1 \times 10^5$  cells) were injected in to 6-8 week old NOD/SCID mice. Tumors were allowed to grow for 35 days until they reached an average volume of 140 mm<sup>3</sup>. The mice were randomized (n = 10 per group) and treated with anti-RSPO3 antibody 131R002,

anti-RSPO1 antibody 89M5, taxol, a combination of 89M5 and taxol, a combination of 131R002 and taxol, a combination of 89M5 and 131R002, a combination of 89M5, 131R002 and taxol, or a control antibody. Antibodies were dosed at 20 mg/kg once a week, and taxol was dosed at 15 mg/ml once a week through day 46 and subsequently dosed at 7.5 mg/kg. Administration of the antibodies and taxol was performed via injection into the intraperitoneal cavity. Tumor growth was monitored and tumor volumes were measured with electronic calipers at the indicated time points. Data are expressed as mean  $\pm$  S.E.M.

[0330] As shown in Figure 6, a combination of anti-RSPO1 antibody 89M5 and anti-RSPO3 antibody 131R002 inhibited OMP-OV38 ovarian tumor growth as compared to control antibody. Combinations of anti-RSPO1 antibody 89M5 and taxol or anti-RSPO3 antibody 131R002 and taxol had no effect relative to taxol alone. However, surprisingly a combination of anti-RSPO1 89M5, anti-RSPO3 antibody 131R002, and taxol showed activity that was greater than taxol alone.

#### Example 7

Inhibition of lung tumor growth *in vivo* by anti-RSPO3 antibodies

[0331] In OMP-LU45 non-small cell lung tumors, it has been observed that CD201<sup>+</sup> cells are more tumorigenic than CD201<sup>-</sup> cells. Furthermore, RSPO3 was found to be highly expressed in the CD201<sup>+</sup> cell population. Dissociated and sorted OMP-LU45 CD44<sup>+</sup>CD201<sup>+</sup> lung tumor cells ( $5 \times 10^4$  cells) were injected into 6-8 week old NOD/SCID mice. Tumors were allowed to grow for 38 days until they reached an average volume of 140 mm<sup>3</sup>. The mice were randomized (n = 10 per group) and treated with anti-RSPO3 antibody 131R002 or a control antibody. Antibodies were dosed at 25 mg/kg once a week and administration of the antibodies was performed via injection into the intraperitoneal cavity. Tumor growth was monitored and tumor volumes were measured with electronic calipers at the indicated time points. Data are expressed as mean  $\pm$  S.E.M.

[0332] In a study with a second lung tumor, dissociated OMP-LU25 lung tumor cells ( $5 \times 10^4$  cells) were injected into 6-8 week old NOD/SCID mice. Tumors were allowed to grow for 48 days until they reached an average volume of 110 mm<sup>3</sup>. The mice were randomized (n = 9 per group) and treated with anti-RSPO3 antibody 131R002 or a control antibody. Antibodies were dosed at 25 mg/kg once a week and administration of the antibodies was performed via injection into the intraperitoneal cavity. Tumor growth was monitored and tumor volumes were measured with electronic calipers at the indicated time points. Data are expressed as mean  $\pm$  S.E.M.

[0333] As shown in Figure 7A and 7B, anti-RSPO antibody 131R002 inhibited growth of both lung tumors OMP-LU45 and OMP-LU25 as compared to a control antibody.

#### Example 8

Inhibition of  $\beta$ -catenin activity by anti-RSPO3 antibodies

[0334] HEK-293 cells were transfected with a 6xTCF-luciferase reporter vector (TOPflash, Millipore, Billerica, MA). After 24-48 hrs, the transfected HEK-293 cells were incubated with a combination of WNT3a conditioned medium (5ng/ml) and human RSPO3 (10ng/ml, R&D BioSystems) in the presence of anti-RSPO3 antibodies 131R002, 131R006B, or 131R007. Antibodies 131R002, 131R006 or 131R007 were added to the cells in 5-fold serial dilutions from 20 $\mu$ g/ml to 0.0064 $\mu$ g/ml. As controls, cells were incubated with WNT3a conditioned medium alone, a combination of WNT3a conditioned medium and human RSPO3, or with no addition to cells. The cells were incubated for 16 hours and luciferase activity was measured using Steady-Glo® Luciferase Assay System according to the manufacturer's instructions (Promega, Madison, WI).

[0335] As shown in Figure 8, all three anti-RSPO3 antibodies reduced WNT3a/RSPO3-induced  $\beta$ -catenin signaling in a dose-dependent manner. The humanized antibodies 131R006B and 131R007 appeared to have a greater ability to inhibit  $\beta$ -catenin activity than antibody 131R002. These results demonstrated that humanized antibodies 131R006B and 131R007 are stronger inhibitors of RSPO3 than 131R002 and are capable of reducing and/or blocking WNT3a/RSPO3-induced  $\beta$ -catenin signaling.

#### Example 9

##### Inhibition of RSPO3 binding to LGR5

[0336] HEK-293T cells were transfected with a cDNA expression vector that encoded the extracellular domain of human LGR5 (FLAG-LGR5-CD4TM-GFP). Transfected cells were incubated with recombinant RSPO3-biotin fusion protein in the presence of anti-RSPO3 antibodies 131R006B or 131R007. Cells were incubated without antibody as a control. Cells were washed in PBS and binding of RSPO3 to LGR5-expressing transfected cells was determined by addition of PE-conjugated streptavidin and analysis by flow cytometry.

[0337] As shown in Figure 9, anti-RSPO3 antibodies 131R006B and 131R007 were highly effective in blocking binding of RSPO3 to LGR5-expressing cells.

#### Example 10

##### Binding affinities of RSPO3 antibodies

[0338] The  $K_D$  of RSPO3 antibodies 131R002, 131R003, 131R003 CDR3 variant, h131R007, h131R008, and h131R011 were determined using a Biacore 2000 system from Biacore LifeSciences (GE Healthcare). The method used was different than described in Example 3. A goat anti-human IgG antibody was coupled to a carboxymethyl-dextran (CM5) SPR chip using standard amine-based chemistry (NHS/EDC) and blocked with ethanolamine. Antibodies (purified antibody or culture supernatant) were diluted to a concentration of 10 $\mu$ g/ml in HBS-P-BSA (0.01M HEPES pH7.4, 0.15M NaCl, 0.005% v/v Polysorbate 20, 100ug/ml BSA) and captured onto the chip via the anti-human IgG antibody. Human RSPO3 (R&D

Systems) was serially diluted 2-fold from 300nM to 37.5nM in HBS-P-BSA and injected sequentially over the captured anti-RSPO3 antibodies. RSPO3 association and dissociation was measured at each concentration. After each antigen injection 5 $\mu$ l of 100mM H<sub>3</sub>PO<sub>4</sub> was injected to remove the antigen-antibody complex and a subsequent injection performed. Kinetic data were collected over time and were fit using the simultaneous global fit equation to yield affinity constants ( $K_D$  values) for each antibody (Table 4).

Table 4

RSPO3 Antibody	$K_D$
131R002 (IgG2)	1.3nM
131R003 (IgG2)	1.9nM
131R003 CDR3 variant (IgG2)	1.7nM
h131R007 (IgG2)	654pM
h131R008 (IgG1)	876pM
h131R010 (IgG1)	ND
h131R011 (IgG2)	686pM

[0339] In additional experiments, antibody h131R008 was shown to have a  $K_D$  as low as 448pM for human RSPO3, no detectable binding to human RSPO1 or RSPO2, and weak binding to human RSPO4. Antibody h131R008 was shown to have a  $K_D$  of 248pM for murine RSPO3, no detectable binding to murine RSPO1 or RSPO2 and weak binding to murine RSPO4.

#### Example 11

Inhibition of lung tumor growth *in vivo* by anti-RSPO3 antibodies

[0340] The non-small cell lung cancer (NSCLC) cell line NCI-H2030 was selected for testing based on a high level of RPSO3 expression in microarray data. NCI-H2030 cells ( $1 \times 10^6$ ) were injected into NOD-SCID mice. Tumors were allowed to grow for approximately 60 days until they reached an average volume of 100 mm<sup>3</sup>. Tumor-bearing mice were randomized into 4 groups (n = 7-9 per group). Tumor-bearing mice were treated with anti-RSPO3 antibody 131R002, carboplatin, a combination of anti-RSPO3 antibody 131R002 and carboplatin, or a control antibody. Antibodies were dosed at 25 mg/kg once a week. Carboplatin was dosed at 50 mg/kg once a week. Tumor growth was monitored and tumor volumes were measured with electronic calipers on the indicated days post-treatment.

[0341] As shown in Figure 10, treatment with anti-RSPO3 antibody in combination with carboplatin inhibited NCI-H2030 tumor growth better than carboplatin alone or the antibody alone.

[0342] OMP-LU102 is a patient-derived non-small cell lung cancer (NSCLC) xenograft that was selected for testing based on a high level of RPSO3 expression in microarray data. OMP-LU102 lung tumor cells



( $1 \times 10^5$ ) were injected into NOD-SCID mice. Tumors were allowed to grow for 22 days until they reached an average volume of  $90 \text{ mm}^3$ . Tumor-bearing mice were randomized into 4 groups ( $n = 10$  per group). Tumor-bearing mice were treated with anti-RSPO3 antibody 131R002, carboplatin, a combination of anti-RSPO3 antibody 131R002 and carboplatin, or a control antibody. Antibodies were dosed at 25 mg/kg once a week. Carboplatin was dosed at 50 mg/kg once a week. Tumor growth was monitored and tumor volumes were measured with electronic calipers on the indicated days post-treatment.

[0343] As shown in Figure 11A, treatment with anti-RSPO3 antibody inhibited OMP-LU102 lung tumor growth as a single agent but had much greater effect in combination with carboplatin.

[0344] RNA was prepared from tumors from each of the four experimental groups following the treatment. Gene expression was characterized by microarray analysis. Gene set enrichment analysis indicated that anti-RSPO3 antibody treatment (either as a single agent or in combination with carboplatin) inhibited the expression of various gene sets characteristic of normal stem cells or cancer stem cells as shown in Figure 11B. Treatment with carboplatin alone did not have this effect on gene expression.

#### Example 12

Inhibition of pancreatic tumor growth *in vivo* by anti-RSPO3 antibodies

[0345] OMP-PN35 is patient-derived pancreatic ductal adenocarcinoma (PDAC) xenograft that was selected for testing based on high level of RPSO3 expression in microarray data. OMP-PN35 ( $1 \times 10^5$ ) tumor cells were injected into NOD-SCID mice. Tumors were allowed to grow for 30 days until they reached an average volume of  $90 \text{ mm}^3$ . Tumor-bearing mice were randomized into 4 groups ( $n = 10$  per group). Tumor-bearing mice were treated with anti-RSPO3 antibody 131R002, gemcitabine plus nab-paclitaxel (ABRAXANE), a combination of anti-RSPO3 antibody and gemcitabine and nab-paclitaxel (ABRAXANE). Antibodies were dosed at 25 mg/kg once a week. Gemcitabine was dosed at 20 mg/kg once a week and nab-paclitaxel (ABRAXANE) was dosed at 30 mg/kg once a week. Tumor growth was monitored and tumor volumes were measured with electronic calipers on the indicated days post-treatment.

[0346] In Figure 12A the results from all four treatment groups are shown and in Figure 12B only the combination treatments are shown on an expanded scale. Figure 12A and 12B show that anti-RSPO3 antibody in combination with gemcitabine and nab-paclitaxel (ABRAXANE) inhibited OMP-PN35 pancreatic tumor growth better than gemcitabine and nab-paclitaxel (ABRAXANE) alone.

#### Example 13

Inhibition of  $\beta$ -catenin activity by anti-RSPO3 antibodies

[0347] HEK-293 cells were transfected with a 6xTCF-luciferase reporter vector (TOPflash, Millipore, Billerica, MA). After 24-48 hrs, the transfected HEK-293 cells were incubated with a combination of WNT3a conditioned medium (5ng/ml) and human RSPO3 (2ng/ml, R&D BioSystems) in the presence of anti-RSPO3 antibodies h131R007 or h131R010. Antibodies h131R007 or h131R010 were added to the cells in 5-fold serial dilutions from 20µg/ml to 0.0064µg/ml. As controls, cells were incubated with WNT3a conditioned medium alone, a combination of WNT3a conditioned medium and human RSPO3, or with no addition to cells. The cells were incubated for 16 hours and luciferase activity was measured using Steady-Glo® Luciferase Assay System according to the manufacturer's instructions (Promega, Madison, WI).

[0348] As shown in Figure 13, antibody h131R010 reduced WNT3a/RSPO3-induced β-catenin signaling in a dose-dependent manner and to a similar extent as h131R007. Since h131R010 inhibited β-catenin signaling to the same extent as h131R007, it is clear that activity of the anti-RSPO3 antibody was not affected by conversion to an IgG1 isotype.

#### Example 14

Inhibition of lung tumor growth *in vivo* by anti-RSPO3 antibodies

[0349] OMP-LU25 is a patient-derived non small cell lung cancer (NSCLC) xenograft that was selected for testing based on high level of RPSO3 expression in microarray data. OMP-LU25 tumor cells ( $5 \times 10^4$ ) were injected into NOD-SCID mice. Tumors were allowed to grow for 33 days until they reached an average volume of 120 mm<sup>3</sup>. Tumor-bearing mice were randomized into 4 groups (n = 9 per group). Tumor-bearing mice were treated with either control antibody, anti-RSPO3 antibody 131R008, paclitaxel, or the combination of anti-RSPO3 antibody 131R008 and paclitaxel. Antibodies were dosed weekly at 20mg/kg. Paclitaxel was dosed weekly at 15mg/kg. Tumor growth was monitored and tumor volumes were measured with electronic calipers on the indicated days post-treatment.

[0350] As shown in Figure 14, anti-RSPO3 antibody 131R008 inhibited OMP-LU25 tumor growth as a single agent and in combination with chemotherapy. Furthermore, the combination of anti-RSPO3 antibody 131R008 with paclitaxel led to tumor regression.

[0351] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to person skilled in the art and are to be included within the spirit and purview of this application.

[0352] All publications, patents, patent applications, internet sites, and accession numbers/database sequences including both polynucleotide and polypeptide sequences cited herein are hereby incorporated by reference herein in their entirety for all purposes to the same extent as if each individual publication,

patent, patent application, internet site, or accession number/database sequence were specifically and individually indicated to be so incorporated by reference.

[0353] The sequences disclosed in the application are:

**Human RSPO1 protein sequence with signal sequence (SEQ ID NO:1)**

MRLGLCVVALVLSWTHLTISSRGIKGKRRQRRI SAEGSQACAKGCELCSEVNGCLKCSPKL  
FILLERNDIRQVGVCPLPSCPPGYFDARNPDMNKCIKCKIEHCEACFSHNFCTKCKEGLYL  
HKGRCPACPEGSSAANGTMECSSPAQCEMSEWSPWGPCSKKQQLCGFRRGSEERTRRVL  
HAPVGDHAACSDTKETRRC TVRRVPCPEGQKRRKGGQGRRENANRNLARKESKEAGAGSR  
RRKGQQQQQQGTVGPLTSAGPA

**Human RSPO2 protein sequence with signal sequence (SEQ ID NO:2)**

MQFRLFSFALIILNCMDYSHCQGNRWRRSKRASYVSNPICKGCLSCSKDNGCSRQQQKLF  
FFLRREGMRQYGECLHSCPSGYYGHRAPDMNRCARCRTE NCDS CF SKDFCTKCKVGFYLH  
RGRCFDECPDGFAPLEETMECEVGEVGHWSEWGTC SRNNRTCGFKWGLETRTRQIVKKP  
VKDTIPCPPTIAESRRCKMTMRHCPGGKRTPKAKEKRNKKKKRKLIERAQEQHSVFLATDR  
ANQ

**Human RSPO3 protein sequence with signal sequence (SEQ ID NO:3)**

MHLRLISWLFIIILNFMEYIGSQNASRGRRRQRMHPNVSQGCQGGCATCSDYNGCLSCCKPR  
LFFALERIGMKQIGVCLSSCPSGYYGTRYPDINKCTKCKADCDTCFNKNFCTKCKSGFYL  
HLGKCLDNCPEGLEANNHTMECVSIVHCEVSEWNPWSPCTKKGKTCGFKRGTTETRVREII  
QHPSAKGNLCPPTNETRKCTVQRKKCQKGERGKKGRERKRKKPNKGESKEAIPDSKSLES  
SKEIPEQRENKQQQKKRKVQDKQKSVSVSTVH

**Human RSPO4 protein sequence with signal sequence (SEQ ID NO:4)**

MRAPLCLLLLLVAHAVDMLALNRRKKQVGTGLGGNCTGCIICSEENGCTCQQLFLFIRR  
EGIRQYGKCLHDCPPGYFGIRGQEVNRCKKCGATCESCFSDFCIRCKRQFYLYKGKCLP  
TCPGTLAHQNTRECQGECELGPGWGWSPCTHNGKTCGSAWGLESRVREAGRAGHEEAAT  
CQVLSESRKCPIQRPCPGERSPGQKKGRKDRRPRKDRKLDRLDVRPRQPLQP

**Human RSPO3 protein sequence without predicted signal sequence (SEQ ID NO:5)**

QNASRGRRRQRMHPNVSQGCQGGCATCSDYNGCLSCCKPRLFFALERIGMKQIGVCLSSCP  
SGYYGTRYPDINKCTKCKADCDTCFNKNFCTKCKSGFYLHLGKCLDNCPEGLEANNHTME  
CVSIVHCEVSEWNPWSPCTKKGKTCGFKRGTTETRVREIIQHPSAKGNLCPPTNETRKCTV  
QRKKCQKGERGKKGRERKRKKPNKGESKEAIPDSKSLESSKEIPEQRENKQQQKKRKVQD  
KQKSVSVSTVH

**Human RSPO3 furin-like domain 1 (SEQ ID NO:6)**

PNVSQGCQGGCATCSDYNGCLSCCKPRLFFALERIGMKQIGVCLSSCPSGYYG

**Human RSPO3 furin-like domain 2 (SEQ ID NO:7)**

INKCTKCKADCDTCFNKNFCTKCKSGFYLHLGKCLDNCPEGLEA

**Human RSPO3 thrombospondin domain (SEQ ID NO:8)**

HCEVSEWNPWSPCTKKGKTCGFKRGTTETRVREIIQHPSAKGNLCPPTNETRKCTVQRKKCQ

**131R002/131R003 Heavy chain CDR1 (SEQ ID NO:9)**

KASGYTFTDYS

**131R002/131R003 Heavy chain CDR2 (SEQ ID NO:10)**

IYPSNGDS

131R002/131R003 Heavy chain CDR3 (SEQ ID NO:11)  
ATYFANYFDY

131R002/131R003 Light chain CDR1 (SEQ ID NO:12)  
QSVDDYDGDSYM

131R002/131R003 Light chain CDR2 (SEQ ID NO:13)  
AAS

131R002/131R003 Light chain CDR3 (SEQ ID NO:14)  
QQSNEPLT

131R002 Heavy chain variable region (SEQ ID NO:15)  
QVQLQESGPPELVKPGASVKISCKASGYTFDYSIHVVKQNHGKSLDWIGYIYPSNGDSGYN  
QKFKNRATLTVDTSSTAYLEVRRLTFEDSAVYYCATYFANYFDYWGQGTTLTVSSAST

131R003 Heavy chain variable region (SEQ ID NO:16)  
QVQLKQSGPELVKPGASVKISCKASGYTFDYSIHVVKQNHGKSLDWIGYIYPSNGDSGYN  
QKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANYFDYWGQGTTLTVSSAST

131R002/131R003 Light chain variable region (SEQ ID NO:17)  
DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGDSYMNWYQQKPGQPPKLLIYAASNLESG  
IPARFSGSGSGTDFTLNHPVEEEDAATYYCQQSNEPLTFGAGTKLELKR

131R002 Heavy chain variable region nucleotide sequence (SEQ ID NO:18)  
CAGGTACAATTGCAAGAATCCGGACCCGAACCTTGTGAAGCCCGGAGCGTCAGTCAAGATC  
TCGTGTAAGGCCAGCGGGTACACCTTTACGGATTATTCGATCCATTGGGTAAAACAGAAT  
CACGGGAAGTCGCTCGACTGGATTGGTTATATCTACCCGTCCAACGGTGATTTCGGGATAC  
AACCAGAAGTTCAAAAATCGGGCCACACTTACAGTGGACACATCGTCGTCAACTGCATAT  
CTCGAGGTCCGCAGACTGACGTTTGAGGACTCAGCTGTCTACTATTGCGCGACTTATTTTC  
GCCAACTACTTCGATTACTGGGGCCAGGGGACGACACTGACGGTCAGCTCCGCGAGCACC

131R003 Heavy chain variable region nucleotide sequence (SEQ ID NO:19)  
CAGGTGCAACTTAAACAGTCGGGGCCTGAGTTGGTCAAACCAGGAGCCTCAGTAAAGATT  
AGCTGCAAAGCATCAGGTTATACCTTTACGGATTACTCGATCCACTGGGTGAAGCAGAAC  
CACGGAAAGTCACTGGATTGGATCGGGTACATCTACCCCTCGAATGGAGATTTCGGGGTAT  
AACCAAAAGTTCAAAAACCGGGCCACGCTGACTGTGGACACGTCGTATTCCACCGCATAT  
TTGGAAGTCCGCAGACTCACGTTTCGAGGACTCCGCGGTATACTATTGTGCCACATACTTT  
GCGAATTACTTTGACTACTGGGGTCAGGGCACAACGCTTACTGTCTCCAGCGCGTCAACA

131R002/131R003 Light chain variable region nucleotide sequence (SEQ ID NO:20)  
GACATCGTGCTCACACAGAGCCCTGCATCGCTCGCAGTATCGCTTGGTCAGCGAGCGACC  
ATTTTCATGCAAAGCGTCACAATCGGTAGATTACGACGGAGACTCCTACATGAAGTGGTAT  
CAGCAGAAACCAGGGCAGCCCCCGAAGTTGCTCATCTACGCCGCGTCCAATCTGGAGTCA  
GGCATTCCCGCCAGATTTCAGCGGGAGCGGGTCAGGAACGGATTTTACCCTCAATATCCAT  
CCGGTAGAGGAGGAAGATGCGGCGACTTACTATTGTCTCAGCAGTCGAATGAGGACCCACTC  
ACGTTTCGGGGCTGGAACAAAACCTGGAACCTTAAACGG

131R002 Heavy chain amino acid sequence with predicted signal sequence underlined (SEQ ID NO:21)  
MKHLWFFLLLVAAPRVLSQVQLQESGPPELVKPGASVKISCKASGYTFDYSIHVVKQNH  
GKSLDWIGYIYPSNGDSGYNQKFKNRATLTVDTSSTAYLEVRRLTFEDSAVYYCATYFA

NYFDYWGGTTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
ALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCV  
ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVH  
NAKTKPREEQFNSTFRVSVLTVVHVDWLNQKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFF  
LYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

**131R003 Heavy chain amino acid sequence with predicted signal sequence underlined (SEQ ID NO:22)**

MKHLWFFLLLVAAPRWVLSQVQLKQSGPELVKPGASVKISCKASGYTFTDYSIHWVKQNH  
GKSLDWIGYIYPSNGDSGNQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFA  
NYFDYWGGTTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
ALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCV  
ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVH  
NAKTKPREEQFNSTFRVSVLTVVHVDWLNQKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFF  
LYSKLTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPGK

**131R002/131R003 Light chain amino acid sequence with predicted signal sequence underlined (SEQ ID NO:23)**

MKHLWFFLLLVAAPRWVLSDIVLTQSPASLAVSLGQRATISCKASQSVDDYDGDSYMNWYQ  
QKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQQSNEDPLT  
FGAGTKLELKRVAAPSVFI FPPSDEQLKSGTASVCLNNFYPREAKVQWKVDNALQSG  
NSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**131R002 Heavy chain nucleotide sequence with predicted signal sequence (SEQ ID NO:24)**

ATGAAACACTTGTGGTTCCTTTCTTTGCTGGTGGCAGCGCCTAGGTGGGTGCTCAGCCAG  
GTACAATTGCAAGAATCCGACCCGAACCTTGTGAAGCCCGGAGCGTCAGTCAAGATCTCG  
TGTAAGGCCAGCGGGTACACCTTTACGGATTATTCGATCCATTGGGTAAACAGAATCAC  
GGGAAGTCGCTCGACTGGATTGGTTATATCTACCCGTCCAACGGTGATTCGGGATACAAC  
CAGAAGTTCAAAAATCGGGCCACACTTACAGTGGACACATCGTCGTCAACTGCATATCTC  
GAGGTCCGCAGACTGACGTTTGAGGACTCAGCTGTCTACTATTGCGCGACTTATTTCCGC  
AACTACTTCGATTACTGGGGCCAGGGGACGACACTGACGGTCAGCTCCGCGAGCACCAAG  
GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCCCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGCTCCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGAC  
GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGACACGAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAATAACAAGACCACCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGA

**131R003 Heavy chain nucleotide sequence with predicted signal sequence (SEQ ID NO:25)**

ATGAAGCATCTTTGGTTCCTTCTGCTCTTGGTGGCTGCGCCGAGGTGGGTGCTCAGCCAG  
GTGCAACTTAAACAGTCGGGGCCTGAGTTGGTCAAACCAGGAGCCTCAGTAAAGATTAGC

TGCAAAGCATCAGGTTATACCTTTACGGATTACTCGATCCACTGGGTGAAGCAGAACCAC  
 GGAAAGTCACTGGATTGGATCGGGTACATCTACCCCTCGAATGGAGATTGGGGGTATAAC  
 CAAAAGTTCAAAAACCGGGCCACGCTGACTGTGGACACGTCGTATTCCACCGCATATTTG  
 GAAGTCCGCAGACTCACGTTTCGAGGACTCCGCGGTATACTATTGTGCCACATACTTTGCG  
 AATTACTTTGACTACTGGGGTCAGGGCACAACGCTTACTGTCTCCAGCGCGTCAACAAAG  
 GGCCCTCCGTGTTCCCTCTGGCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
 CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCTGGC  
 GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
 CTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
 GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGTGCGTG  
 GAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
 AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGAC  
 GTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
 AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
 CTGACCGTGGTGCACCAGGACTGGCTGAACGGC~~AAA~~GAATACAAGTGCAAGGTGTCCAAC  
 AAGGGCCTGCCTGCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
 CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
 ACCTGTCTGGTGAAGGGCTCTACCCTTCCGATATCGCCGTGGAGTGGAGTCTAACGGC  
 CAGCCTGAGAACAACCTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
 CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
 TCCGTGATGCACGAGGCCCTGCACAACCCTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
 GGCAAGTGA

**131R002/131R003 Light chain nucleotide sequence with predicted signal sequence (SEQ ID NO:26)**

ATGAAGCACCTCTGGTTCCTTCTTCTTCTGGTTCGAGCGCCGAGATGGGTACTTAGCGAC  
 ATCGTGTCTACACAGAGCCCTGCATCGCTCGCAGTATCGCTTGGTCAGCGAGCGACCAT  
 TCATGCAAAGCGTCACAATCGGTAGATTACGACGGAGACTCCTACATGAACTGGTATCAG  
 CAGAAACCAGGGCAGCCCCCGAAGTTGCTCATCTACGCCGCGTCCAATCTGGAGTCAGGC  
 ATTCCCGCCAGATTCAGCGGGAGCGGGTCAGGAACGGATTTTACCCTCAATATCCATCCG  
 GTAGAGGAGGAAGATGCGGCGACTTACTATTGTGTCAGCAGTCAATGAGGACCCACTCACG  
 TTCGGGGCTGGAACAAACTGGAACCTAAACGGACTGTGGCGGCTCCCTCAGTGTTCATC  
 TTCCCTCCCTCCGACGAACAATTGAAGTCGGGTACTGCCTCCGTGCTCTGTTTGTGAAC  
 AACTTTTATCCGAGGGAAGCCAAGGTGCAGTGAAGGTGGATAATGCGCTGCAGAGCGGT  
 AACTCGCAAGAGTCAGTCACAGAGCAAGACTCGAAGGATTGACGTATTCGCTCAGCAGC  
 ACATTGACGCTGTGCAAGGCAGATTACGAGAAACACAAGGTGTACGCGTGCAGAGGTACC  
 CATCAGGGATTGTGCTCACCCGTGACGAAATCCTTTAACC GCGGAGAATGCTGA

**131R002 Heavy chain amino acid sequence without predicted signal sequence (SEQ ID NO:27)**

QVQLQESGPPELVKPGASVKISCKASGYTFDYSIHVVKQNHGKSLDWIGYIYPSNGDSGY  
 NQKFKNRATLTVDTSSTAYLEVRRLTFEDSAVYYCATYFANYFDYWGGTTLTVSSAST  
 KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY  
 SLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCVECPPCPAPPVAGPSVFLFP  
 PKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
 VLTVVHQLDNLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS  
 LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGSAFLYSLKLTVDKSRWQQGNVFS  
 CSVMHEALHNHYTQKSLSLSPGK

**131R003 Heavy chain amino acid sequence without predicted signal sequence (SEQ ID NO:28)**

QVQLKQSGPELVKPGASVKISCKASGYTFDYSIHVVKQNHGKSLDWIGYIYPSNGDSGY  
 NQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANYFDYWGGTTLTVSSAST  
 KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY  
 SLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCVECPPCPAPPVAGPSVFLFP  
 PKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
 VLTVVHQLDNLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS

LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**131R002/131R003 Light chain amino acid sequence without predicted signal sequence (SEQ ID NO:29)**

DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGDSYMNWYQQKPGQPPKLLIYAASNLES  
GIPARFSGSGSGTDFTLNHPVEEEDAATYYCQQSNEDPLTFGAGTKLELKRTVAAPSVF  
IFPPSDEQLKSGTASVVCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS  
STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**131R002 Heavy chain amino acid sequence without predicted signal sequence (SEQ ID NO:30)**

CAGGTACAATTGCAAGAATCCGGACCCGAACCTTGTGAAGCCCGGAGCGTCAGTCAAGATC  
TCGTGTAAGGCCAGCGGGTACACCTTTACGGATTATTTCGATCCATTGGGTAAAACAGAAT  
CACGGGAAGTCGCTCGACTGGATTGGTTATATCTACCCGTCCAACGGTGATTTCGGGATAC  
AACCAGAAGTTCAAAAATCGGGCCACACTTACAGTGGACACATCGTCGTCAACTGCATAT  
CTCGAGGTCCGCAGACTGACGTTTGAGGACTCAGCTGTCTACTATTGCGCGACTTATTTTC  
GCCAACTACTTTCGATTACTGGGGCCAGGGGACGACACTGACGGTCAGCTCCGCGAGCACC  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCCTCCGGCCTGTAC  
TCCCTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGC  
AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
GTGGAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCCCTGTTCCCT  
CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTG  
GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
CACAACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT  
GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
AACAAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCC  
CTGACCTGTCTGGTGAAGGGCTTCTACCCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
GGCCAGCCTGAGAACAACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
TTCCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
CCTGGCAAGTGA

**131R003 Heavy chain amino acid sequence without predicted signal sequence (SEQ ID NO:31)**

CAGGTGCAACTTAAACAGTCGGGGCCCTGAGTTGGTCAAACCAGGAGCCTCAGTAAAGATT  
AGCTGCAAAGCATCAGGTTATACCTTTACGGATTACTCGATCCACTGGGTGAAGCAGAAC  
CACGGAAAGTCACTGGATTGGATCGGGTACATCTACCCCTCGAATGGAGATTTCGGGGTAT  
AACCAAAAGTTCAAAAACCGGGCCACGCTGACTGTGGACACGTCGTATTCCACCGCATAT  
TTGGAAGTCCGCAGACTCACGTTGAGGACTCCGCGGTATACTATTGTGCCACATACTTT  
GCGAATTACTTTGACTACTGGGGTCAGGGCACAACGCTTACTGTCTCCAGCGCGTCAACA  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCCTCCGGCCTGTAC  
TCCCTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGC  
AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
GTGGAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCCCTGTTCCCT  
CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTG  
GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
CACAACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT  
GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
AACAAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCC

CTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
GGCCAGCCTGAGAACAACACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
TTCCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
CCTGGCAAGTGA

**131R002/131R003 Light chain amino acid sequence without predicted signal sequence (SEQ ID NO:32)**

GACATCGTGCTCACACAGAGCCCTGCATCGCTCGCAGTATCGCTTGGTCAGCGAGCGACC  
ATTTTCATGCAAAGCGTCACAATCGGTAGATTACGACGGAGACTCCTACATGAACTGGTAT  
CAGCAGAAACCAGGGCAGCCCCCGAAGTTGCTCATCTACGCCGCTCCAATCTGGAGTCA  
GGCATTCCCGCCAGATTTCAGCGGGAGCGGGTTCAGGAACGGATTTTACCCTCAATATCCAT  
CCGGTAGAGGAGGAAGATGCGGCGACTTACTATTGTGTCAGCAGTCAATGAGGACCCACTC  
ACGTTCCGGGGCTGGAACAAAACCTGGAACCTTAAACGGACTGTGGCGGCTCCCTCAGTGTTT  
ATCTTCCCTCCCTCCGACGAACAATTGAAGTCCGGTACTGCCTCCGTCGTCTGTTTGTG  
AACAACTTTTATCCGAGGGAAGCCAAGGTGCAGTGGAAAGTGGATAATGCGCTGCAGAGC  
GGTAACTCGCAAGAGTCAAGTACAGAGCAAGACTCGAAGGATTTCGACGTATTCGCTCAGC  
AGCACATTGACGCTGTCTGAAGGCAGATTACGAGAAACACAAGGTGTACGCGTGCAGAGTC  
ACCCATCAGGGATTGTCGTACCCCGTGACGAAATCCTTTAACCGCGGAGAATGCTGA

**FLAG Tag (SEQ ID NO:33)**

DYKDDDDK

**131R003 Heavy chain CDR1 variant (SEQ ID NO:34)**

KASGYTFTSYTF

**131R003 Heavy chain CDR3 variant (SEQ ID NO:35)**

ATYFANNFDY

**131R003 Heavy chain variable region - Variant 1 (SEQ ID NO:36)**

QVQLKQSGPELVKPGASVKISCKASGYTFTDYSIHVVKQNHGKSLDWIGYIYPSNGDSGY  
NQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANNFDYWQGTTTLTVSS

**131R003 Heavy chain variable region - Variant 2 (SEQ ID NO:37)**

QVQLKQSGPELVKPGASVKISCKASGYTFTSYTFHWVKQNHGKSLDWIGYIYPSNGDSGY  
NQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANNFDYWQGTTTLTVSS

**131R003 Heavy chain - Variant 1 with predicted signal sequence underlined (SEQ ID NO:38)**

MKHLWFFLLLVAAPRWVLSQVQLKQSGPELVKPGASVKISCKASGYTFTDYSIHVVKQNH  
GKSLDWIGYIYPSNGDSGYNQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFA  
NNFDYWQGTTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCV  
ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVH  
NAKTKPREEQFNSTFRVSVLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGSFF  
LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

**131R003 Heavy chain - Variant 1 without predicted signal sequence (SEQ ID NO:39)**

QVQLKQSGPELVKPGASVKISCKASGYTFTDYSIHVVKQNHGKSLDWIGYIYPSNGDSGY  
NQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANNFDYWQGTTTLTVSSAST  
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY  
SLSSVVTVPSSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLF  
PKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
VLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS



LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**131R003 Heavy chain - Variant 1 nucleic acid with predicted signal sequence (SEQ ID NO:40)**

ATGAAGCATCTTTGGTTCTTCCTGCTCTTGGTGGCTGCGCCGAGGTGGGTGCTCAGCCAG  
GTGCAACTTAAACAGTCGGGGCCTGAGTTGGTCAAACCAGGAGCCTCAGTAAAGATTAGC  
TGCAAAGCATCAGGTTATACCTTTACGGATTACTCGATCCACTGGGTGAAGCAGAACCAC  
GGAAAGTCACTGGATTGGATCGGGTACATCTACCCCTCGAATGGAGATTCGGGGTATAAC  
CAAAAGTTCAAAAACCGGGCCACGCTGACTGTGGACACGTCGTATTCCACCGCATATTTG  
GAAGTCCGCAGACTCACGTTTCGAGGACTCCGCGGTATACTATTGTGCCACATACTTTGCG  
AATAACTTTGACTACTGGGGTCAGGGCACAACGCTTACTGTCTCCAGCGCGTCAACAAAG  
GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCTCCGTGGTGACCGTGCCTTCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGGAC  
GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGACACGAGTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGA

**131R003 Heavy chain - Variant 2 with predicted signal sequence underlined (SEQ ID NO:41)**

MKHLWFFLLLVAAPRWLSQVQLKQSGPELVKPGASVKISCKASGYTFTSYTFHWVKQNH  
GKSLDWIGYIYPSNGDSGYNQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFA  
NNFDYWGGTTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
ALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVRKKCV  
ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVH  
NAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFF  
LYSKLTVDKSRWQQGNVFS

**131R003 Heavy chain - Variant 2 without predicted signal sequence (SEQ ID NO:42)**

QVQLKQSGPELVKPGASVKISCKASGYTFTSYTFHWVKQNHGKSLDWIGYIYPSNGDSGY  
NQKFKNRATLTVDTSYSTAYLEVRRLTFEDSAVYYCATYFANNFDYWGGTTTLTVSSAST  
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY  
SLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVRKKCVVECPPCAPPVAGPSVFLFP  
PKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
VLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS  
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**131R003 Heavy chain - Variant 2 nucleic acid with predicted signal sequence (SEQ ID NO:43)**

ATGAAGCATCTTTGGTTCTTCCTGCTCTTGGTGGCTGCGCCGAGGTGGGTGCTCAGCCAG  
GTGCAACTTAAACAGTCGGGGCCTGAGTTGGTCAAACCAGGAGCCTCAGTAAAGATTAGC  
TGCAAAGCATCAGGATACACCTTCACTAGCTATACATTCCACTGGGTGAAGCAGAACCAC

GGAAAGTCACTGGATTGGATCGGGTACATCTACCCCTCGAATGGAGATTCGGGGTATAAC  
 CAAAAGTTCAAAAACCGGGCCACGCTGACTGTGGACACGTCGTATTCCACCGCATATTTG  
 GAAGTCCGCGAGACTCACGTTTCGAGGACTCCGCGGTATACTATTGTGCCACATACTTTGCG  
 AATAACTTTGACTACTGGGGTCAGGGCACAACGCTTACTGTCTCCAGCGCGTCAACAAAG  
 GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
 CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCTGGC  
 GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
 CTGTCCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
 GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
 GAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
 AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGAC  
 GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
 AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
 CTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
 AAGGGCCTGCCTGCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
 CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
 ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
 CAGCCTGAGAACAACCTACAAGACCACCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
 CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
 TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
 GGCAAGTGA

### Humanized 131R003 Antibodies

Humanized 131R005/131R007/131R008/131R010/131R011 Heavy chain variable region (SEQ ID NO:44)

QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYSIHWVRQAPGQGLEWIGYIYPSNGDSGY  
 NQKFKNRVTMTRDTSTSTAYMELSRLRSED TAVYYCATYFANNFDYWGQGTTLTVSS

Humanized 131R006A Heavy chain variable region (SEQ ID NO:45)

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYTFHWVRQAPGQGLEWIGYIYPSNGDSGY  
 NQKFKNRVTMTRDTSTSTAYMELSRLRSED TAVYYCATYFANNFDYWGQGTTLTVSS

Humanized 131R005/131R007/131R011 Heavy chain (IgG2) with predicted signal sequence underlined (SEQ ID NO:46)

MKHLWFFLLLVAAPRWVLSQVQLVQSGAEVKKPGASVKVSCKASGYTFTDYSIHWVRQAP  
 GQGLEWIGYIYPSNGDSGYNQKFKNRVTMTRDTSTSTAYMELSRLRSED TAVYYCATYFA  
 NNFYDWGQGTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
 ALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCV  
 ECPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVH  
 NAKTKPREEQFNSTFRVSVLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
 PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFF  
 LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Humanized 131R006A Heavy chain with predicted signal sequence underlined (SEQ ID NO:47)

MKHLWFFLLLVAAPRWVLSQVQLVQSGAEVKKPGASVKVSCKASGYTFTSYTFHWVRQAP  
 GQGLEWIGYIYPSNGDSGYNQKFKNRVTMTRDTSTSTAYMELSRLRSED TAVYYCATYFA  
 NNFYDWGQGTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
 ALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCV  
 ECPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVH  
 NAKTKPREEQFNSTFRVSVLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
 PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFF  
 LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R005/131R007/131R011 Heavy chain (IgG2) without predicted signal sequence (SEQ ID NO:48)**

QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYSIHVVRQAPGQGLEWIGYIYPSNGDSGY  
NQKFKNRVTMTRDTSTSTAYMELSLRSEDYAVYYCATYFANNFDYWGQGTTLTVSSAST  
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLY  
SLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCVECPPEPPVAGPSVFLFP  
PKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
VLTQVHQLDNLGKEYKCKVSNKGLPAPIEKTIKTKGQPREPQVYTLPPSREEMTKNQVS  
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R006A Heavy chain without predicted signal sequence (SEQ ID NO:49)**

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYTFHWVRQAPGQGLEWIGYIYPSNGDSGY  
NQKFKNRVTMTRDTSTSTAYMELSLRSEDYAVYYCATYFANNFDYWGQGTTLTVSSAST  
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLY  
SLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTVKCCVECPPEPPVAGPSVFLFP  
PKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS  
VLTQVHQLDNLGKEYKCKVSNKGLPAPIEKTIKTKGQPREPQVYTLPPSREEMTKNQVS  
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGGSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R005/131R007 Heavy chain variable region nucleic acid (SEQ ID NO:50)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCGGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTAC  
ATGGAAGTCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCA

**Humanized 131R006A Heavy chain variable region nucleic acid (SEQ ID NO:51)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTAGCTATACATTCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCGGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTAC  
ATGGAAGTCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCA

**Humanized 131R005/131R007 Heavy chain nucleic acid with predicted signal sequence (SEQ ID NO:52)**

ATGAAGCATCTGTGGTTTTTTCCTCCTCCTTGTGCGCCGCTCCACGCTGGGTGCTTTCCCAA  
GTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTGAGC  
TGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCACCT  
GGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTACAAC  
CAAAAGTTCAAGAACCGGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTACATG  
GAACTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTCGCT  
AATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACCAAG  
GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAAGTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCCTCCGTGGTGACCGTGCCTTCCTCCAAGTTCGGCACCCAGACCTACACCTGCAAC

GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGTCCTGCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGGAC  
GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAACCTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGA

**Humanized 131R006A Heavy chain nucleic acid with predicted signal sequence (SEQ ID NO:53)**

ATGAAGCATCTGTGGTTTTTTCCTCCTCCTTGTCGCGCTCCACGCTGGGTGCTTTCCCAA  
GTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTGAGC  
TGCAAGGCTTCTGGATACACCTTCACTAGCTATACATTCCACTGGGTGAGACAGGCACCT  
GGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTACAAC  
CAAAAGTTCAAGAACCAGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTACATG  
GAACTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTCGCT  
AATAACTTCGACTATTGGGGGAGGGCACCACCCTGACTGTGAGCTCAGCCTCAACCAAG  
GGCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCCTCCGGCCTGTACTCC  
CTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGTCCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGGAC  
GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAACCTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGA

**Humanized 131R005/131R007 Heavy chain nucleic acid without predicted signal sequence (SEQ ID NO:54)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCAGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTAC  
ATGGAACCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAACTTCGACTATTGGGGGAGGGCACCACCCTGACTGTGAGCTCAGCCTCAACC  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCCTCCGGCCTGTAC  
TCCCTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGC  
AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
GTGGAGTGCCCTCCTTGTCCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCT

CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTG  
 GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
 CACAACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT  
 GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
 AACAAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
 GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCC  
 CTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
 GGCCAGCCTGAGAACAACACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
 TTCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
 TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
 CCTGGCAAGTGA

**Humanized 131R006A Heavy chain - nucleic acid without predicted signal sequence (SEQ ID NO:55)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
 AGCTGCAAGGCTTCTGGATACACCTTCACTAGCTATACATTCCACTGGGTGAGACAGGCA  
 CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
 AACCAAAAGTTCAAGAACCGGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTAC  
 ATGGAACCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
 GCTAATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACC  
 AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCC  
 GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
 GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTAC  
 TCCCTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTCGGCACCCAGACCTACACCTGC  
 AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
 GTGGAGTGGCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCCCT  
 CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTG  
 GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
 CACAACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT  
 GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
 AACAAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
 GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCC  
 CTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
 GGCCAGCCTGAGAACAACACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
 TTCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
 TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
 CCTGGCAAGTGA

**Human IgG1 Heavy chain constant region (SEQ ID NO:56)**

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTCTPPCPAPELLGG  
 PSVFLFPPKPKDITLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN  
 STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDE  
 LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRW  
 QQGNVVFSCSVMEALHNHYTQKSLSLSPGK

**Human IgG2 Heavy chain constant region (SEQ ID NO:57)**

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSNFGTQTYTCNVNHNKPSNTKVDKTVKCCVECPPCPAPPVAGPSVF  
 LFPPKPKDITLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFR  
 VVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKN  
 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGN  
 VFSCSVMEALHNHYTQKSLSLSPGK

**Human IgG3 Heavy chain constant region (SEQ ID NO:58)**

ASTKGPSVFPLAPCSRSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSSLGTQTYTCNVNHKPSNTKVDKRVELKTPLGDTHTCPRCPEPKSC  
 DTPPPCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCPAPELLGGPSVFLFPPKPKDT  
 LMISRTPEVTCVVVDVSHEDPEVQFKWYVDGVEVHNAKTKPREEQYNSTFRVSVLTVLH  
 QDWLNGKEYKCKVSNKALPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVK  
 GFYPSDIAVEWESSGQPENNYNTTPMLDSGDSFFLYSKLTVDKSRWQQGNIFSCSVME  
 ALHNRFTQKSLSLSPGK

**Human IgG4 Heavy chain constant region (SEQ ID NO:59)**

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSSLGTQTYTCNVNHNKPSNTKVDKRVESKYGPPCPSCPAPEFLGGPSV  
 FLFPPKPKDTLMISRTPEVTCVVVDVSDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY  
 RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTKAKAGQPREPQVYTLPPSQEEMTK  
 NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSRLTVDKSRWQEG  
 NVFSCSVMEALHNHYTQKSLSLSLGK

**Human IgG2 Heavy chain constant region (13A Chain variant) (SEQ ID NO:60)**

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSNFGTQTYTCNVNHNKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVF  
 LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFR  
 VVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREKMTKN  
 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLKSDGDSFFLYSKLTVDKSRWQQGN  
 VFSCSVMEALHNHYTQKSLSLSPGK

**Human IgG2 Heavy chain constant region (13B Chain variant) (SEQ ID NO:61)**

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
 GLYSLSSVVTVPSSNFGTQTYTCNVNHNKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVF  
 LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFR  
 VVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKN  
 QVSLTCLVEGFYPSDIAVEWESNGQPENNYKTTTPMLDSGDSFFLYSELTVDKSRWQQGN  
 VFSCSVMEALHNHYTQKSLSLSPGK

**Humanized 131R006B Heavy chain variable region (SEQ ID NO:62)**

QVQLVQSGAEVKKPGASVKVSCKASGYTFDYSIHWRQAPGQGLEWIGYIYPSNGDSGY  
 NQKFKNRVTMTVDTSYSTAYMELSRLRSED TAVYYCATYFANNFDYWGGTTTLTVSS

**Humanized 131R006B Heavy chain with predicted signal sequence underlined (SEQ ID NO:63)**

MKHLWFFLLLVAAPRWVLSQVQLVQSGAEVKKPGASVKVSCKASGYTFDYSIHWRQAP  
 GQGLEWIGYIYPSNGDSGYNQKFKNRVTMTVDTSYSTAYMELSRLRSED TAVYYCATYFA  
 NNFYDWGGTTTLTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSG  
 ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQTYTCNVNHNKPSNTKVDKTKVERKCCV  
 ECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVH  
 NAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPRE  
 PQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLDSGDSFF  
 LYSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK

**Humanized 131R006B Heavy chain without predicted signal sequence (SEQ ID NO:64)**

QVQLVQSGAEVKKPGASVKVSCKASGYTFDYSIHWRQAPGQGLEWIGYIYPSNGDSGY  
 NQKFKNRVTMTVDTSYSTAYMELSRLRSED TAVYYCATYFANNFDYWGGTTTLTVSSAST  
 KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY  
 SLSSVVTVPSSNFGTQTYTCNVNHNKPSNTKVDKTKVERKCCVECPPCPAPPVAGPSVFLFPP  
 KPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS

VLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS  
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPMLDSGDSFFLYSKLTVDKSRWQQGNVFS  
CSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R006B Heavy chain variable region nucleic acid (SEQ ID NO:65)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCGGGTGACTATGACCGTGGATACCTCATACTCTACTGCCTAC  
ATGGAACCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCA

**Humanized 131R006B Heavy chain nucleic acid with sequence signal (SEQ ID NO:66)**

ATGAAGCATCTGTGGTTTTTCTCCTCCTTGTGCGCCGCTCCACGCTGGGTGCTTTCCCAA  
GTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTGAGC  
TGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCACCT  
GGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTACAAC  
CAAAAGTTCAAGAACCGGGTGACTATGACCGTGGATACCTCATACTCTACTGCCTACATG  
GAACTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTCGCT  
AATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACCAAG  
GGCCCTCCGTGTTCCCTCTGGCCCTTGTCTCCCGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGTCTCTCTCTCTCTGGCTGGCCCTTCTGTGTTCTGTCTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGGAC  
GTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAACCTACAAGACCACCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGA

**Humanized 131R006B Heavy chain nucleic acid without predicted sequence signal (SEQ ID NO:67)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCGGGTGACTATGACCGTGGATACCTCATACTCTACTGCCTAC  
ATGGAACCTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACC  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCTTGTCTCCCGTCCACCTCTGAGTCTACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCTCCGGCCTGTAC  
TCCCTGTCTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGC  
AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
GTGGAGTGCCTCCTTGTCTCTCTCTCTCTGGCTGGCCCTTCTGTGTTCTGTCTCCCT  
CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGT  
GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
CACAACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT



GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
AACAAGGGCCTGCCTGCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAAAGACCAAGAACCAGGTGTCC  
CTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
GGCCAGCCTGAGAACAACACTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
TTCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
CCTGGCAAGTGA

**Humanized 131R008/131R010 Heavy chain (IgG1) with predicted signal sequence underlined (SEQ ID NO:68)**

MKHLWFFLLLVAAPRWVLSQVQLVQSGAEVKKPGASVKVSCKASGYTFTDYSIHWRQAP  
QGQLEWIGYIYPSNGDSGYNQKFKNRVTMTRDTSTSTAYMELSRLRSEDVAVYYCATYFA  
NNFDYWQGQTTTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG  
ALTSKVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCD  
KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG  
VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG  
QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD  
GSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R008/131R010 Heavy chain (IgG1) without predicted signal sequence (SEQ ID NO:69)**

QVQLVQSGAEVKKPGASVKVSCKASGYTFTDYSIHWRQAPQGQLEWIGYIYPSNGDSGY  
NQKFKNRVTMTRDTSTSTAYMELSRLRSEDVAVYYCATYFANNFDYWQGQTTTLTVSSAST  
KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSKVHTFPAVLQSSGLY  
SLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSV  
FLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY  
RVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK  
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG  
NVFSCFSVMHEALHNHYTQKSLSLSPGK

**Humanized 131R008 Heavy chain (IgG1) with signal sequence nucleic acid (SEQ ID NO:70)**

ATGAAGCATCTGTGGTTTTTCTCCTCCTTGTCGCCGCTCCACGCTGGGTGCTTTCCCAA  
GTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTGAGC  
TGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCACCT  
GGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTACAAC  
CAAAAGTTCAAGAACCGGGTGACTATGACCAGAGATACCTCAACATCTACTGCCTACATG  
GAACTCAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTCGCT  
AATAACTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACCAAG  
GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTCCCTCCAAGTCCACCTCCGGCGGCACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCAGCCGTGCTGCAGTCCTCCGGCCTGTACTCC  
CTGTCTCCTCCGTGGTGACCGTGCTTCCCTCCTCCTGGGCACCCAGACCTACATCTGCAAC  
GTGAACCACAAGCCTTCCAACACCAAGGTGGACAAGCGGGTGGAGCCTAAGTCCTGCGAC  
AAGACCCACACCTGCCCTCCCTGCCCTGCCCCTGAGCTGCTGGGCGGACCTTCCGTGTTT  
CTGTTCCCTCCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAGGTGACCTGC  
GTGGTGGTGGACGTGTCCCACGAGGATCCTGAGGTGAAGTTCAATTGGTACGTGGACGGC  
GTGGAGGTGCACAACGCTAAGACCAAGCCAAGGGAGGAGCAGTACAACCTCCACCTACCGG  
GTGGTGTCTGTGCTGACCGTGTGCAACAGGACTGGCTGAACGGCAAAGAATACAAGTGC  
AAGGTCTCCAACAAGGCCCTGCCCGCTCCCATCGAGAAAACCATCTCCAAGGCCAAGGGC  
CAGCCTCGCGAGCCTCAGGTGTACACCCTGCCACCCAGCCGGGAGGAGATGACCAAGAAC  
CAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGG  
GAGTCTAACGGCCAGCCCGAGAACAACACTACAAGACCACCCCTCCTGTGCTGGACTCCGAC  
GGCTCCTTCTTCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGCAAC  
GTGTTCTCCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGAGCCTG



TCTCTGTCTCCTGGCAAGTGA

**Humanized 131R008 Heavy chain (IgG1) without predicted signal sequence nucleic acid (SEQ ID NO:71)**

CAAGTCCAATTGGTCCAGAGCGGTGCCGAAGTGAAGAAACCGGGAGCTTCCGTGAAAGTG  
AGCTGCAAGGCTTCTGGATACACCTTCACTGACTATTCAATCCACTGGGTGAGACAGGCA  
CCTGGTCAGGGACTGGAGTGGATTGGATACATCTACCCCTCAAATGGGGACTCTGGCTAC  
AACCAAAAGTTCAAGAACCGGGTGAATGACCAGAGATACCTCAACATCTACTGCCTAC  
ATGGAAGTCAAGCAGGCTGCGCTCAGAGGACACCGCAGTGTATTACTGTGCCACCTACTTC  
GCTAATAAATTTCGACTATTGGGGGCAGGGCACCACCCTGACTGTCAGCTCAGCCTCAACC  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTCCCTCCAAGTCCACCTCCGGCGGCACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCAGCCGTGCTGCAGTCTCCGGCCTGTAC  
TCCCTGTCTCTCCGTGGTGAACCGTGCCTTCCCTCCCTGGGCACCCAGACCTACATCTGC  
AACGTGAACCACAAGCCTTCCAACACCAAGGTGGACAAGCGGGTGGAGCCTAAGTCTCTGC  
GACAAGACCCACACCTGCCCTCCCTGCCCTGCCCTGAGCTGCTGGGCGGACCTTCCGTG  
TTCCTGTTCCCTCCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAGGTGACC  
TGCGTGGTGGTGGACGTGTCCACGAGGATCTGAGGTGAAGTTCAATTGGTACGTGGAC  
GGCGTGGAGGTGCACAACGCTAAGACCAAGCCAAGGGAGGAGCAGTACAACCTCCACCTAC  
CGGGTGGTGTCTGTGCTGACCGTGTGTCACCAGGACTGGCTGAACGGCAAAGAATACAAG  
TGCAAGGTCTCCAACAAGGCCCTGCCCGTCCCATCGAGAAAACCATCTCCAAGGCCAAG  
GGCCAGCCTCGCGAGCCTCAGGTGTACACCCTGCCACCCAGCCGGGAGGAGATGACCAAG  
AACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAG  
TGGGAGTCTAACGGCCAGCCCGAGAACAATAACAAGACCACCCCTCCTGTGCTGGACTCC  
GACGGCTCCTTCTCTCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGC  
AACGTGTTCTCTCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGAGC  
CTGTCTCTGTCTCTGGCAAGTGA

**Humanized 131R005/131R007/131R008 Light chain variable region (SEQ ID NO:72)**  
DIVLTQSPASLAVSLGQRATITCKASQSVDDYDGDSYMNWYQQKPGQPPKLLIYAASNLES  
GIPARFSGSGSGTDFTLTINPVEAEDVATYYCQQSNEDPLTFGAGTKLELKR

**Humanized 131R005/131R007/131R008 Light chain with predicted signal sequence underlined (SEQ ID NO:73)**

MKHLWFFLLLVAAPRVLSDIVLTQSPASLAVSLGQRATITCKASQSVDDYDGDSYMNWYQ  
QKPGQPPKLLIYAASNLES  
GIPARFSGSGSGTDFTLTINPVEAEDVATYYCQQSNEDPLTFGAGTKLELKR  
TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**Humanized 131R005/131R007/131R008 Light chain without predicted signal sequence underlined (SEQ ID NO:74)**

DIVLTQSPASLAVSLGQRATITCKASQSVDDYDGDSYMNWYQQKPGQPPKLLIYAASNLES  
GIPARFSGSGSGTDFTLTINPVEAEDVATYYCQQSNEDPLTFGAGTKLELKR  
TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS  
STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**Humanized 131R005/131R007/131R008 Light chain variable region nucleic acid (SEQ ID NO:75)**

GATATCGTCTCTGACCCAAAGCCCTGCTTCACTTGCTGTGAGCCTGGGGCAACGCGCCACC  
ATCACTTGCAAGGCATCTCAGAGCGTGGACTATGATGGAGACTCTTACATGAATTGGTAT  
CAACAGAAGCCAGGTCAACCTCCCAAAGTCTGATCTACGCCGCATCTAATCTTGAAAGC  
GGCATCCCGGCTCGGTTCTCTGGTTCTGGATCAGGAACCGACTTCACCCTCACCATTAAC  
CCAGTGGAGGCCGAGGACGTGGCTACTTACTACTGCCAGCAGTCAAACGAGGACCCCTG  
ACTTTCGGAGCCGGGACCAAGCTGGAGCTTAAGCGT

**Humanized 131R005/131R007/131R008 Light chain with signal sequence nucleic acid (SEQ ID NO:76)**

ATGAAACATCTTTGGTTCTTCCTTCTGCTGGTCGCTGCTCCTCGGTGGGTGCTTAGCGAT  
ATCGTCCTGACCCAAAGCCCTGCTTCACTTGCTGTGAGCCTGGGGCAACGCGCCACCATC  
ACTTGCAAGGCATCTCAGAGCGTGGACTATGATGGAGACTCTTACATGAATTGGTATCAA  
CAGAAGCCAGGTCAACCTCCCAAAGTCTGATCTACGCCGCATCTAATCTTGAAAGCGGC  
ATCCCGGCTCGGTTCTCTGGTTCTGGATCAGGAACCGACTTCACCCTCACCATTAAACCA  
GTGGAGGCCGAGGACGTGGCTACTTACTACTGCCAGCAGTCAAACGAGGACCCCTGACT  
TTCGGAGCCGGGACCAAGCTGGAGCTTAAGCGTACGGTGGCCGCACCGTCAGTCTTTATC  
TTTCCACCCTCCGACGAACAGCTTAAGTCAGGCACTGCCTCAGTCGTGTGTCTCCTCAAT  
AACTTCTACCCCAAGGAGGCCAAGGTGCAGTGGAAAGTGGACAACGCCCTCCAGTCCGGG  
AACTCTCAAGAAAGCGTCACCGAGCAGGACAGCAAGGACTCCACCTACTCACTGTCAAGC  
ACTCTCACCCTCTCAAAGGCCGATTATGAGAAGCACAAGGTGTACGCATGCGAAGTGACC  
CATCAGGGTCTGTCCTCTCCTGTACCAAGTCCTTCAATAGAGGAGAATGTTGA

**Humanized 131R005/131R007/131R008 Light chain without predicted signal sequence nucleic acid (SEQ ID NO:77)**

GATATCGTCCTGACCCAAAGCCCTGCTTCACTTGCTGTGAGCCTGGGGCAACGCGCCACC  
ATCACTTGCAAGGCATCTCAGAGCGTGGACTATGATGGAGACTCTTACATGAATTGGTAT  
CAACAGAAGCCAGGTCAACCTCCCAAAGTCTGATCTACGCCGCATCTAATCTTGAAAGC  
GGCATCCCGGCTCGGTTCTCTGGTTCTGGATCAGGAACCGACTTCACCCTCACCATTAAAC  
CCAGTGGAGGCCGAGGACGTGGCTACTTACTACTGCCAGCAGTCAAACGAGGACCCCTG  
ACTTTCGGAGCCGGGACCAAGCTGGAGCTTAAGCGTACGGTGGCCGCACCGTCAGTCTTT  
ATCTTTCACCCTCCGACGAACAGCTTAAGTCAGGCACTGCCTCAGTCGTGTGTCTCCTC  
AATAACTTCTACCCCAAGGAGGCCAAGGTGCAGTGGAAAGTGGACAACGCCCTCCAGTCC  
GGGA ACTCTCAAGAAAGCGTCACCGAGCAGGACAGCAAGGACTCCACCTACTCACTGTCA  
AGCACTCTCACCCTCTCAAAGGCCGATTATGAGAAGCACAAGGTGTACGCATGCGAAGTG  
ACCCATCAGGGTCTGTCCTCTCCTGTACCAAGTCCTTCAATAGAGGAGAATGTTGA

**Variant Heavy chain CDR1 (SEQ ID NO:78)**

DYSIH

**Variant Heavy chain CDR2 (SEQ ID NO:79)**

YIYPSNGDSGYNQKFK

**Variant Heavy chain CDR3 (SEQ ID NO:80)**

TYFANNFD

**Variant Light chain CDR1 (SEQ ID NO:81)**

KASQSVDYDGD SYMN

**Variant Light chain CDR2 (SEQ ID NO:82)**

AASNLES

**Variant Light chain CDR3 (SEQ ID NO:83)**

QQSNEDPLTF

**Humanized 131R010 Heavy chain (IgG1) with signal sequence nucleic acid (SEQ ID NO:84)**

ATGAAACACTTGTGGTTCTTTCTGCTCCTTGTCGCAGCACCACGGTGGGTGCTGTGCGAA  
GTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTCTCA  
TGCAAGGCCAGCGGATACACTTTACCCGACTACTCCATCCATTGGGTGAGGCAGGCTCCG  
GGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTCGAACGGAGATTCGGGGTACAAT  
CAGAAGTTCAAGAACCGCGTGACCATGACTCGGGACACCTCAACTTCACGGCTTATATG

GAAGTGAAGCCGCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTTGCG  
AACAATTTTCGATTACTGGGGACAAGGAACACGCTCACTGTCAGCTCAGCCAGCACCAAG  
GGCCCTCCGTGTTCCCTCTGGCCCTTCCCTCCAAGTCCACCTCCGGCGGCACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCAGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCTCCGTGGTGACCGTGCCTTCCCTCCCTGGGCACCCAGACCTACATCTGCAAC  
GTGAACCACAAGCCTTCCAACACCAAGGTGGACAAGCGGGTGGAGCCTAAGTCTGCGAC  
AAGACCCACACCTGCCCTCCCTGCCCTGCCCTGAGCTGCTGGGCGGACCTTCCGTGTTT  
CTGTTCCCTCCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAGGTGACCTGC  
GTGGTGGTGGACGTGTCCCACGAGGATCCTGAGGTGAAGTTCAATTGGTACGTGGACGGC  
GTGGAGGTGCACAACGCTAAGACCAAGCCAAGGGAGGAGCAGTACAACCTCCACCTACCGG  
GTGGTGTCTGTGCTGACCGTGTGACCAAGGACTGGCTGAACGGCAAAGAATACAAGTGC  
AAGGTCTCCAACAAGGCCCTGCCCGCTCCCATCGAGAAAACCATCTCCAAGGCCAAGGGC  
CAGCCTCGCGAGCCTCAGGTGTACACCCTGCCACCCAGCCGGGAGGAGATGACCAAGAAC  
CAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCTCCGATATCGCCGTGGAGTGG  
GAGTCTAACGGCCAGCCCGAGAACAACCTACAAGACCACCCCTCCTGTGCTGGACTCCGAC  
GGCTCCTTCTTCCCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGCAAC  
GTGTTCTCCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGAGCCTG  
TCTCTGTCTCCTGGCAAGTGATAA

**Humanized 131R010 Heavy chain (IgG1) without signal sequence nucleic acid (SEQ ID NO:85)**

CAAGTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTC  
TCATGCAAGGCCAGCGGATACACTTTCACCGACTACTCCATCCATTGGGTGAGGCAGGCT  
CCGGGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTGCAACGGAGATTCCGGGTAC  
AATCAGAAGTTCAAGAACCAGCGTGACCATGACTCGGGACACCTCAACTTCCACGGCTTAT  
ATGGAACCTGAGCCGCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTT  
GCGAACAATTTTCGATTACTGGGGACAAGGAACACGCTCACTGTCAGCTCAGCCAGCACC  
AAGGGCCCCTCCGTGTTCCCTCTGGCCCTTCCCTCCAAGTCCACCTCCGGCGGCACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTGGAACCTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCAGCCGTGCTGCAGTCTCCGGCCTGTAC  
TCCCTGTCTCCTCCGTGGTGAACCGTGCCTTCCCTCCCTGGGCACCCAGACCTACATCTGC  
AACGTGAACCACAAGCCTTCCAACACCAAGGTGGACAAGCGGGTGGAGCCTAAGTCTCTGC  
GACAAGACCCACACCTGCCCTCCCTGCCCTGCCCTGAGCTGCTGGGCGGACCTTCCGTG  
TTCCTGTTCCCTCCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAGGTGACC  
TGCGTGGTGGTGGACGTGTCCCACGAGGATCCTGAGGTGAAGTTCAATTGGTACGTGGAC  
GGCGTGGAGGTGCACAACGCTAAGACCAAGCCAAGGGAGGAGCAGTACAACCTCCACCTAC  
CGGTGGTGTCTGTGCTGACCGTGTGACCAAGGACTGGCTGAACGGCAAAGAATACAAG  
TGCAAGGTCTCCAACAAGGCCCTGCCCGCTCCCATCGAGAAAACCATCTCCAAGGCCAAG  
GGCCAGCCTCGCGAGCCTCAGGTGTACACCCTGCCACCCAGCCGGGAGGAGATGACCAAG  
AACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCTCCGATATCGCCGTGGAG  
TGGGAGTCTAACGGCCAGCCCGAGAACAACCTACAAGACCACCCCTCCTGTGCTGGACTCC  
GACGGCTCCTTCTTCCCTGTACTCCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCAGGGC  
AACGTGTTCTCCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGAGC  
CTGTCTCTGTCTCCTGGCAAGTGATAA

**Humanized 131R010/131R011 Light chain variable region (SEQ ID NO:86)**

DIQMTQSPSSLSASVGDRVTITCKASQSVSDYDGDSYMNWYQQKPKAPKLLIYAASNLES  
GVPSRFSGSGSGTDFTLTISPVAEDFATYYCQQSNEDPLTFGAGTKLELKR

**Humanized 131R010/131R011 Light chain with predicted signal sequence underlined (SEQ ID NO:87)**

MKHLWFFLLLVAAPRVLSDIQMTQSPSSLSASVGDRVTITCKASQSVSDYDGDSYMNWYQ  
QKPKAPKLLIYAASNLES GVPSRFSGSGSGTDFTLTISPVAEDFATYYCQQSNEDPLT  
FGAGTKLELKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG  
NSQESVTEQDSKDSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**Humanized 131R010/131R011 Light chain without predicted signal sequence (SEQ ID NO:88)**

DIQMTQSPSSLSASVGDRVTITCKASQSVDDYDGDSYMNWYQQKPGKAPKLLIYAASNLES  
GVPSRFSGSGSGTDFTLTISPVAEDFATYYCQQSNEDPLTFGAGTKLELKRITVAAPSVF  
IFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLS  
STLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

**Humanized 131R010/131R011 Light chain variable region nucleic acid (SEQ ID NO:89)**

GATATCCAGATGACTCAGTCGCCCTCATCGTTGAGCGCCTCGGTGCGGGATCGCGTGACT  
ATTACTTGTAAGCGTCCCAGAGCGTGGACTACGACGGAGATTCTTACATGAAGTGGTAT  
CAGCAAAAACCGGGAAAGGCTCCTAACTTCTCATCTACGCAGCCTCGAATCTGGAATCA  
GGAGTCCCAGAGCGGTTTACGCGGATCAGGCTCCGGTACTGATTTTACCCTCACGATCTCG  
CCAGTGCAAGCCGAGGACTTCGCGACCTACTACTGCCAACAGTCCAACGAGGACCCGCTG  
ACCTTCGGCGCAGGGACCAAGCTGGAAGTGAAGCGT

**Humanized 131R010/131R011 Light chain with signal sequence nucleic acid (SEQ ID NO:90)**

ATGAAACACCTGTGGTTCCTCCTGCTGGTGGCAGCTCCCAGATGGGTCTGTCCGAT  
ATCCAGATGACTCAGTCGCCCTCATCGTTGAGCGCCTCGGTGCGGGATCGCGTGACTATT  
ACTTGTAAGCGTCCCAGAGCGTGGACTACGACGGAGATTCTTACATGAAGTGGTATCAG  
CAAAAACCGGGAAAGGCTCCTAACTTCTCATCTACGCAGCCTCGAATCTGGAATCAGGA  
GTCCCAGAGCGGTTTACGCGGATCAGGCTCCGGTACTGATTTTACCCTCACGATCTCGCA  
GTGCAAGCCGAGGACTTCGCGACCTACTACTGCCAACAGTCCAACGAGGACCCGCTGACC  
TTCGGCGCAGGGACCAAGCTGGAAGTGAAGCGTACGGTGGCCGCTCCATCCGTGTTTATC  
TTTCCGCCGTCCGATGAGCAGCTCAAGTCGGGCACTGCCAGCGTGGTCTGCCTGCTTAAC  
AATTTCTACCCTAGGGAAGCCAAGGTGCAGTGAAGGTGGATAACGCGCTCCAATCCGGT  
AACTCGCAAGAGAGCGTGACCGAACAGGACTCAAAGGACTCGACGTACAGCCTGTATCG  
ACCTTGACTCTCTCAAAGGCCGACTACGAAAAGCACAAGGTCTACGCGTGCGAAGTCACC  
CATCAGGGACTGTCTCGCCTGTGACCAAGAGCTTCAATCGCGGAGAGTGCTGA

**Humanized 131R010/131R011 Light chain without signal sequence nucleic acid (SEQ ID NO:91)**

GATATCCAGATGACTCAGTCGCCCTCATCGTTGAGCGCCTCGGTGCGGGATCGCGTGACT  
ATTACTTGTAAGCGTCCCAGAGCGTGGACTACGACGGAGATTCTTACATGAAGTGGTAT  
CAGCAAAAACCGGGAAAGGCTCCTAACTTCTCATCTACGCAGCCTCGAATCTGGAATCA  
GGAGTCCCAGAGCGGTTTACGCGGATCAGGCTCCGGTACTGATTTTACCCTCACGATCTCG  
CCAGTGCAAGCCGAGGACTTCGCGACCTACTACTGCCAACAGTCCAACGAGGACCCGCTG  
ACCTTCGGCGCAGGGACCAAGCTGGAAGTGAAGCGTACGGTGGCCGCTCCATCCGTGTTT  
ATCTTTCCGCCGTCCGATGAGCAGCTCAAGTCGGGCACTGCCAGCGTGGTCTGCCTGCTT  
AACAATTTCTACCCTAGGGAAGCCAAGGTGCAGTGAAGGTGGATAACGCGCTCCAATCC  
GGTAAGTTCGCAAGAGAGCGTGACCGAACAGGACTCAAAGGACTCGACGTACAGCCTGTCA  
TCGACCTTGACTCTCTCAAAGGCCGACTACGAAAAGCACAAGGTCTACGCGTGCGAAGTC  
ACCATCAGGGACTGTCTCGCCTGTGACCAAGAGCTTCAATCGCGGAGAGTGCTGA

**Humanized 131R011 Heavy chain variable region nucleic acid (SEQ ID NO:92)**

CAAGTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTC  
TCATGCAAGGCCAGCGGATACACTTTCACCGACTACTCCATCCATTGGGTGAGGCAGGCT  
CCGGGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTGCAACGGAGATTCGGGGTAC  
AATCAGAAGTTCAAGAACCAGCGTGACCATGACTCGGGACACCTCAACTTCCACGGCTTAT  
ATGGAAGTGCAGCCGCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTT  
GCGAACAATTTGATTACTGGGGACAAGGAACACGCTCACTGTCAGCTCA

**Humanized 131R011 Heavy chain (IgG2) with signal sequence nucleic acid (SEQ ID NO:93)**

ATGAAACACTTGTGGTTCCTTCTGCTCCTTGTGCGCAGCACACGGTGGGTGCTGTGCGAA  
GTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTCTCA

TGCAAGGCCAGCGGATACACTTTACCGACTACTCCATCCATTGGGTGAGGCAGGCTCCG  
GGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTCGAACGGAGATTCCGGGTACAAT  
CAGAAGTTCAAGAACCGCGTGACCATGACTCGGGACACCTCAACTTCCACGGCTTATATG  
GAACTGAGCCGCCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTTGCG  
AACAATTTTCGATTACTGGGGACAAGGAACCACGCTCACTGTCAGCTCAGCCAGCACCAAG  
GGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCCGCT  
CTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCTGGC  
GCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTACTCC  
CTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGCAAC  
GTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGCGTG  
GAGTGCCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCTCCT  
AAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTGGAC  
GTGTCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTGCAC  
AACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCTGTG  
CTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCCAAC  
AAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGCGAG  
CCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTG  
ACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAACGGC  
CAGCCTGAGAACAACCTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTCTTC  
CTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGC  
TCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCTCCT  
GGCAAGTGATAA

**Humanized 131R011 Heavy chain (IgG2) without signal sequence nucleic acid (SEQ ID NO:94)**

CAAGTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTC  
TCATGCAAGGCCAGCGGATACACTTTACCGACTACTCCATCCATTGGGTGAGGCAGGCT  
CCGGGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTCGAACGGAGATTCCGGGTAC  
AATCAGAAGTTCAAGAACCGCGTGACCATGACTCGGGACACCTCAACTTCCACGGCTTAT  
ATGGAAGTGAAGCCCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTT  
GCGAACAATTTTCGATTACTGGGGACAAGGAACCACGCTCACTGTCAGCTCAGCCAGCACC  
AAGGGCCCCCTCCGTGTTCCCTCTGGCCCCCTTGCTCCCGGTCCACCTCTGAGTCTACCGCC  
GCTCTGGGCTGCCTGGTGAAGGACTACTTCCCTGAGCCTGTGACCGTGTCTTGGAACTCT  
GGCGCCCTGACCTCTGGCGTGCACACCTTCCCTGCCGTGCTGCAGTCTCCGGCCTGTAC  
TCCCTGTCTCCGTGGTGACCGTGCCTTCCCTCCAACCTTCGGCACCCAGACCTACACCTGC  
AACGTGGACCACAAGCCTTCCAACACCAAGGTGGACAAGACCGTGGAGCGGAAGTGCTGC  
GTGGAGTGGCCTCCTTGTCTGCTCCTCCTGTGGCTGGCCCTTCTGTGTTCTGTTCCCT  
CCTAAGCCTAAGGACACCCTGATGATCTCCCGGACCCCTGAAGTGACCTGCGTGGTGGTG  
GACGTGTCCCACGAGGACCCTGAGGTGCAGTTCAATTGGTACGTGGACGGCGTGGAGGTG  
CACACGCCAAGACCAAGCCTCGGGAGGAACAGTTCAACTCCACCTTCCGGGTGGTGTCT  
GTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAATACAAGTGCAAGGTGTCC  
AACCAAGGGCCTGCCTGCCCCCTATCGAAAAGACCATCTCTAAGACCAAGGGCCAGCCTCGC  
GAGCCTCAGGTCTACACCCTGCCTCCTAGCCGGGAGGAAATGACCAAGAACCAGGTGTCC  
CTGACCTGTCTGGTGAAGGGCTTCTACCCTTCCGATATCGCCGTGGAGTGGGAGTCTAAC  
GGCCAGCCTGAGAACAACCTACAAGACCACCCCTCCTATGCTGGACTCCGACGGCTCCTTC  
TTCTGTACTCCAAGCTGACAGTGGACAAGTCCCGGTGGCAGCAGGGCAACGTGTTCTCC  
TGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGTCCCTGTCCCTGTCT  
CCTGGCAAGTGATAA

**Humanized 131R010 Heavy chain variable region (SEQ ID NO:95)**

CAAGTGCAATTGGTGCAGTCCGGAGCGGAAGTGAAGAAGCCTGGTGCCTCGGTCAAAGTC  
TCATGCAAGGCCAGCGGATACACTTTACCGACTACTCCATCCATTGGGTGAGGCAGGCT  
CCGGGCCAGGGCCTGGAGTGGATTGGGTACATCTACCCGTCGAACGGAGATTCCGGGTAC  
AATCAGAAGTTCAAGAACCGCGTGACCATGACTCGGGACACCTCAACTTCCACGGCTTAT  
ATGGAAGTGAAGCCCTGAGATCCGAGGACACTGCGGTGTACTACTGTGCCACCTACTTT

GCGAACAATTTGATTACTGGGGACAAGGAACCACGCTCACTGTCAGCTC

## WHAT IS CLAIMED IS:

1. An isolated antibody that specifically binds human R-spondin 3 (RSPO3), which comprises:  
(a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9), KASGYTFTSYTF (SEQ ID NO:34), or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10) or YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11), ATYFANNFDY (SEQ ID NO:35), or TYFANNFD (SEQ ID NO:80); and/or  
(b) a light chain CDR1 comprising QSVDYDGDSYM (SEQ ID NO:12) or KASQSVDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AAS (SEQ ID NO:13) or AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14) or QQSNEPLTF (SEQ ID NO:83).
2. The antibody of claim 1, which comprises:  
(a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or KASGYTFTSYTF (SEQ ID NO:34), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising ATYFANYFDY (SEQ ID NO:11) or ATYFANNFDY (SEQ ID NO:35), and/or  
(b) a light chain CDR1 comprising QSVDYDGDSYM (SEQ ID NO:12), a light chain CDR2 comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEPLT (SEQ ID NO:14).
3. The antibody of claim 1, which comprises:  
(a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising YIYPSNGDSGYNQKFK (SEQ ID NO:79), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80); and/or  
(b) a light chain CDR1 comprising KASQSVDYDGDSYMN (SEQ ID NO:81), a light chain CDR2 comprising AASNLES (SEQ ID NO:82), and a light chain CDR3 comprising QQSNEPLTF (SEQ ID NO:83).
4. The antibody of claim 1, which comprises:  
(a) a heavy chain CDR1 comprising KASGYTFTDYS (SEQ ID NO:9) or DYSIH (SEQ ID NO:78), a heavy chain CDR2 comprising IYPSNGDS (SEQ ID NO:10), and a heavy chain CDR3 comprising TYFANNFD (SEQ ID NO:80); and/or  
(b) a light chain CDR1 comprising QSVDYDGDSYM (SEQ ID NO:12), a light chain CDR2

comprising AAS (SEQ ID NO:13), and a light chain CDR3 comprising QQSNEEDPLT (SEQ ID NO:14).

5. An isolated antibody that specifically binds human RSPO3, which comprises:
  - (a) a heavy chain variable region having at least 90% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, OR SEQ ID NO:62; and/or
  - (b) a light chain variable region having at least 90% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.
6. The antibody of any one of claims 1-5, which comprises:
  - (a) a heavy chain variable region having at least 95% sequence identity to SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:44, SEQ ID NO:45, or SEQ ID NO:62; and/or
  - (b) a light chain variable region having at least 95% sequence identity to SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.
7. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:15; and
  - (b) a light chain variable region comprising SEQ ID NO:17 or SEQ ID NO:72.
8. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:16; and
  - (b) a light chain variable region comprising SEQ ID NO:17 or SEQ ID NO:72.
9. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:36; and
  - (b) a light chain variable region comprising SEQ ID NO:17 or SEQ ID NO:72.
10. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:37; and
  - (b) a light chain variable region comprising SEQ ID NO:17 or SEQ ID NO:72.



11. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:44; and
  - (b) a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.
12. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:45; and
  - (b) a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.
13. The antibody of claim 6, which comprises:
  - (a) a heavy chain variable region comprising SEQ ID NO:62; and
  - (b) a light chain variable region comprising SEQ ID NO:17, SEQ ID NO:72, or SEQ ID NO:86.
14. An isolated antibody that competes with the antibody according to any one of claims 1-13 for specific binding to RSPO3.
15. An isolated antibody that binds the same epitope on RSPO3 as the antibody according to any one of claims 1-13.
16. An isolated antibody that binds an epitope on RSPO3 that overlaps with the epitope on RSPO3 bound by the antibody according to any one of claims 1-13.
17. The antibody according to any one of claims 1-16, which is a recombinant antibody, a monoclonal antibody, a chimeric antibody, or a bispecific antibody.
18. The antibody according to any one of claims 1-17, which is a humanized antibody.
19. The antibody according to any one of claims 1-17, which is a human antibody.
20. The antibody according to any one of claims 1-19, which is an IgG1 antibody.
21. The antibody according to any one of claims 1-19, which is an IgG2 antibody.
22. The antibody according to any one of claims 1-19, which is an antibody fragment comprising an antigen binding site.

23. An antibody comprising:
  - (a) a heavy chain sequence of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:64, or SEQ ID NO:69; and
  - (b) a light chain sequence of SEQ ID NO:29, SEQ ID NO:74, or SEQ ID NO:88.
24. The antibody of claim 23, which comprises a heavy chain sequence of SEQ ID NO:48 and a light chain sequence of SEQ ID NO:74 or SEQ ID NO:88.
25. The antibody of claim 23, which comprises a heavy chain sequence of SEQ ID NO:64 and a light chain sequence of SEQ ID NO:74 or SEQ ID NO:88.
26. The antibody of claim 23, which comprises a heavy chain sequence of SEQ ID NO:69 and a light chain sequence of SEQ ID NO:74 or SEQ ID NO:88.
27. An antibody comprising the heavy chain variable region and the light chain variable region from an antibody selected from the group consisting of: 131R102, 131R103, a variant of 131R103, a humanized version of 131R103, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, and h131R011.
28. An antibody selected from the group consisting of: 131R102, 131R103, a variant of 131R103, a humanized version of 131R103, h131R006A, h131R006B, h131R005/131R007, h131R008, h131R010, and h131R011.
29. An antibody comprising the heavy chain variable region encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.
30. An antibody comprising the heavy chain encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.
31. An antibody comprising the light chain variable region encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.
32. An antibody comprising the light chain encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.

33. An antibody comprising the heavy chain variable region encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_ and the light chain variable region encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.
34. An antibody comprising the heavy chain encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_ and the light chain encoded by the plasmid deposited with ATCC as PTA-\_\_\_\_\_.
35. The antibody according to any one of claims 1-34, which inhibits binding of RSPO3 to at least one leucine-rich repeat containing G protein coupled receptor (LGR).
36. The antibody of claim 35, wherein the LGR is selected from the group consisting of LGR4, LGR5, and LGR6.
37. The antibody of claim 36, wherein the LGR is LGR5.
38. The antibody according to any one of claims 1-37, which inhibits RSPO3 signaling.
39. The antibody according to any one of claims 1-38, which inhibits activation of  $\beta$ -catenin.
40. The antibody according to any one of claims 1-39, which inhibits  $\beta$ -catenin signaling.
41. The antibody according to any one of claims 1-40, which inhibits tumor growth.
42. The antibody according to any one of claims 1-41, which induces expression of differentiation markers in a tumor.
43. The antibody according to any one of claims 1-42, which induces cells in a tumor to differentiate.
44. The antibody according to any one of claims 1-41, which reduces the frequency of cancer stem cells in a tumor.
45. A polypeptide comprising a sequence selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ

ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64  
SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID  
NO:86, SEQ ID NO:87, and SEQ ID NO:88.

46. The polypeptide of claim 45, which is an antibody.
47. A cell comprising or producing the antibody or polypeptide according to any one of claims 1-46.
48. An isolated polynucleotide molecule comprising a polynucleotide that encodes an antibody or polypeptide according to any one of claims 1-46.
49. An isolated polynucleotide molecule comprising a polynucleotide sequence selected from the group consisting of: SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, and SEQ ID NO:95.
50. A vector comprising the polynucleotide of claim 48 or claim 49.
51. A cell comprising the polynucleotide of claim 48 or claim 49 or the vector of claim 50.
52. A pharmaceutical composition comprising the antibody or polypeptide according to any one of claims 1-46 and a pharmaceutically acceptable carrier.
53. A method of inhibiting growth of a tumor, wherein the method comprises contacting the tumor with an effective amount of an antibody according to any of claims 1-44.
54. A method of inhibiting growth of a tumor in a subject, wherein the method comprises administering to the subject a therapeutically effective amount of an antibody according to any of claims 1-44.

55. A method of inducing differentiation of tumor cells in a subject, comprising administering to the subject a therapeutically effective amount of an antibody according to any one of claims 1-44.
56. A method of reducing the frequency of cancer stem cells in a tumor in a subject, comprising administering to the subject a therapeutically effective amount of an antibody according to any one of claims 1-44.
57. A method of inhibiting  $\beta$ -catenin signaling in a cell, comprising contacting the cell with an effective amount of an antibody according to any one of claims 1-44.
58. The method of claim 57, wherein the cell is a tumor cell.
59. The method according to any one of claims 53-56 or 58, wherein the tumor is selected from the group consisting of colorectal tumor, ovarian tumor, pancreatic tumor, lung tumor, liver tumor, breast tumor, kidney tumor, prostate tumor, gastrointestinal tumor, melanoma, cervical tumor, bladder tumor, glioblastoma, and head and neck tumor.
60. The method of claim 59, wherein the tumor is a pancreatic tumor.
61. The method of claim 59, wherein the tumor is a lung tumor.
62. The method of claim 59, wherein the tumor is a colorectal tumor.
63. The method of claim 59, wherein the tumor is an ovarian tumor.
64. A method of treating cancer in a subject, wherein the method comprises administering to the subject a therapeutically effective amount of an antibody according to any of claims 1-44.
65. The method of claim 64, wherein the cancer is selected from the group consisting of colorectal cancer, ovarian cancer, pancreatic cancer, lung cancer, liver cancer, breast cancer, kidney cancer, prostate cancer, gastrointestinal cancer, melanoma, cervical cancer, bladder cancer, glioblastoma, and head and neck cancer.
66. The method of claim 65, wherein the cancer is colorectal cancer.

67. The method of claim 65, wherein the cancer is pancreatic cancer.
68. The method of claim 65, wherein the cancer is lung cancer.
69. The method of claim 65, wherein the cancer is ovarian cancer.
70. The method according to any one of claims 53-56 or 58-69, wherein the tumor or the cancer expresses elevated levels of RSPO3 as compared to levels of RSPO3 in a reference sample or to a pre-determined level of RSPO3.
71. The method according to any one of claims 53-56 or 58-70, further comprising a step of determining the level of RSPO3 expression in the tumor or cancer.
72. The method of claim 71, wherein determining the level of RSPO3 expression is done prior to treatment or contact with the antibody.
73. The method of claim 71 or claim 72, wherein if the tumor or cancer has an elevated expression level of RSPO3, the antibody is:
  - (a) administered to the subject; or
  - (b) contacted with the tumor or tumor cell.
74. The method according to any one of claims 53-56 or 58-69, wherein the tumor or the cancer has a RSPO gene fusion.
75. The method according to any one of claims 53-56 or 58-69, further comprising a step of determining if the tumor or cancer has a RSPO gene fusion.
76. The method of claim 74 or claim 75, wherein the RSPO gene fusion is a RSPO2 gene fusion.
77. The method of claim 74 or claim 75, wherein the RSPO gene fusion is a RSPO3 gene fusion.
78. The method according to any one of claims 53-56 or 58-77, wherein the subject has had a tumor or a cancer removed.

79. A method of treating a disease in a subject wherein the disease is associated with activation of  $\beta$ -catenin, comprising administering a therapeutically effective amount of an antibody according to any one of claims 1-44.
80. The method according to any one of claims 53-56 or 58-79, which further comprises administering at least one additional therapeutic agent.
81. The method of claim 80, wherein the additional therapeutic agent is a chemotherapeutic agent.
82. The method of claim 80, wherein the additional therapeutic agent is an angiogenesis inhibitor.
83. The method of claim 80, wherein the additional therapeutic agent is an antibody.
84. The method of claim 83, wherein the antibody is an anti-RSPO1 antibody.
85. The method of claim 83, wherein the antibody is an anti-RSPO2 antibody.
86. The method according to any one of claims 53-56 or 58-85, wherein the subject is human.
87. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising: determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3.
88. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3.
89. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising: determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3, wherein if the tumor has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody.

90. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3,wherein if the tumor has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody.
91. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising: determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3.
92. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3.
93. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising: determining if the subject has a tumor that has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3, wherein if the tumor has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody.
94. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has an elevated expression level of RSPO3 as compared to a reference sample or a pre-determined level of RSPO3,wherein if the tumor has an elevated expression level of RSPO3 the subject is selected for treatment with the antibody.
95. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising: determining if the subject has a tumor that has a RSPO gene fusion.



96. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has a RSPO gene fusion.
97. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising: determining if the subject has a tumor that has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with the antibody.
98. A method of identifying a human subject for treatment with an antibody that specifically binds RSPO3 comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has a RSPO gene fusion,wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with the antibody.
99. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising: determining if the subject has a tumor that has a RSPO gene fusion.
100. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has a RSPO gene fusion.
101. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising: determining if the subject has a tumor that has a RSPO gene fusion, wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with the antibody.
102. A method of selecting a human subject for treatment with an antibody that specifically binds RSPO3, comprising:
  - (a) obtaining a tumor sample from the subject, and
  - (b) determining if the tumor has a RSPO gene fusion,wherein if the tumor has a RSPO gene fusion, then the subject is selected for treatment with the antibody.

103. The method of any one of claims 95-102, wherein the RSPO gene fusion is a RSPO2 gene fusion.
104. The method of any one of claims 95-102, wherein the RSPO gene fusion is a RSPO3 gene fusion.
105. The method of any one of claims 87-104, wherein the antibody is the antibody of any one of claims 1-44.
106. The method of any one of claims 87-105, wherein the tumor is selected from the group consisting of colorectal tumor, colon tumor, ovarian tumor, pancreatic tumor, lung tumor, liver tumor, breast tumor, kidney tumor, prostate tumor, gastrointestinal tumor, melanoma, cervical tumor, bladder tumor, glioblastoma, and head and neck tumor.
107. The method of claim 106, wherein the tumor is a lung tumor.
108. The method of claim 106, wherein the tumor is a pancreatic tumor.
109. The method of claim 106, wherein the tumor is an ovarian tumor.
110. The method any one of claims 87-94 or 105-109, wherein the expression level of RSPO3 is determined by a PCR-based assay, microarray analysis, or immunohistochemistry.
111. The method of any one of claims 95-109, wherein the RSPO gene fusion is determined by a PCR-based assay, microarray analysis, or nucleotide sequencing.
112. The method of claim 111, wherein the nucleotide sequencing is NextGen sequencing or whole-genome sequencing.
113. The method of claim 88, 90, 92, 94, 96, 98, 100, or 102-109, wherein the sample is a fresh tumor sample, a frozen tumor sample, or a formalin-fixed paraffin-embedded sample.

Figure 1A

RSPO1

ID	Original Source	Purity	Treatment	Tissue Type	Avg Sig	STDev	P	A	M	138	278	417	558
Colon Diseased	Colon	DISEASED	Colon DISE		4.85	0.14	0	21	0				
Colon Benign	Colon	BENIGN	Colon BENI		4.82	0.09	0	24	0				
Breast Normal	Breast	NORMAL	Breast NOR		11.25	13.18	5	17	0	81.88	1		
Breast Malignant	Breast	MALIGNANT	Breast MAL		4.86	0.87	1	105	0				
Breast Benign	Breast	BENIGN	Breast BEN		42.26	12.70	2	7	0	8	7		
Brain Benign	Brain	BENIGN	Brain BENI		4.82	0.05	0	16	0				
Brain Malignant	Brain	MALIGNANT	Brain MALI		4.82	0.18	0	23	0				
Liver Benign	Liver	BENIGN	Liver BENI		4.81	0.14	0	4	0				
Kidney Normal	Kidney	NORMAL	Cortex of k		4.83	0.72	0	61	0				
Kidney Malignant	Kidney	MALIGNANT	Kidney MAL		6.73	5.34	3	88	0	21.1			
Kidney Benign	Kidney	BENIGN	Kidney BEN		4.84	0.12	0	15	0				
Endometrium Malignant	Endometrium	MALIGNANT	Endometrium		11.78	22.09	5	52	0	86.1	1.1		
Endometrium Benign	Endometrium	BENIGN	Endometrium		16.08	33.46	3	7	0	86.1	1.1		
Colon Normal	Colon	NORMAL	Ascending c		4.82	0.39	0	74	0				
Colon Malignant	Colon	MALIGNANT	Colon MALI		4.83	0.65	1	140	0	81			
Ovary Malignant	Ovary	MALIGNANT	Ovary MALI		33.80	85.09	32	102	1	88.0	8.8	1.8	1
Ovary Normal	Ovary	NORMAL	Ovary NORM		5.81	1.28	0	7	0	81			
Lung Benign	Lung	NORMAL	Lung NORMA		5.35	0.85	1	63	0				
Lung Malignant	Lung	BENIGN	Ovary BENI		20.32	38.55	5	30	0	81	1.1		
Lung Benign	Lung	MALIGNANT	Lung MALIG		4.84	0.73	0	124	0	81			
Liver Diseased	Liver	BENIGN	Lung BENIG		4.80	0.08	0	5	0				
Liver Malignant	Liver	DISEASED	Liver DISE		4.89	0.22	0	22	0	81			
Liver Normal	Liver	MALIGNANT	Liver MALI		4.82	0.15	0	25	0				
Prostate Malignant	Prostate	NORMAL	Liver NORM		4.85	0.13	0	6	0				
Prostate Normal	Prostate	MALIGNANT	Prostate M		5.00	0.69	0	73	0	81			
Pancreas Normal	Pancreas	NORMAL	Prostate N		6.03	8.23	0	32	0	81			
Pancreas Benign	Pancreas	NORMAL	Pancreas N		4.89	0.11	0	13	0				
Pancreas Malignant	Pancreas	DISEASED	Prostate D		5.87	2.10	2	18	0	81			
Pancreas Benign	Pancreas	BENIGN	Pancreas B		4.83	0.13	0	5	0				
Pancreas Malignant	Pancreas	MALIGNANT	Pancreas M		4.79	0.11	0	66	0				







Figure 2




Figure 3




Figure 4






Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

FIG. 11A




Figure 11

FIG. 11B

Anti-RSPO3 antibody as single agent

Geneset	SIZE	p-val
OMP_DLL4_UP	88	0.00E+00
WEINBERG_ES_1	371	7.30E-03
OMP_NEWCSC_UP	66	8.32E-03
CURATED_TGFB	200	2.23E-02
CURATED_STEMCELL	280	2.48E-02

Carboplatin as single agent

Geneset	SIZE	p-val
ASSOU_ESC_DN	69	1.43E-03

Anti-RSPO3 antibody + Carboplatin

Geneset	SIZE	p-val
BATTLE_HU_PROLIFERATION	184	0.00E+00
TIAN_GBM_CD133_UP	83	0.00E+00
NEVINS_CSR	85	0.00E+00
OMP_CD201+_HIGH	240	0.00E+00
WONG_EMBRYONIC_STEM_CELL_CORE	329	0.00E+00
WEINBERG_PROLIFERATION	147	0.00E+00
WEINBERG_ES_1	371	0.00E+00
RICKMAN_TUMOR_DIFFERENTIATED_W	106	0.00E+00
WEINBERG_OCT4_TARGETS	286	1.33E-02
MILANO_GSI_RAT_DN	57	1.66E-02
WEINBERG_ES_2	35	2.34E-02
PN_CD201_LOGIT18	18	4.71E-02



Figure 12




Figure 13




Figure 14

